

78810_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 78810

Manuscript Type: ORIGINAL ARTICLE

Clinical and Translational Research

Molecular mechanisms of Baihedihuang Decoction as a treatment for breast cancer related anxiety: A network pharmacology and molecular docking study

breast cancer related anxiety

Zhong-Hui Li, guohua yang, fang wang

Abstract

BACKGROUND

The therapeutic effects of a combination of Chinese medicines called Baihedihuang Decoction have been clinically verified, although its molecular targets in breast cancer related anxiety remain unknown.

AIM

To explore the molecular mechanisms of Baihedihuang decoction for breast cancer related anxiety treatment.

METHODS

We used the Traditional Chinese Medicine Systems Pharmacology database to screen the active ingredients and potential targets of Baihedihuang Decoction, and constructed the "drug-ingredient-target" network map with the help of Cytoscape 3.8 software. Also, we used Online Mendelian Inheritance in Man, DrugBank and Gencards databases to collect the disease targets of breast cancer related anxiety, and used the STRING platform to construct the Protein-protein interaction network. STRING platform was for protein interaction analysis and Protein-protein interaction network construction; Metascape platform was for Gene Ontology and Kyoto Encyclopedia of Genes and Genomes pathway enrichment analysis of key targets; Molecular docking technology was used to verify the drug component/target disease network.

RESULTS

We screened 16 active ingredients of Baihedihuang Decoction for breast cancer related anxiety, with 113 target proteins. There are 931 disease targets of breast cancer related anxiety, and finally, 43 key targets and 305 Kyoto Encyclopedia of Genes and Genomes pathways were generated. The main active ingredients of Baihedihuang Decoction for breast cancer related anxiety are Verbascoside, β -sitosterol, Stigmasterol, catalpol, *etc.* CDK2, TP53, HTR2A, ESR1, *etc.* are its key targets, and the main involved signaling

pathways may include neuroactive ligand-receptor interaction pathway, 5-hydroxytryptaminergic synapse, P53 signaling pathway, cGMP-PKG signaling pathway, the cAMP signaling pathway, *etc.* Finally, molecular docking was performed by Vina software to validate the key active ingredients in Baihedihuang Decoction with the selected key action targets. The molecular docking results showed that Verbascoside, β -sitosterol, Stigmasterol and CDK2 could stably bind and interact through amino acid residues SER249, ARG260, PRO228, ALA282, SER276, LYS273, ASN272, *etc.*

CONCLUSION

Baihedihuang Decoction for breast cancer related anxiety may be multi-level, multi-target and multi-pathway, providing ideas and basis for further research.

Key Words: Network pharmacology; Molecular docking; Baihedihuang Decoction; breast cancer related anxiety; Mechanism of action

Li ZH, yang G, wang F. Molecular mechanisms of Baihedihuang Decoction as a treatment for breast cancer related anxiety: A network pharmacology and molecular docking study. *World J Clin Cases* 2022; In press

Core Tip: Baihedihuang decoction has significant effects on breast cancer related anxiety; however, its molecular targets in breast cancer related anxiety remain unknown. We conducted a network pharmacology and molecular docking study to determine whether Baihedihuang decoction ingredients target molecules and signaling pathways related to breast cancer related anxiety pathogenesis. The main active ingredients identified include Verbascoside, β -sitosterol, Stigmasterol, catalpol. Target genes included CDK2, TP53, HTR2A, ESR1, which are involved in cell cycle, apoptosis, cell autophagy, blood circulation and other biological processes. Our results confirm that Baihedihuang decoction can treat breast cancer related anxiety through

multi-component, multi-target, and multi-channel synergy.

INTRODUCTION

Breast cancer is one of the most common malignant tumors in American women and one of the main causes of cancer death. According to the statistics of the American Cancer Association, the prevalence and mortality of breast cancer are rising[1]. Although the treatment of breast cancer has made great progress due to the development of medicine, surgery, radiotherapy and chemotherapy for breast cancer still have a series of negative effects on women, affecting their mental state, reducing their quality of life, and causing irreparable traumatic experiences[2]. Some studies have shown that nearly half of the patients with malignant tumors have anxiety and depression tendencies [3]. The majority of patients with breast cancer are women, and their likelihood of anxiety and depression is higher than that of other tumors and patients with breast cancer not only have a relatively high incidence of anxiety and depression, but also have a high morbidity and mortality rate, and the morbidity and mortality rate of cancer patients with depression is 19% higher than that of cancer patients alone [4]. Clinical studies have shown that the incidence of depression in patients before breast cancer diagnosis is about 57%, and the incidence of depression after breast cancer surgery is about 63%[5], and surgical treatment can effectively improve the survival rate of patients, but due to the influences of physiological and psychological factors, postoperative patients are mostly accompanied by different degrees of depression and anxiety, which is not conducive to the implementation of postoperative treatment plans and disease recovery, with serious impacts on the survival quality of patients. This is harmful to the implementation of postoperative treatment plans and disease recovery, with serious impacts on patients' survival quality [6, 7]. The pathogenesis of breast cancer related anxiety(BCA) is not clear, but according to animal experiments, it may involve the neurological, endocrine and immune systems. Although there are many types of clinical drugs for the treatment of BCA, they have obvious adverse effects and poor patient compliance. In contrast, Chinese herbal

medicine is effective and safe in the supporting positive energy and alleviating the adverse effects of western medical treatment[8], and the role of Chinese herbal medicine compounds in the treatment of BCA deserves further study.

The clinical treatment of BCA with Baihedihuang Decoction (BD) is effective, but the related mechanism of action has been less studied. Network pharmacology is a powerful research method and it has been used in recent years to study the mechanism of action of the herbal compounding for diseases, and it also plays a non-negligible role in the studying the laws of the herbal compounding and guiding the development of new drugs[9]. In this paper, we will use network pharmacology to investigate the mechanism of anxiety effect of BD associated with breast cancer from various perspectives in order to lay the theoretical foundation for subsequent related research.

MATERIALS AND METHODS

Drug and gene data

The chemical composition of each herbal medicine in BD was searched in the Traditional Chinese Medicine Systems Pharmacology database (TCMSP)[10]. According to the principle of pharmacokinetics, the compounds that meet the conditions of Oral Bioavailability $\geq 30\%$ and Drug-Likeness ≥ 0.18 [11] should be screened, and the active ingredient-related protein targets should be found by the TCMSP. Since the chemical composition of Baihe and groundnut was altered accordingly after maceration, Regaloside B, caffeate, catalpol, and Verbascoside were added as active ingredients by searching the published relevant literature reports. Then, we standardized the protein target information in the Uniprot database (http://www.the_uniprot.org) and searched for their corresponding gene symbols. Finally, we used Cytoscape 3.8.1 software to construct a multi-level interaction of the herbal-compound-target network schematic.

Screening of breast cancer-related anxiety targets

By using "breast cancer", "anxiety" as keywords, we searched for targets in the Online Mendelian Inheritance in Man (OMIM). The disease-related targets in OMIM

(<http://www.omim.org>), DrugBank (<https://go.drugbank.com>) and Genecards (<https://www.genecards.org>) disease gene databases were searched, and the search results were finally combined and deleted. Duplicate values were finally obtained for breast cancer-related anxiety depression targets.

Protein interaction network construction and key target screening

the STRING11.0 Database (<https://www.string-db.org>) and Cytoscape3.8.1 software were used to construct a Protein interaction network (PPI) network map of BD-BCA targets, and then the active ingredient targets of BD and BCA were imported into Biso. The BCA targets were imported into BisoGenet, and the intersection network of the PPI networks was extracted by using the Merge function in Cytoscape, and the attribute values of each node in the intersection network were analyzed, and the results were filtered step by step to obtain the key targets.

Pathway enrichment analysis

The screened targets were imported into the Metascape platform (<https://metascape.org>) for Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis. GO mainly describes the specific roles and functions of the targets, including cellular component and molecular function, and biological GO mainly illustrates the specific actions and functions of the targets, including cellular component, molecular function and biological process; KEGG illustrates the signaling pathways of BD affecting BCA.

Molecular docking

The PDB database was searched for core genes to perform molecular docking experiments. The structures of the corresponding proteins were downloaded. PyMOL (version 2.4.0) software was used to remove water molecules and small ligand molecules from these receptor macromolecular structures. AutoDockTools (version 1.5.6) software was used to add polar hydrogen ions and set the grid box to determine the search range of the active pockets of the proteins, which were unknown. To include as many receptor structures as possible, the size of the active pocket had to be increased appropriately to improve the docking results. According to the core genes identified,

the drug components that interacted with the core genes were sought in the constructed drug-disease network. The two-dimensional structures of these small ligand molecules were downloaded from the PubChem database, and the MM2 calculation tool in ChemBio3D (version 14.0) software was used to optimize the three-dimensional structure of the small ligand molecule with the smallest free energy. Vina (version 1.1.2) software was used to perform the docking work.

RESULTS

A total of 160 the chemical components of BD were retrieved from the the TCMSP database. After the screening of Oral Bioavailability not less than 30% and Drug-Likeness not less than 0.18, a total of 9 active ingredients were obtained. Then, after reviewing the relevant literature for supplementation, 9 active ingredients were obtained for Baihe and 7 for Dihuang, Please see Table 1.

BD-BCA network construction

By using Cytoscape 3.8.1 software, the network diagram of 16 active ingredients of BD and their 113 targets after de-weighting was drawn, as shown in figure1.

BCA disease gene screening

By searching OMIM, Drugbank, and Genecards disease databases, 931 BCA targets were obtained after screening and removing duplicate values. The intersection of drug targets and disease targets was 42, as shown in figure 2.

The targets were submitted to the STRING11.0 database and Cytoscape3.8.1 software to obtain the PPI network map of BD for BCA target, as shown in Figure 3. The key targets were obtained after 2 screenings using BisoGenet to calculate the network topology characteristic attribute values of the intersection PPI, and the screening process is shown in Figure 4.

BD-BCA enrichment analysis

Gene enrichment analysis of the above 42 key nodes at the Metascape website included GO functional analysis, involving biological process, cellular component, and molecular function, and KEGG pathway analysis was performed, and the top 20 Log10 (P) values

were selected from the analysis results for visualization. From the results, it can be seen that the biological processes of BD are mainly the chemical synaptic transmission process, blood circulation process, cellular response process to organic cyclic compounds, and the regulation process of postsynaptic membrane potential, as shown in Figure 5. The functions of relevant targets for BCA are mainly enriched in the molecular functions of neurotransmitter receptor activity, G protein-coupled amine receptor activity, G protein-coupled adenosine receptor activity, and actions of relevant targets for BCA are mainly enriched in the molecular neuroactive ligand-receptor interaction pathway, 5-hydroxytryptaminergic synapses, P53 signaling pathway, cGMP-PKG signaling pathway, and cAMP signaling pathway, as shown in Figure 6.

Molecular docking simulation of active ingredients and target genes

CDK2 was selected as the core target in the PPI core network, and Stigmasterol, β -sitosterol and Verbascoside were selected as the key compounds based on the BD-BCA gene target map, and then the screened compounds and CDK2 were molecularly docked using AutoDock Vina software, and finally visualized using PyMOL software (Figure 7). The smaller the binding energy, the higher the binding activity and the easier the ligand binds to the receptor [12, 13], and the results are shown in Table 2. The results indicate that the core target CDK2 binds better to the key compounds, Stigmasterol, β -sitosterol and Verbascoside, and the predictions are relatively reliable.

DISCUSSION

Chinese medicine is an important component of complementary and alternative therapies and is more widely used in Asian countries. TCM has a diverse composition and is often useful in the treatment of various complex diseases through various biological pathways. While modern medicine has limited therapeutic effects and toxic side effects, studies have shown that TCM is more effective and safer than modern medicine in the treatment of BCA, and its use with modern therapies may also enhance its efficacy [14]. Therefore, we used network pharmacology and molecular docking to explore the mechanism of action of BD for BCA to help provide a safer and more effective treatment option for BCA patients.

BD is a summary of clinical experience over the ages and is the most essential part of the great treasury of Chinese medicine. The clinical efficacy of BD with addition and subtraction in the treatment of BCA is remarkable, but the mechanism of action is still unclear. Hu Huanping *et al* found that Baihe has a sedative-hypnotic effect, which shortens the sleep latency in animal models of insomnia [15]. Some experiments and related studies [16-18] have shown that BD can exert antidepressant effects through apoptosis and neurotransmitter regulation, and also has some antitumor effects.

In this paper, we screened the active ingredients of BD for the treatment of BCA through network pharmacology, mainly including Verbascoside, β -sitosterol, catalpol, *etc*. It has been shown that sitosterol has antidepressant effects and can act on the HTR2A target by inhibiting the hyperfunction of 5-HT_{2A}, thus inhibiting the phospholipase C system to restore the normal level of intracytoplasmic calcium ion concentration [19]. β -sitosterol has anti-inflammatory and anti-tumor effects [20]. Verbascoside can achieve antidepressant effects by stimulating the expression of 5-HT and DA genes and increasing the content of monoamine neurotransmitters in the brain [21]. Catalpol may inhibit the apoptosis of rat hippocampal astrocytes by promoting the expression of the Bcl-2 gene and thus inhibit depressive activity [22].

The key targets are CDK2, TP53, HTR2A, the ESR1, *etc*. The p53 protein encoded by the TP53 gene regulates cell division and proliferation, and it is one of its important roles and is also known as an "oncogene" [23]. HTR2A is a common target in the neuroactive ligand-receptor interaction signaling pathway and 5-hydroxytryptaminergic synapses. It has been shown [24] that inhibiting the function of 5-HT_{2A}, it can further inhibit the phospholipase C system, thus restoring the normal level of calcium ion concentration in the cytoplasm and producing antidepressant effects. CDK2 is closely related to tumor development, and by selectively inhibiting the activity of CDK2, the concentration of E2F increases, which in turn leads to cell cycle arrest in the S phase or apoptosis, thus achieving the therapeutic purpose of tumor treatment [25, 26]. It has also been shown that CDK2 is closely related to the growth of breast cancer cells and can promote

tamoxifen resistance in hormone receptor-positive breast cancer [27].The ER protein encoded by the ESR1 gene can regulate the metabolic level of estrogen by binding to the estrogen receptor, which is closely related to the level of female hormones and the development of breast cancer[28].

According to the KEGG pathway analysis, the main pathways involved in the treatment of BCA with BD include the 5-hydroxytryptaminergic synaptic pathway, P53 signaling pathway, cAMP signaling pathway, cGMP-PKG signaling pathway, *etc.* The 5-hydroxytryptaminergic synaptic pathway is composed of 5-hydroxytryptamine and various types of 5-hydroxytryptamine receptors on the presynaptic and postsynaptic membranes distributed in different parts of the brain, which can affect neurotransmitter releases, such as dopamine and γ -aminobutyric acid, and mediate higher brain activities such as learning, memory, and emotion, and its altered function can cause the occurrence of affective disorders such as anxiety disorders, depression, and Alzheimer's disease [29].P53 can regulate tumor development by participating in the regulation of cell signaling, induction of apoptosis, and autophagic processes. It has been shown that p53 transcriptionally up-regulates the expression of Bcl-2 family protein BAD when DNA is damaged[230],while p53 is carried by BAD to the outer mitochondrial membrane through interaction with BAD, which in turn increases the permeability of the outer mitochondrial membrane and induces apoptosis [31],which is corroborated with our derived key target TP53. Adenylate cyclase has a G-protein the S signaling coupling signaling function, which catalyzes the hydrolysis of adenosine triphosphate into cAMP, activates PKA in turn, and phosphorylates a variety of target proteins[32]. One study found that cAMP levels and PKA activity in the hippocampus of depressed rats were significantly lower than those in normal rats, and cAMP levels and PKA activity in CUMS depressed rats were significantly higher after administration of Baihezhiu Decoction [33].Therefore, the cAMP signaling pathway is closely related to the pathogenesis of depression. cGMP signaling pathway is involved in neuroplasticity, and its main downstream effector is protein kinase G(PKG) [34].In neuronal cells, the activation of PKG or increase in cGMP can prevent apoptosis, and the cGMP-PKG

signaling pathway plays an important anti-apoptotic role[35, 36].One study showed that hippocampal cGMP levels increased after 8 wk of fluoxetine and amitriptyline treatment, which in turn activated the downstream PKG-dominated cGMP signaling cascade response to exert antidepressant effects [34], which suggested that BD could act on the cGMP-PKG signaling pathway to exert antidepressant effects.

CONCLUSION

In the Summary, the active ingredients in BD, such as Verbascoside, β -sitosterol, Stigmasterol, mainly act on CDK2, TP53, HTR2A, ESR1 and other key targets, thus regulating the neuroactive ligand-receptor interaction pathway, 5-hydroxytryptaminergic synapse, P53 signaling pathway, cGMP-PKG signaling pathway, cAMP signaling pathway, etc, which are involved in cell cycle, apoptosis, cell autophagy, blood circulation and other biological processes. It has potential effects on cancer inhibition, anti-anxiety, etc. In addition, it has been demonstrated that the main active ingredients of BD can interact with the core targets, which indicates that it has the characteristics of multi-component, multi-target and multi-pathway co-regulation. The network pharmacology approach can help to interpret and predict the targets of the Chinese medicine compound more graphically and provide new ideas for the development of new drugs. However, network pharmacology is also limited by the available database information and experimental data, and the final results cannot reflect the whole real cell network without guaranteeing the complete authenticity of the database and experimental results [37]; To further study the treatment of BC with BD, relevant clinical studies will be carried out based on this network pharmacology study, etc, in order to obtain the stronger evidence-based medical evidence.

ARTICLE HIGHLIGHTS

Research background

The treatment effect of breast cancer related anxiety is not good and the side effects are

obvious. The clinical effect of Baihedihuang decoction is better, but its mechanism is unknown.

Research motivation

To explore the mechanism of Baihedihuang Decoction in treating breast cancer related anxiety.

Research objectives

To explore the relationship between drug targets of Baihedihuang Decoction and breast cancer related anxiety disorders.

Research methods

Network pharmacology and molecular docking technology were mainly used in this study.

Research results

We screened 16 active ingredients of Baihedihuang Decoction for breast cancer related anxiety, with 113 target proteins. Finally, molecular docking was performed by Vina software to validate the key active ingredients in Baihedihuang Decoction with the selected key action targets. The molecular docking results showed that Verbascoside , β -sitosterol , Stigmasterol and CDK2 could stably bind and interact through amino acid residues SER249, ARG260, PRO228, ALA282, SER276, LYS273, ASN272, etc.

1

Research conclusions

BD can the pathogenesis of BCA with multiple targets.

Research perspectives

The results of this study could serve as a fundamental basis for the further exploration in the *in vitro* and *in vivo* experiments for clinical promotions.

3%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

www.wjgnet.com
Internet

37 words — 1%
- 2

Yao Wang, Junbo Zou, Yanzhuo Jia, Yulin Liang, Xiaofei Zhang, Chang-Li Wang, Xiao Wang, Dongyan Guo, Yajun Shi, Ming Yang. "A Study on the Mechanism of Lavender in the Treatment of Insomnia Based on Network Pharmacology", Combinatorial Chemistry & High Throughput Screening, 2020
Crossref

14 words — < 1%
- 3

bmcbgenomics.biomedcentral.com
Internet

13 words — < 1%
- 4

Xiao Dong, Xiuxiu Fang, Yongyong Wang, Xiaohui Song. "Raman spectra and optical properties of the chalcogen-hyperdoped silicon: a first-principles study", Optics Express, 2018
Crossref

12 words — < 1%
- 5

Youyang Shi, Yuanyuan Wu, Feifei Li, Yang Zhang et al. "Identifying the anti-metastasis effect of Anhydroicaritin on breast cancer: Coupling network pharmacology with experimental validation", Journal of Ethnopharmacology, 2022
Crossref

12 words — < 1%

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 12 WORDS
EXCLUDE BIBLIOGRAPHY	OFF	EXCLUDE MATCHES	< 12 WORDS