

63663_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 63663

Manuscript Type: ORIGINAL ARTICLE

Observational Study

11

Prognostic values of intraorbital segment of optic nerve sheath diameter (ONSD) and ONSD/eyeball transverse diameter ratio for comatose patients with acute stroke : An observational study

Prognostic values of ONSD and ONSD/ ETD

Sha Zhu, Chao Chen, Liu-Liu Wang, Dian-Jiang Zhao, Yuan-Li Zhao, Xian-Zeng Liu

Abstract

BACKGROUND

Optic nerve sheath diameter (ONSD) measurement is one of the non-invasive methods recommended for increased intracranial pressure (ICP) monitoring.

AIM

This study aimed to evaluate the roles of optic nerve sheath diameter (ONSD) and ONSD/eyeball transverse diameter (ETD) ratio in predicting prognosis of death in comatose patients with acute stroke during their hospitalization.

METHODS

A total of 67 comatose patients with acute stroke were retrospectively recruited. The ONSD and ETD were measured by cranial computed tomography (CT) scan. All patients underwent cranial CT scan within 24 h after coma onset. Patients were divided into death group and survival group according to their survival status at discharge. The

differences of the ONSD and ONSD/ETD ratio between the two groups and their prognostic values were compared.

RESULTS

The ONSD and ONSD/ETD ratio were 6.07 ± 0.72 mm and 0.27 ± 0.03 in the comatose patients, respectively. The ONSD was significantly greater in the death group than that in the survival group (6.32 ± 0.67 vs. 5.65 ± 0.62 mm, $t = 4.078$, $P < 0.0001$). The ONSD/ETD ratio was significantly higher in the death group than that in the survival group (0.28 ± 0.03 vs. 0.25 ± 0.02 , $t = 4.625$, $P < 0.0001$). The area under the receiver operating characteristic (ROC) curve (AUC) was 0.760 (95% confidence interval (CI): 0.637-0.882, $P < 0.0001$) for the ONSD and 0.808 (95% CI: 0.696-0.920, $P < 0.0001$) for the ONSD/ETD ratio.

CONCLUSION

The mortality increased in comatose patients with acute stroke when the ONSD was >5.7 mm or the ONSD/ETD ratio was >0.25 . Both indexes could be used as prognostic tools for comatose patients with acute stroke. The ONSD/ETD ratio was more stable than the ONSD alone, which would be preferred in clinical practice.

Key Words: optic nerve sheath diameter, eyeball transverse diameter, coma, intracranial pressure, stroke, prognosis

Zhu S, Chen C, Wang LL, Zhao DJ, Zhao YL, Liu XZ. Prognostic values of intraorbital segment of optic nerve sheath diameter (ONSD) and ONSD/eyeball transverse diameter ratio for comatose patients with acute stroke : An observational study. *World J Clin Cases* 2022; In press

Core Tip: Optic nerve sheath diameter (ONSD) and ONSD/eyeball transverse diameter (ETD) were correlated with intracranial pressure and prognosis. This study

aimed to evaluate the roles of ONSD and ONSD/ETD ratio in predicting prognosis of death in comatose patients with acute stroke during hospitalization. 67 comatose patients were retrospectively recruited. ONSD and ETD were measured by cranial CT scans within 24 h after coma onset. It was found that the mortality increased when $\text{ONSD} > 5.7 \text{ mm}$ or $\text{ONSD/ETD ratio} > 0.25$. The ONSD/ETD ratio was more stable than ONSD alone, which be preferred in clinical practice.

Optic nerve sheath diameter (ONSD) and ONSD/eyeball transverse diameter(ETD) were correlated with intracranial pressure and prognosis. This study aimed to evaluate the roles of ONSD and ONSD/ETD ratio in predicting prognosis of death in comatose patients with acute stroke during hospitalization. 67 comatose patients were retrospectively recruited. ONSD and ETD were measured by cranial CT scans within 24 h after coma onset. It was found that the mortality increased when $\text{ONSD} > 5.7 \text{ mm}$ or $\text{ONSD/ETD ratio} > 0.25$. The ONSD/ETD ratio was more stable than ONSD alone, which be preferred in clinical practice.

INTRODUCTION

Stroke is a leading cause of death and disability worldwide¹. As the world's largest developing country, China had the largest number of incident cases and deaths related to stroke². Stroke is mainly accompanied by an increased intracranial pressure (ICP), leading to more severe brain damage *via* secondary cerebral ischemia and hernia³. Coma is the most severe disturbance of consciousness. A sudden rise of ICP is typically observed in comatose patients with acute stroke. Therefore, ICP monitoring is of particular importance for comatose patients.

In recent years, non-invasive ICP monitoring methods have been developed, such as transcranial Doppler ultrasound, cerebral blood flow velocity measurement, venous intraocular pressure (IOP) measurement, retinal optical coherence tomography, optic nerve sheath diameter (ONSD) measurement, *etc.* They provided an achievable ICP measurement method for specific patients, especially when invasive monitoring is contraindicated or unavailable. Among all the non-invasive ICP monitoring methods,

ONSD measurement is one of the most convenient and user-friendly methods for patients with severe neurological diseases.

Optic nerve sheath is a continuation of the dura mater and contains a subarachnoid space. It is continuous with the intracranial subarachnoid space, and an increased ICP can be directly transmitted to the optic nerve sheath^{4,5}. Previous studies have shown that computed tomography (CT) can be used to accurately measure ONSD. The strong correlation between ICP and ONSD has been generally identified⁶⁻⁸. Therefore, ONSD measurement is regarded as a reliable and noninvasive tool for ICP monitoring. However, the ONSD of normal and sick individuals mainly varies noticeably, indicating a large standard deviation (SD) range. In subsequent studies, it was found that the ONSD was strongly correlated with eyeball transverse diameter (ETD) in healthy individuals⁹. The ONSD/ETD ratio slightly varies, indicating a small SD range. Therefore, the ONSD/ETD ratio has a smaller variability and a higher stability, and it may be more appropriate for ICP monitoring^{7,10}. To date, several studies have shown that the ONSD/ETD ratio is more related to ICP than the ONSD^{6,7,10}. Therefore, it is reasonable to speculate that the ONSD/ETD ratio has a higher value in predicting the prognosis of neurological function. Previous studies have largely concentrated on the correlation between the ONSD and prognosis. It has been reported that the ONSD can help predict the prognosis of patients with cerebral hemorrhage (CH)¹¹. However, there are few studies on the prognostic role of the ONSD/ETD ratio in stroke.

Unenhanced cranial CT scan is the most common examination for acute stroke, especially for comatose patients, which can rapidly identify the type of stroke and determine the scope of associated structural changes. In the current study, unenhanced cranial CT scan was used to measure the ONSD and ETD, and the prognostic values of the ONSD and ONSD/ETD ratio in comatose patients with acute stroke were compared.

MATERIALS AND METHODS

2. Methods

2.1. Subjects

Comatose patients with acute stroke who were admitted to the neurological intensive care unit of Peking University International Hospital (Beijing, China) from August 2015 to September 2021 were retrospectively enrolled. The inclusion criteria were as follows: 1) patients who were aged ≥ 18 and ≤ 80 years old; 2) meeting the diagnostic criteria of acute cerebral infarction (ACI), CH, and spontaneous subarachnoid hemorrhage (SAH); 3) comatose patients with Glasgow coma scale (GCS) scores ≤ 8 upon admission. The exclusion criteria were as follows: 1) patients with a history of glaucoma, thyroid-associated ophthalmopathy, or optic neuropathy; 2) patients with acute brain stem and cerebellar strokes, and a history of SAH; 3) severe complications (e.g., hematological diseases and tumors) that might affect life expectancy. This retrospective study was approved by the Ethics Committee of Peking University International Hospital (Approval No. 2021-001 (BMR)).

2.2 Collection of baseline clinical data

Patients' baseline clinical data were collected, including age, gender, body weight, body mass index (BMI), mean arterial pressure (MAP), stroke type, GCS scores, surgery during hospitalization (e.g., clearance of hematoma and decompressive craniectomy), and the survival status at discharge (alive or dead).

2.3 ONSD and ETD measurements

CT scan was performed using a 64-row spiral CT scanner (Siemens, Munich, Germany), and the scanning parameters were as follows: tube voltage of 120 kV, tube current of 200-300 mA, slice thickness of 2 mm, slice interval of 3 mm, and pitch of 1. All patients received cranial CT scan within 24 h upon coma onset. The ONSD and ETD were measured by two experienced radiologists, who were blinded to patients' clinical data. The ONSD and ETD were measured at fixed mediastinal window setting (width, 300; level, 35). The direction of the optic nerve was determined by three-dimensional reconstruction of CT data. The ONSD was measured vertically at 3 mm behind the eyeball (Figure 1). ETD was measured from one side of the retina behind the lens to the other side for the maximum diameter (Figure 1). The values measured by two

radiologists were averaged. The ONSD and ETD were measured bilaterally, and their average was taken for each case to calculate the ONSD/ETD ratio.

Fig. 1 (A) The ONSD was measured at 3 mm behind the eyeball by the cranial CT scan; (B) ETD was measured by the cranial CT scan.

2.4 Statistics

The statistical analysis was performed using MedCalc 19.6 software (MedCalc Software Ltd., Ostend, Belgium). Continuous variables were expressed as mean \pm SD. Categorical variables were expressed as count (percentage). Death was set to 0 and survival to 1 depending on the patient's survival status at discharge. The χ^2 test was used for the statistical analysis of categorical variables, such as gender, GCS, and stroke type. The independent-samples t-test was applied for the statistical analysis of continuous variables, including age, body weight, BMI, ONSD, and ONSD/ETD ratio. The Mann-Whitney U test was utilized for the statistical analysis of height and MAP. The ONSD and ONSD/ETD ratio were included in the logistic regression model for multivariate analysis. The diagnostic performance of the ONSD and ONSD/ETD ratio was assessed by the receiver operating characteristic (ROC) curve analysis. All tests were two-sided, and $P < 0.05$ was considered statistically significant.

RESULTS

3.1 Participants' baseline data

A total of 67 comatose patients with acute stroke were included. The baseline characteristics of the comatose patients and healthy controls at baseline are summarized in Table 1. In the ACI group, 3 patients (23%) underwent decompressive craniectomy. In the CH group, 15 patients (38%) received hematoma evacuation. In the SAH group, 1 patient (7%) underwent ventricular puncture and drainage.

3.2 Comparison of clinical data between the survival group and the death group

In this study, 42 (62.7%) patients died, while 25 (37.3%) patients survived at discharge. As shown in Table 2, there were significant differences in the GCS score ($\chi^2 = 49.809$,

¹
 $P < 0.0001$), stroke type ($\chi^2 = 11.981$, $P = 0.003$), ONSD ($t = 4.078$, $P < 0.0001$), ONSD/ETD ratio ($t = 4.625$, $P < 0.0001$) and surgery ($\chi^2 = 4.803$, $P = 0.048$) between the two groups.

¹ 3.3 Evaluation of the ONSD and ONSD/ETD ratio for predicting prognosis of comatose patients

The ONSD and ONSD/ETD ratio were included in the logistic regression model for multivariate analysis, respectively. After adjusting for age, MAP, GCS score, stroke type, and surgery, the associations among the ONSD ($P = 0.04$), ONSD/ETD ratio ($P = 0.036$), and mortality were still significant. The performance of the ONSD and ONSD/ETD ratio in predicting prognosis of comatose patients with acute stroke is shown in Figure 2. The area under the curve (AUC) of the ONSD was 0.760 (95% confidence interval (CI): 0.637-0.882, $P < 0.0001$), with a sensitivity of 81.0% and a specificity of 64.0% at a cut-off value of 5.7 mm. The AUC of the ONSD/ETD ratio was 0.808 (95% CI: 0.696-0.920, $P < 0.0001$), with a sensitivity of 92.9% and a specificity of 68.0% at a cut-off value of 0.25. The AUC values of the ROC curves of the ONSD and ONSD/ETD ratio were compared. There was no significant difference in the AUC values of the two indices ($Z = 1.333$, $P = 0.1826$).

¹
Fig 2. ROC curves for comparing the efficiency of the ONSD and ONSD/ETD ratio in predicting prognosis of comatose patients with acute stroke.

DISCUSSION

In the present study, the ONSD measured in patients who died was 6.32 ± 0.67 mm, while that in patients who were alive was 5.65 ± 0.62 mm. The ONSD/ETD ratio was 0.28 ± 0.03 in the death group compared with 0.25 ± 0.02 in the survival group. These two parameters had almost the same predictive performance for prognosis of comatose patients with acute stroke.

A large number of studies on the ONSD for ICP assessment have been published since 1996. The ¹⁰ direct relationship between the increased ICP and ONSD in patients with traumatic and nontraumatic brain injuries was reported^{7,8,12}. However, the normal ranges and critical values of ICP and the ONSD for predicting the prognosis of such

patients have still remained elusive. Jenjitranan *et al* showed that the cut-off value of the ONSD for predicting the increased ICP on cranial CT was 4.8 mm¹³. Vaiman *et al* found that ONSD>5.5 mm on CT could predict an increased ICP in patients with hemorrhagic stroke⁶. A study on CH showed that the ONSD was 6.40±0.70 mm in patients with an increased ICP compared with 4.70 ±0.40 mm in those without an increased ICP, indicating a significant difference¹⁴. In another study, the ONSD was measured by ultrasound to predict early prognosis of patients with stroke. It was found that the ONSD in the death group and survival group was 5.5±0.4 and 4.4±0.5 mm, respectively¹⁵. The results of our study revealed that the ONSD in the death group was significantly greater than that in the survival group, demonstrating that the ONSD was an indirect monitoring marker for ICP. A higher mortality was predicted when the ONSD was >5.7 mm.

The clinical application of the ONSD has some limitations. First, factors affecting the ONSD are not clear. Second, the normal range or threshold of the ONSD for the diagnosis of ICP is uncertain, and the two ranges may overlap. Third, the reliability of the ONSD needs to be improved. Therefore, researchers have attempted to apply different indices to various imaging techniques, such as ONSD/ETD ratio.

The ONSD/ETD ratio was first described in 2014 and has recently attracted scholars' attention. To date, it has not been determined whether the ONSD could be related to age, sex, height, body weight, BMI, MAP, and head circumference. However, the ONSD/ETD ratio is irrelevant to the abovementioned factors based on previous studies^{16,17}. In addition, the ONSD/ETD ratio has a lower variability than the ONSD alone. Overall, the ONSD/ETD ratio is more stable than the ONSD. In healthy adults, the ONSD/ETD ratio on MRI was 0.22±0.01¹⁶, while that was 0.19 ± 0.02 on CT¹⁸. The ONSD/ETD ratio is a better predictor of increased ICP compared with the ONSD in brain-injured patients¹⁹. The ONSD/ETD ratio of 0.21 can be used as a simple reference tool for the diagnosis of papilledema and elevated ICP in pediatric patients²⁰. The mortality noticeably increased in comatose patients with acute stroke when the ONSD was >5.0 mm or the ONSD/ETD ratio was >0.25¹⁵. Our previous study indicated that

ONSD>6.4 mm or ONSD/ETD ratio>0.25 predicted the poor prognosis of comatose patients with supratentorial lesions²¹. The ONSD/ETD ratio was negatively correlated with Glasgow outcome scale (GOS) score ($r=-.64$)²².

In the present study, the mortality increased in comatose patients with acute stroke when the ONSD was >5.7 mm or the ONSD/ETD ratio was >0.25. The abovementioned studies indicated a higher consistency of the ONSD/ETD ratio in the prognostic prediction. In addition, the AUC values of the ONSD and ONSD/ETD ratio for predicting the prognosis of comatose patients with acute stroke were 0.760 (95% CI: 0.637-0.882) and 0.808 (95% CI: 0.696-0.920), respectively. Although there was no significant difference in the performance of the two parameters in predicting the prognosis, the AUC of the ONSD/ETD ratio seemed to be higher than that of the ONSD. Therefore, additional large-scale studies are required to confirm the cut-off values of the two parameters and their predictive values.

Moreover, we found a significant difference in MAP between the death group and the survival group, which could be attributed to a higher hemodynamic instability in the death group. Although ischemic stroke is prevalent, hemorrhagic stroke can lead to a higher mortality²³. In the present study, we reported the highest mortality in the SAH group, followed by ACI and CH groups. During hospitalization, surgeries were performed on 23% of the patients with ACI, 38% of the patients with CH, and 7% of the patients with SAH. The highest mortality of SAH might be explained by a higher ICP and a greater incidence of seizures. The lowest mortality observed in CH might be related to active decompressive surgery during hospitalization. These results confirmed the necessity of decompressive surgery for comatose patients with acute stroke to prevent cerebral hernia and to improve the prognosis.

However, the present study had some limitations. Firstly, the study was retrospective and had a small sample size, with unavoidable biases. Secondly, we only classified patients based on their survival status upon discharge, and a long-term follow-up was not conducted. In the future, the prognostic values of the ONSD and ONSD/ETD ratio

in comatose patients with acute stroke will be further verified by large-scale multi-center studies.

CONCLUSION

According to our study, ¹ the ONSD and ONSD/ETD ratio were significantly higher in comatose patients with acute stroke in the death group than those in the survival group. The mortality increased in comatose patients with acute stroke when the ONSD was >5.7 mm or the ONSD/ETD ratio was >0.25 . Both indexes could be used as prognostic indicators for comatose patients with acute stroke. However, the ONSD/ETD ratio was more stable than the ONSD and would be preferred in clinical practice.

ARTICLE HIGHLIGHTS

Research background

Stroke is mainly accompanied by an increased intracranial pressure (ICP), leading to more severe brain damage via secondary cerebral ischemia and hernia. A sudden rise of ICP is typically observed in comatose patients with acute stroke. Measuring optic nerve sheath diameter (ONSD) is noninvasive and convenient in comatose patients. Previous studies have shown that computed tomography (CT) can be used to accurately measure ONSD. The strong correlation between ICP and ONSD has been generally identified.

Research motivation

However, the ONSD of normal and sick individuals mainly varies noticeably. The ONSD/ETD ratio slightly varies. Therefore, the ONSD/ETD ratio has a smaller variability and a higher stability, and it may be more appropriate for ICP monitoring. To date, ⁴ several studies have shown that the ONSD/ETD ratio is more related to ICP than the ONSD. Therefore, it is reasonable to speculate that the ONSD/ETD ratio has a higher value in predicting the prognosis of neurological function.

Research objectives

In the current study, unenhanced cranial CT scan was used to measure the ONSD and ETD, and the prognostic values of the ONSD and ONSD/ETD ratio in comatose patients with acute stroke were compared. The AUC values of the ONSD and ONSD/ETD ratio for predicting the prognosis of comatose patients with acute stroke were 0.760 (95% CI: 0.637-0.882) and 0.808 (95% CI: 0.696-0.920), respectively. Although there was no significant difference in the performance of the two parameters in predicting the prognosis, the AUC of the ONSD/ETD ratio seemed to be higher than that of the ONSD. This study confirmed that ONSD/ETD can better predict neurological outcomes, and the variation is small, which provides a reference for non-invasive ICP monitoring and prediction of neurological outcomes in the future.

Research methods

A total of 67 comatose patients with acute stroke were retrospectively recruited. The ONSD and ETD were measured by cranial computed tomography (CT) scan. The ONSD was measured vertically at 3 mm behind the eyeball. ETD was measured from one side of the retina behind the lens to the other side for the maximum diameter. All patients underwent cranial CT scan within 24 h after coma onset. Patients' baseline clinical data were collected, including age, gender, body weight, body mass index (BMI), mean arterial pressure (MAP), stroke type, GCS scores, surgery during hospitalization (e.g., clearance of hematoma and decompressive craniectomy). Patients were divided into death group and survival group according to their survival status at discharge. The differences of the ONSD and ONSD/ETD ratio between the two groups and their prognostic values were compared by MedCalc software.

Research results

In this study, 42 (62.7%) patients died, while 25 (37.3%) patients survived at discharge. There were significant differences in the GCS score ($\chi^2=49.809$, $P<0.0001$), stroke type ($\chi^2=11.981$, $P=0.003$), ONSD ($t=4.078$, $P<0.0001$), ONSD/ETD ratio ($t=4.625$, $P<0.0001$), and surgery ($\chi^2=4.803$, $P=0.048$) between the two groups. The ONSD and

ONSD/ETD ratio were included in the logistic regression model for multivariate analysis, respectively. After adjusting for age, MAP, GCS score, stroke type, and surgery, the associations among the ONSD ($P=0.04$), ONSD/ETD ratio ($P=0.036$), and mortality were still significant.

² The area under the curve (AUC) of the ONSD was 0.760 (95% confidence interval (CI): 0.637-0.882, $P<0.0001$), with a sensitivity of 81.0% and a specificity of 64.0% at a cut-off value of 5.7 mm. The AUC of the ONSD/ETD ratio was 0.808 (95% CI: 0.696-0.920, $P<0.0001$), with a sensitivity of 92.9% and a specificity of 68.0% at a cut-off value of 0.25. The AUC values of the ROC curves of the ONSD and ONSD/ETD ratio were compared. There was no significant difference in the AUC values of the two indices ($Z = 1.333$, $P = 0.1826$).

The results of this study provide a reference for noninvasive ICP monitoring and prediction of neurological outcomes in the future.

³ In the future, we need to expand the sample size to determine the value of ONSD/ETD ratio in predicting increased ICP and poor prognosis.

Research conclusions

In this study, the area under the ROC curve of ONSD and ONSD/ETD ratio was compared by MedCalc for the first time. Although there was no significant statistical difference in P value, the area under the ROC curve of ONSD/ETD ratio tended to increase compared with ONSD, and the degree of variation was smaller. Therefore, ONSD/ETD ratio is more recommended for ICP monitoring and predicting prognosis.

Research perspectives

Increase the sample size to study single etiology

13%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|----------------|
| <div style="background-color: red; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">1</div> | bmcneurol.biomedcentral.com
<small>Internet</small> | 175 words — 5% |
| <hr/> | | |
| <div style="background-color: magenta; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">2</div> | Sha Zhu, Chao Cheng, Dianjiang Zhao, Yuanli Zhao, Xianzeng Liu, Jun Zhang. "Clinical Value of Optic Nerve Sheath Diameter Assessment in Prognosis of Comatose Patients with Supratentorial Lesions", Research Square, 2021
<small>Crossref Posted Content</small> | 89 words — 2% |
| <hr/> | | |
| <div style="background-color: purple; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">3</div> | www.ncbi.nlm.nih.gov
<small>Internet</small> | 54 words — 1% |
| <hr/> | | |
| <div style="background-color: teal; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">4</div> | Jin Young Youm, Jae Hoon Lee, Hyun Seok Park. "Comparison of transorbital ultrasound measurements to predict intracranial pressure in brain-injured patients requiring external ventricular drainage", Journal of Neurosurgery, 2022
<small>Crossref</small> | 41 words — 1% |
| <hr/> | | |
| <div style="background-color: green; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">5</div> | bmcgastroenterol.biomedcentral.com
<small>Internet</small> | 32 words — 1% |
| <hr/> | | |
| <div style="background-color: brown; color: white; text-align: center; width: 40px; height: 40px; line-height: 40px; margin: 0 auto;">6</div> | Jie Du, Yanjun Deng, Hua Li, Shigang Qiao, Mengnan Yu, Qingya Xu, Chen Wang. "Ratio of Optic Nerve Sheath Diameter to Eyeball Transverse Diameter by Ultrasound Can Predict Intracranial Hypertension in Traumatic Brain Injury Patients: A Prospective Study", Neurocritical Care, 2019
<small>Crossref</small> | 31 words — 1% |
-

- 7 assets.researchsquare.com 18 words — < 1%
Internet
-
- 8 Xuan Zhang, Joshua Medow, Bermans Iskandar, Fa Wang, Mehdi Shokoueinejad, Joyce Koueik, John Webster. "Invasive and noninvasive means of measuring intracranial pressure: a review", *Physiological Measurement*, 2017 17 words — < 1%
Crossref
-
- 9 advances.umw.edu.pl 16 words — < 1%
Internet
-
- 10 T. Ertekin, M. G. Boyaci, A. Bilir, A. Yucel, A. Ertekin, O. Turamanlar, R. Duman. "Optic nerve sheath diameter measurement: a means of detecting increased intracranial pressure in pseudotumor cerebri patients", *Folia Morphologica*, 2022 15 words — < 1%
Crossref
-
- 11 pesquisa.bvsalud.org 13 words — < 1%
Internet
-
- 12 www.mryuan.net 12 words — < 1%
Internet

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS
EXCLUDE MATCHES < 12 WORDS