

82782_Auto_EditedC.docx

Name of Journal: *World Journal of Psychiatry*

Manuscript NO: 82782

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Evaluation of chronic idiopathic tinnitus and its psychosocial triggers

Hamed SA *et al.* Psychosocial triggers of tinnitus

Sherifa Ahmed Hamed, Fadia Ahmed Attiah, Mohamed Fawzy, Mohamed Azzam

Abstract

BACKGROUND

The tinnitus susceptibility patterns in relation to different psychological and life stressors are unknown in different cultures.

AIM

To determine the comorbid psychosocial factors and behaviors associated with tinnitus and the predictors for the increase in its severity.

METHODS

Participants were 230 adults (males = 70; females = 160; mean age = 38.6 ± 3.3). They underwent audiograms, speech discrimination and masking testing and neuropsychiatric evaluation. Measures used for assessment included tinnitus handicap inventory, ¹⁷depression anxiety stress scale 21 (DASS-21), perceived stress scale (PSS) and insomnia severity index (ISI).

RESULTS

Patients had mean duration of tinnitus of 11.5 ± 2.5 mo. They had intact hearing perception at 250-8000 Hz and 95 (41.3%) had aggravation of tinnitus loudness by masking noise. Decompensated tinnitus was reported in 77% ($n = 177$). The majority had clinically significant insomnia (81.3%), somatic symptoms (75%) other than tinnitus and perceived moderate (46.1%) and high (44.3%) stress to tinnitus. The severe/extremely severe symptoms of depression, anxiety and stress were reported in 17.4%, 35.7% and 44.3%, respectively. Patients with decompensated type had significantly higher scores for ISI ($P = 0.001$), DASS-21 (depression = 0.02, anxiety = 0.01, stress = 0.001) compared to those with compensated tinnitus. Psychiatric interviewing showed that 35.7% had non-specific anxiety disorder, 17.4% had major depression and 19.6% fulfilled the criteria of somatization disorder. Multivariate analysis showed that the only independent predictors for tinnitus severity were the duration of tinnitus [odd ratios (OR) = 0.832, 95%CI: 0.640-1.158, $P = 0.001$] and PSS (OR = 0.835, 95%CI: 0.540-1.125, $P = 0.001$) scores.

CONCLUSION

Up to our knowledge, this is the first study in our culture which evaluated the causal relationship between psychological factors and tinnitus onset, severity and persistence. Tinnitus could be the earliest and dominant somatic symptom induced by life stressors and psychological vulnerabilities. Therefore, multidisciplinary consultation (psychologists, psychiatrists and neurologists) is important to acknowledge among the audiologists and otolaryngologists who primarily consult patients.

Key Words: Chronic tinnitus; Idiopathic; Somatization; Psychosocial factors

Hamed SA, Attiah FA, Fawzy M, Azzam M. Evaluation of chronic idiopathic tinnitus and its psychosocial triggers. *World J Psychiatry* 2023; In press

Core Tip: Tinnitus is a very frequent subjective symptom in adults. It has been found that 15%-30% of patients with tinnitus had no clinically manifest hearing loss or even had no

subclinical sensorineural hearing loss when evaluated using advanced auditory testing. Previous studies found that there are several comorbid psychiatric conditions and disorders in sufferers of tinnitus, which also may contribute to its persistence and increased severity. They include stress, anxiety, depression, sleep disturbance, increased forgetfulness, major depression and anxiety and somatoform disorders. However, the relationship of these comorbidities to the onset of tinnitus is still unclear. Also the psychosocial triggers for initiation, increased severity and chronicity of tinnitus are understudied in many areas of the world.

INTRODUCTION

⁶ Tinnitus is the most common subjective auditory symptom with an estimated prevalence of 8%-20% (~14.4%) in adults and 35%-45% (~40%) in individuals aged ≥ 55 years old. The prevalence of tinnitus is reduced after the 7th decade^[1]. It has also been reported that tinnitus is a life-lasting disabling problem in 0.5%-2.5% of patients^[1-3] and a handicapping problem in 1%-2%^[4]. Tinnitus is defined as a sound heard in the ears or ears and head in absence of external auditory source. It is commonly described as ringing, whistling, buzzing, roaring and clicking and rarely as a combination of sounds or a non-defined sound. Some audiologists suggested that the majority of patients with tinnitus (~90%) might have a degree of hearing loss which varied from severe to very mild or subclinical^[5]. However, others found tinnitus in 30% of patients without clinically manifest sensorineural hearing loss (SNHL) and in 15% of patients without evidence of hearing impairment using advanced testing for detection of subclinical deficits^[6]. They also suggested that persistence of tinnitus and aggravation of its severity overtime (decompensated tinnitus) highly indicate the presence of psychological or other physical (to less extent) comorbid conditions, because tinnitus due to lesions in the auditory system is often transient and disappears overtime due to brain filtering mechanisms regardless to the cause^[6]. The common comorbid conditions in patients with chronic tinnitus include emotional stresses, mood swings and anxiety symptoms (28%-49%)^[7-9], sleep problems (10%-80%)^[10,11], poor attention and concentration (up to 70%)^[9], and chronic physical diseases (21%-47%)^[12]. Research studies also reported that 45%-78% of patients with chronic tinnitus satisfied the criteria of at least one psychiatric disorder(s), for example affective (~23.5%), anxiety (~35%) and somatoform (~25%) disorders^[2,8,9,13-20].

Various medications and interventions are used in the treatment of tinnitus including vasodilators, vitamins, calcium channel blockers, antihistamines, anticonvulsants (*e.g.* valproate), gabapentin, antidepressants, anxiolytics, psychotherapy, biofeedback and hypnosis. However, none ²⁴ have been approved by the Food and Drug

Administration. Many clinicians agreed that combined management strategies is more effective for treatment of tinnitus^[7,21-24].

Whether psychological/psychiatric factors are risks for bringing tinnitus is understudied in different countries, therefore, in this work, we aimed to determine: (1) The comorbid psychosocial factors and behaviors associated with tinnitus's onset, severity and chronicity; and (2) The predictors for the increase in its severity.

MATERIALS AND METHODS

Study design and participants

This is a cross-sectional study. It included 230 adults (males = 70; females = 160) with chronic tinnitus. Patients were recruited from January 2020 to July 2022 from the outpatient Otolaryngology and Neuropsychiatry Clinics of Assiut University Hospital, Assiut Egypt. Inclusion criteria: (1) Adults (20-55 years old); (2) Self-reported tinnitus as a main physical problem at presentation; (3) Duration of tinnitus ≥ 6 mo; (4) Normal results of previous routine audiometric testing which were done at different time points; and (5) Normal imaging (computed tomography or magnetic resonance imaging) of the brain and cervical spine or cord. Exclusion criteria: (1) Pulsatile tinnitus; (2) Manifest hearing loss; (3) Local cause which explain tinnitus, for example impacted wax, infections/inflammation of the ear, Meniere disease, otosclerosis, presbycusis, etc; (4) Medical, neurological or psychiatric disease which can cause hearing impairment (e.g. diabetes, anemia, thyroid disease, stroke, small vessel disease, demyelinating lesions, migraine, epilepsy, tumor as acoustic neuromas, psychosis, etc); (5) Head or cervical Vertebra injuries; and (6) Ototoxic drug use (e.g. aspirin, quinine, aminoglycosides, cancer chemotherapy as cisplatin, etc).

Procedures

Data collection included: (1) Demographics: age, gender, residence, marital state, educational level (classified as low if can't read or write, can read, primary or secondary school education or high if high school, collage, etc) and socioeconomic state. The socio-

economic scale (SES) was used for determination of socioeconomic state, SES is a structured questionnaire used to gather data related to parents' education, family income per month, sanitation and crowding index. SES total score is 30. SES was classified as high (score 25 to ≥ 30), middle (score > 20 to ≥ 25), low (score > 15 to < 20) or very low (score < 15)^[25]; and (2) Clinical characteristics: Otological, medical, neurological and psychiatric histories and examinations were done. Tinnitus inquiries included: its duration, origin, description, laterality, intensity, pattern and time of occurrence. The results of previous evaluations (audiograms, laboratory investigations and drug prescriptions for tinnitus) were also gathered. The diagnosis of a psychiatric disorder was done according to the Structured Clinical Interviewing using Arabic version of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Revised (DSM-IV-TR)^[26,27].

The diagnostic work-up questionnaires included: Tinnitus handicap inventory (THI)^[28]: It consists of 25 items, with 3 responses for each which are 1st response or "yes" equals 4 points, or 2nd response or "sometimes" equals 2 points or a 3rd response "no" equals 0. Its total scores range from 0 to 100 points. Higher scores indicate greater perceived handicap. Tinnitus severity was classified as: (1) Grade 1 or slight (score 0-16) if tinnitus was heard only in quiet environment, masked easily with environmental noise and did not interfere with sleep or daily activities; (2) Grade 2 or mild (score 18-36) if tinnitus was masked easily by environmental noise, neglected during life activities and occasionally interfered with sleep, but did not interfere with daily activities; (3) Grade 3 or moderate (score 38-56) if tinnitus could be noticed in noisy environment but the individual could still perform the daily activities; (4) Grade 4 or severe (score 58-76) if tinnitus was always heard, disturbed sleep and could interfere with daily activity; and (5) Grade 5 or catastrophic (score 78-100) if tinnitus was severe enough to interfere with any life activity. For comparative statistics, tinnitus was divided according to its severity into compensated tinnitus if THI grades were 1 or 2 (*i.e.* slight or mild) and decompensated tinnitus if THI grades were 3, 4 or 5 (*i.e.* moderate, severe or catastrophic).

Insomnia severity index (ISI): It is 7 items questionnaire^[29]. The total score categories are: (1) No clinically significant insomnia (score 0-7); (2) Sub-threshold insomnia (score 8-14); (3) Moderately severe clinically significant insomnia (score 15-21); and (4) Severe clinically significant insomnia (score 22-28).

Depression anxiety stress scale 21 (DASS-21)^[30]: It is 21 items which produces results for the recent (past week) symptoms. The DASS-21 scores for depression or anxiety or stress symptoms were determined separately. They were classified as normal, mild, moderate, severe and extreme.

Perceived stress scale (PSS)^[31,32]: It is 14 items scale. Each item is rated by a 5-point Likert scale which range from 0: "never" to 4: "very often". The total score of each individual was calculated by reversing the scores on these positive seven items (B4, B5, B6, B7, B9, B10, and B13), followed by adding the responses to the rest of the 14 items. PSS does not reflect a specific diagnosis or course for treatment; therefore, it has no cut-off value. PSS scores range from 0 to 56. Accordingly, subjects were classified according to the perceived stress level to tinnitus (as stressor) into: (1) Perceived low stress if scores were 0 to 13; (2) Perceived moderate stress if scores were 14 to 26; and (3) Perceived high stress if scores were 27 to 40.

A designed questionnaire for determination of life stressors. The term "life stressor" is defined as extraordinary and undesirable major life events or conditions of clear onset and offset. We designed a simple questionnaire based on the common major stressors in our population which included financial, marital, siblings', changes in personal work or home activities', troubles with law and problems with a close relatives' or a friend.

Basic audiologic evaluation: It included pure tone audiometry which was assessed at frequency ranges from 250 Hz to 8000 Hz (interacoustic model AC 40), tympanometry (interacoustic model AZ 26), measurement of acoustic reflex threshold (interacoustic model AZ 26) and speech discrimination test. Speech discrimination score (SDS) is defined as the hearing level to understand and repeat at least 25 Arabic phonetically balanced monosyllabic words. SDS of 90% to 100% was considered normal. For patients

who noticed tinnitus at quite occasions (e.g. at bedtime), a masking test was also done using a white noise machine.

8

Statistical analysis

Data were analyzed with SPSS (IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp). The distributional properties of variables and the appropriateness of the analyses of covariance were confirmed. Comparative statistics were done using two-sided Student's *t* test, Chi square test and one way analysis of variance with Bonferroni post hoc correction. Inferential statistics were performed using Spearman's Correlation Coefficient between: (1) Scores of THI and age, duration of tinnitus and scorings of DASS-21 (depression, anxiety and stress), PSS and ISI; (2) scores of DASS-21 and PSS and ISI; and (3) scores of PSS and ISI. Multivariate analysis was done to detect the independent variables associated with THI. First, univariate analysis was done between scores of THI and duration of tinnitus and scores ISI, DASS-21 and PSS. Variables which showed significant values in univariate model were entered into the multivariate model. Results were expressed as odd ratios (OR) and 95%CI. Significances were calculated with probability value less than 0.05.

RESULTS

Demographics and clinical characteristics

Participants were 230 adults with chronic tinnitus. They had age range of 25 to 50 years (mean = 38.6 ± 3.3). The mean duration of tinnitus at the time of the study was 11.5 ± 2.5 mo (range: 6 to 24 mo). The majority were in their 3rd and 4th decades ($n = 147$, 64%), females (69.6%), rural residents (62.6%) and married (63.5%) and of low education (79.1%) and low/middle socioeconomic states (53%). They had intact hearing perception at 250-8000 Hz, SDS and type A tympanometry. One hundred and forty patients had tinnitus at quite (or bedtime) and 95 of them (or 67.86%) had aggravation of tinnitus loudness by masking noise. Decompensated tinnitus was reported in 77% ($n = 177$; G3 = 52 or 22.6%, G4 = 62 or 27% and G5 = 63 or 27.4%) and compensated tinnitus was reported in 23% (n

= 53; G1 = 15 or 6.5%, and G2 = 38 or 16.5%). Figure 1 showed the demographics of the studied patients and the differences in demographics between patients with compensated *vs* decompensated tinnitus. There were no significant differences between the two groups of patients in relation to different variables. Table 1 showed the characteristics of tinnitus. It showed that the most frequently encountered sounds were whistling (33.04%) and different sounds' descriptions in different occasions (27%), while clicking (7.4%) and sound echo (4%) were the least described types. Bilateral tinnitus of equal intensity in both ears was more frequent than unilateral. Tinnitus was either continuous or periodic. Thirty percent of patients observed that ear plugging resulted in reduced tolerance to tinnitus loudness. There were no significant differences in the tinnitus characteristics between patients with compensated and decompensated types except that the latter had significantly longer duration of tinnitus at presentation ($P = 0.03$).

Psychological characteristics and stressors

Table 2 showed the results of the diagnostic work-up questionnaires and psychiatric interviewing of patients. It showed that the majority had clinically significant insomnia ($n = 187$, 81.3%). The severe and extremely severe symptoms of depression, anxiety and stress were reported in 17.4% ($n = 40$), 35.7% ($n = 82$) and 44.3% ($n = 102$) of patients, respectively. Also, the majority of patients had perceived moderate ($n = 106$ or 46.1%) and high ($n = 102$ or 44.3%) stress to tinnitus. There were no gender differences in scores of ISI, DASS-21 and PSS. Patients with decompensated type had significantly higher scores for ISI ($P = 0.001$), DASS-21 (depression = 0.02, anxiety = 0.01, stress = 0.001) compared to those with compensated tinnitus.

There were high frequencies of patients with somatic symptoms (75%) other than tinnitus. They included non-specific headache (47.9%), gastrointestinal symptoms (41.3%) (*e.g.* vomiting, bloating, diarrhea, abdominal gaseous distention, colic, *etc* without definite etiology or similar concurrent conditions in close household contacts), poor concentration (38.3%) and erectile dysfunction (31.7%). Psychiatric interviewing showed

that 35.7% had non-specific anxiety disorder, 17.4% had major depression and 19.6% fulfilled the criteria of somatization disorder. Higher frequency of somatization disorder (23.7% *vs* 5.7%) and sexual dysfunction (49% *vs* 26.6%) were reported in patients with decompensated compared to those with compensated tinnitus. Family history of psychiatric disorders was reported in only 9.57% of patients.

The most frequently reported life stressors were financial (77.4%), marital (73%) and siblings' (69.6%) issues. Few (8.83%) denied presence of life stressors (Table 3). We observed that there were no significant differences in the frequencies of life stressor for patients with decompensated compared to those with compensated tinnitus.

Results of correlation analyses

Significance correlations were found between duration of tinnitus and its severity and perceived stress towards it and severities of depression, anxiety and stress symptoms and insomnia; between perceived stress towards tinnitus and severities of depression, anxiety and stress symptoms and insomnia; and between severities of insomnia and depression, anxiety and stress symptoms (Table 4). Univariate analysis showed that there were significant correlations between THI scores and duration of tinnitus (OR = 0.913, 95%CI: 0.758-1.262, $P = 0.02$) and scores of ISI (OR = 0.853, 95%CI: 0.650-1.128, $P = 0.001$), DASS21 (depression: OR = 0.430, 95%CI: 0.320-0.656, $P = 0.01$; anxiety: OR = 1.156, 95%CI: 0.886-1.902, $P = 0.001$; stress: OR = 0.932, 95%CI: 0.843-1.230, $P = 0.001$) and PSS (OR = 0.923, 95%CI: 0.643-1.258, $P = 0.001$). Multivariate analysis showed that the only independent predictors for tinnitus severity (THI score) were the duration of tinnitus (OR = 0.832, 95%CI: 0.640-1.158, $P = 0.001$) and PSS score (OR = 0.835, 95%CI: 0.540-1.125, $P = 0.001$).

DISCUSSION

The topic of chronic idiopathic tinnitus and its relation to psychological factors is understudied in different countries. The core outcome domain of this study was identification of psychosocial associates with tinnitus's onset, severity and persistence. Generally speaking, previous studies included the following information about tinnitus:

(1) Lesions in the auditory sensory pathway and its brain circuits often produce high-pitched tinnitus (*i.e.* hearing loss is dominant in high-frequency tone sound stimuli with ranges more than 4000 Hz. These ranges are not routinely tested in audiology clinics^[5]. Tinnitus caused by auditory pathway lesions is often transient and disappears spontaneously within short time regardless to its etiology. This is due to the activation of thalamic filtering processes by the superior brain centers (*i.e.* switch-off the aberrant signals expressed as noisy ear sounds). This process is termed habituation or adaptation^[6]; (2) Clinicians observed that it is impossible at onset to predict the course of tinnitus or whether an individual will develop compensated or decompensated tinnitus overtime. However, audiologist and otologists usually advice the majority (~80%) of individuals with tinnitus against frequent consultations as long as there is no hearing deficits and intact comprehensive of speech, and to become accustomed to tinnitus^[16,20] as there is/are no standard symptomatic treatment(s) for tinnitus. The clinicians also observed that there is no direct relation between tinnitus loudness and the severity of SNHL. There are individuals with very severe hearing loss but never experienced tinnitus^[6]. In contrast, some clinicians reported an association between the estimates of hearing loss and tinnitus loudness. However, such studies were criticized as their authors did not exclude (during analyses) the co-variables which could associate with tinnitus severity (as emotional stress or psychosocial factors)^[5,15]; (3) **Most studies are focused on tinnitus as a somatic stressor**, *i.e.* the cause of adverse mental health. They also suggested that the suffers of tinnitus seem to be more vulnerable to significant stress^[16]. However, few indicated that tinnitus may begin in times of severe stress or after a period of significant stress^[20]; and (4) In many countries, the management of chronic tinnitus may include interdisciplinary approach (*i.e.* neurologists, audiologists, psychologists and psychiatrists)^[1-3].

In our sample, women to men ratio, was 2.3:1. The gender difference of prevalence of chronic tinnitus is debated. Some studies reported higher frequencies in females. Others found the reverse^[12,18]. In this study, the higher frequency of participants from females could be explained by the following: (1) females may be vulnerable to develop

chronic tinnitus than male; (2) Being housewives, females had free times for consultations than males; (3) Females are commonly express their somatic symptoms, feelings and stress more easily and frequently compared to males; and (4) Female s with or without tinnitus had higher psychological risks throughout their lives and were more likely to exhibit emotional disturbances (e.g. mood swings, anxiety and stress), while males tended to experience depression and social isolation^[9].

In this study, psychiatric interviewing revealed that participants had higher frequencies and severities of insomnia, anxiety and stress symptoms. Many patients (75%, $n = 172$) had repeated somatic symptoms in addition to tinnitus which are distressing or result in significant disruption of daily life. They included insomnia, headache, gastrointestinal tract (GIT) symptoms (vomiting, bloating, diarrhea, *etc*) without explanatory cause, erectile dysfunction and memory deficits. These manifestations were marked in patients with decompensated tinnitus.

Considering the age of participants at presentation, ~20% satisfied the DSM-IV criteria of somatization disorder, a class of somatoform disorders^[26,27]. The same criteria were applied for the classification in DSM-IV as “somatic symptom disorder”, a class of somatic symptom and related disorder^[33]. These criteria were previously termed as “psychosomatic or psycho-physiologic disorders”. Non-specific anxiety disorder was reported in 35.65%. Major depressive disorder was reported in 17.39%.

Exploring the psychosocial stressors, the majority (77%) had drastic financial, marital or siblings' stressful life events. In accordance, previous studies reported that 45%-78% of patients with chronic idiopathic tinnitus satisfied the criteria of one or more psychiatric disorders^[2,7-14,16-18]. Some studies considered the role of stress-reactivity in tinnitus onset^[6,19,20]. ³ Belli *et al*^[17] showed that 27% of patients with tinnitus had at least one psychiatric diagnosis (as anxiety disorders in 28%, somatoform and mood disorders in 15%, and personality disorders in 3%) *vs* 5.6% for control subjects. The authors suggested that, the psychological stressor greatly impacts the body physiological function with improper stimulation of the autonomic nervous system, excess adrenaline secretion and severe stress and anxiety and somatization. Fagelson^[34] did a chart review for enrolled

patients to tinnitus service in the Veterans Affairs Medical Center. The authors reported that 34% of patients with tinnitus also fulfilled the criteria for the diagnosis of posttraumatic stress disorder (PTSD). They also reported that the following tinnitus inquiries were more frequent in patients with tinnitus and PTSD compared to those with tinnitus only, they include sudden onset of tinnitus, increased tinnitus severity, poor tolerance to sounds and exaggeration of tinnitus with sounds. The authors suggested that there might be several shared neural mechanisms between tinnitus and PTSD and affect auditory behaviors (*i.e.* mutual reinforcement of symptoms of both tinnitus and PTSD). Furthermore, several studies reported that the chronic and severe tinnitus which followed traumatic brain injury, whether the involved individuals were from military or civil personnel, is due to the emotional trauma or the higher distress level which was experienced during the trauma event and not directly associated with the specific head or neck injury^[35,36].

We observed the following associates and characters of tinnitus in the majority of patients which highly suggest a contribution of non-sensorineural auditory problem: (1) There was no deterioration of hearing over time with repeated audiometric tests despite increase in tinnitus severity in the majority (77%)^[5]; (2) In the same individual in different occasions, tinnitus originated from inside the head or the ear or both (36.09%), alternate in origin from either left or right (15.22%) and had different descriptions (*e.g.* ringing at sometimes and another description at another time) (27%); (3) The bilateral presentation and equal intensity in both ears since onset (43.61%); (4) The descriptions of tinnitus as whistling (33.04%), roaring (14.78%), clicking (7.4%) or sound echo (4%), which are non-familiar descriptions for tinnitus associated with SNHL; (5) The non-continuous or periodic pattern (47.83%); (6) Tinnitus was aggravated or became intolerable with ear plugging or masking noise (41.30%); and (7) Tinnitus severity and its perceived stress were associated with the severity of anxiety, stress and depression symptoms and insomnia^[15,16,20].

Research studies found that insomnia reduced tolerance and increased the perceived annoyance to tinnitus^[10,17,37]. Authors have linked the conscious perception of tinnitus

and its loudness with time and insomnia to the hyper-arousal induced by hyperactivity of the sympathetic nervous system^[38]. We also found that the only independent associates to severe tinnitus were its chronicity and higher perceived stress towards it. We suggested that psychological factors influence the conscious perception of tinnitus, result in worsening of tinnitus (*e.g.* became continuous, increase in its loudness and prolonged its duration) and also generate other functional somatic conditions as headache, poor concentration, GIT manifestations, erectile dysfunction, *etc.* Research studies found an independent association between neuroticism, obsession and anxiety traits and tinnitus severity^[15,16]. It has been suggested that focusing attention on tinnitus increases fear, tension and anxiety and emotional reinforcements followed by emotional exhaustion, neuronal hyper-excitability, establishment of a neuronal plasticity in the limbic system, particularly the amygdala and a vicious circle, resulting in chronic and severe continuous perception of tinnitus^[13,14,17,39].

The results of this study support that the psychological factors play an ¹³important role in the processes of perception, interpretation, treatment, and complications of tinnitus^[40]. Therefore, combined multidisciplinary management of patients with tinnitus is required. Neuropsychiatric evaluation has to include counseling, education, and reassurance, and reduction of depression, anxiety and stress and improving sleep pattern by psychotropic medication, psychotherapy or other interventions^[7,21-24]. Psychotropic drugs improve tinnitus by improving the depression anxiety, stress and sleep disturbance. The previously used medication which have shown efficacy included tricyclic antidepressants (as amitriptyline, nortriptyline and trimipramine) and ²¹selective serotonin reuptake inhibitors (SSRIs) (as sertraline, citalopram, escitalopram, fluoxetine, venlafaxine, fluvoxamine, and paroxetine)^[21]. Folmer ¹*et al*^[21] in their retrospective study on 30 patients with depression and started treatment with SSRIs and psychotherapy for 20.6 mo after tinnitus onset, the authors demonstrated a significant improvement of tinnitus as evidenced by reduction of tinnitus severity index scores. The authors concluded that, SSRIs elevated affectivity which accompanied tinnitus and therefore reduced the intensity and frequency of tinnitus. However, some studies found that

although SSRIs improve symptoms of depression and anxiety but may worsen tinnitus^[22,41]. In tinnitus mouse models, Tang *et al*^[42] analyzed the brain tissue which focused on the response of neurons in the dorsal cochlear nucleus to serotonin. The dorsal nucleus is the portion of the cochlear nucleus with inhibitory characteristics and involved in sensory processing. It is the portion of the cochlear nucleus which is affected by tinnitus. The authors observed hyperactivity and hypersensitivity of the mice to sounds when the fusiform cells in the dorsal cochlear nucleus was exposed to serotonin, indicating that serotonin raises neuronal activity in the dorsal cochlear nucleus. They also suggested that serotonin suppresses signaling through the auditory pathway and enhances transmission through a multisensory pathway. Previous studies also suggested that the main aim for the use of psychotropic medications is to treat the co-morbid anxiety/depression in patients with tinnitus and not as a main symptomatic treatment of tinnitus. Because not all psychotropic medications can worsen tinnitus, it is plausible to suggest that it is better to switch to another psychotropic drug to improve depression and anxiety symptoms without deterioration of tinnitus.

Up to our knowledge, this is the first study in our culture which evaluated the causal relationship between psychological factors and tinnitus onset, severity and persistence. However, this study has limitations which include a female predominance (*i.e.* gender bias), however, we do not consider this a major weakness as statistical analyses were also done after classification of patients from both gender into those with compensated *vs* decompensated tinnitus.

CONCLUSION

Tinnitus was an infrequent earliest and dominant somatic symptom induced by severe life stressors and psychological vulnerabilities. In many patients, the characters of tinnitus were distinguished from that of solely auditory system lesions. Psychiatric interviewing revealed the diagnosis anxiety disorder in 35.7%, somatization disorder in 19.6% and major depression in 17.4% of patients. Tinnitus severity was independently associated with the longer duration of tinnitus and the individual's perceived stress

towards tinnitus. Therefore, multidisciplinary consultation (psychologists, psychiatrists and neurologists) is important to acknowledge.

ARTICLE HIGHLIGHTS

Research background

⁶ Tinnitus is the most common auditory symptom with an estimated prevalence of 8%-40% and life-long disabling problem in ~2.5%. Tinnitus an aberrant sound heard in the ears or in both ears and head. It is believed to be a symptom of hearing impairment. However, many audiologists observed that approximately 15%-30% of subjects with tinnitus had neither manifest nor subclinical hearing impairment. Research studies indicated that psychosocial factors as stress, anxiety, depression and insomnia are strong risks for the chronic and severe tinnitus. They also reported comorbid psychiatric disorders as anxiety, somatization and major depressive disorders in 23%-35% of patients with tinnitus. Others reported improvement of tinnitus after the use of antidepressant or anxiolytic medications. Many audiologist and otolaryngologist believe that tinnitus requires multidisciplinary evaluation and management. As chronic idiopathic tinnitus and evaluation of its psychosocial triggers is an understudied topic in many parts of the world, therefore, it has to be addressed in different populations.

⁴ ***Research motivation***

The research hotspots include determination of: (1) Characteristics of chronic tinnitus in group of patients with no obvious hearing loss as a cause; (2) The comorbid psychosocial stressors and psychiatric conditions and disorders which are associated with tinnitus initiation, aggravation and chronicity; ⁴ and (3) The predictors which are independently associated with increase in the severity of tinnitus.

⁴ ***Research objectives***

This study systematically assessed patients with chronic idiopathic tinnitus and the psychosocial factors associated with its onset, increased severity and chronicity.

Research methods

Tinnitus handicap inventory, depression anxiety stress scale 21, ²perceived stress scale, insomnia severity index and a designed questionnaire for determination of life stressors.

Research results

This study included 230 adult patients from both gender. They had chronic tinnitus with a mean duration of approximately eleven months. Their previous ear, nose and throat and audiology evaluations including pure tone audiometry at frequencies ranged from 250 to 8000 Hz did not reveal obvious cause for tinnitus. Decompensated tinnitus was reported in 77%. The characters of tinnitus in many patients are distinguished from that of tinnitus due to pure auditory pathway lesions, which included being originated from ears and/or head, bilateral or alternate ear side, described as whistling, roaring, clicking, or combined sound, aggravated by ear plugging, stress, anxiety and insomnia and absence of hearing deterioration overtime. Psychiatric evaluation revealed frequent comorbid conditions and disorders including stress, anxiety, somatic problems, depression, enhanced perceived stress towards tinnitus and anxiety, somatization and major depressive disorders. We also reported ²³that the variables which were highly significant and independently associated with tinnitus were its duration and increased stress perception towards tinnitus.

Research conclusions

Tinnitus was a common and predominant somatic symptom in response to severe life stressors and psychological susceptibility. The tinnitus variables showed distinct characteristics which differentiate it from tinnitus caused by pure auditory system lesions. Psychiatric interviewing revealed that the majority of patients had symptoms of stress, anxiety, depression, and sleep disturbance of variable severities. Anxiety and somatization ¹disorders were reported in 35.7% and 19.6% of patients, respectively. Major

depressive disorder **was** reported in 17.4%. The inter-individual variability to stress perception and anxiety were strong triggers for tinnitus, its severity and chronicity.

4

Research perspectives

In our locality, the importance of the psychosocial factors for the initiation, aggravation and chronicity of tinnitus are understudied. Multidisciplinary consultation (psychologists, psychiatrists and neurologists) is important to acknowledge among the audiologists and otolaryngologists who primarily consult patients. Also future longitudinal studies are required to determine the tinnitus outcomes before and after treatment with psychotropic medications.

10%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|--|--|-----------------|
| <div style="background-color: red; color: white; padding: 2px 5px; display: inline-block;">1</div> | www.ncbi.nlm.nih.gov
<small>Internet</small> | 113 words — 2% |
| <hr/> | | |
| <div style="background-color: magenta; color: white; padding: 2px 5px; display: inline-block;">2</div> | www.science.gov
<small>Internet</small> | 52 words — 1% |
| <hr/> | | |
| <div style="background-color: purple; color: white; padding: 2px 5px; display: inline-block;">3</div> | www.dovepress.com
<small>Internet</small> | 34 words — 1% |
| <hr/> | | |
| <div style="background-color: teal; color: white; padding: 2px 5px; display: inline-block;">4</div> | Sherifa Ahmed Hamed, Mohamed Elwasify, Mohamed Abdelhafez, Mohamed Fawzy. "Peripartum depression and its predictors: A longitudinal observational hospital-based study", World Journal of Psychiatry, 2022
<small>Crossref</small> | 33 words — 1% |
| <hr/> | | |
| <div style="background-color: green; color: white; padding: 2px 5px; display: inline-block;">5</div> | www.frontiersin.org
<small>Internet</small> | 29 words — 1% |
| <hr/> | | |
| <div style="background-color: brown; color: white; padding: 2px 5px; display: inline-block;">6</div> | bmcpregnancychildbirth.biomedcentral.com
<small>Internet</small> | 18 words — < 1% |
| <hr/> | | |
| <div style="background-color: brown; color: white; padding: 2px 5px; display: inline-block;">7</div> | Bochra Nourhene Saguem, Amel Braham, Islem Romdhane, Selma Ben Nasr. "Psychological impact of home confinement due to coronavirus disease on medical students and its relationship with cognitive emotion regulation", The Journal of Mental Health Training, Education and Practice, 2021 | 17 words — < 1% |

-
- 8 pdfs.semanticscholar.org 17 words — < 1%
Internet
-
- 9 www.clinicaltrials.gov 17 words — < 1%
Internet
-
- 10 S. E. Shore. "Multisensory integration in the dorsal cochlear nucleus: unit responses to acoustic and trigeminal ganglion stimulation", *European Journal of Neuroscience*, 6/2005 14 words — < 1%
Crossref
-
- 11 www.cureus.com 14 words — < 1%
Internet
-
- 12 aci.health.nsw.gov.au 13 words — < 1%
Internet
-
- 13 www.tinnitusjournal.com 13 words — < 1%
Internet
-
- 14 Janet M. Y. Cheung, Delwyn J. Bartlett, Carol L. Armour, Bandana Saini, Tracey-Lea Laba. "Patient Preferences for Managing Insomnia: A Discrete Choice Experiment", *The Patient - Patient-Centered Outcomes Research*, 2018 12 words — < 1%
Crossref
-
- 15 K-H Lee, S-J Choi, J-H Lee, S Kim, M Seol, Y-S Lee, W-K Kim, J-S Lee, J-H Lee. "Cyclosporine alone vs cyclosporine plus methotrexate for post-transplant immunosuppression after HLA-identical sibling bone marrow transplantation: a randomized prospective study", *Bone Marrow Transplantation*, 2004 12 words — < 1%
Crossref
-

16	ijspt.org Internet	12 words — < 1%
17	pubmed.ncbi.nlm.nih.gov Internet	12 words — < 1%
18	journals.lww.com Internet	11 words — < 1%
19	Asih, Sali, Randy Neblett, Tom G. Mayer, Emily Brede, and Robert J. Gatchel. "Insomnia in a chronic musculoskeletal pain with disability population is independent of pain and depression", The Spine Journal, 2014. Crossref	10 words — < 1%
20	www.pubfacts.com Internet	10 words — < 1%
21	www.sahpra.org.za Internet	10 words — < 1%
22	E. H. Davies, A. Erikson, T. Collin - Histed, E. Mengel, A. Tylki - Szymanska, A. Vellodi. "Outcome of type III Gaucher disease on enzyme replacement therapy: Review of 55 cases", Journal of Inherited Metabolic Disease, 2007 Crossref	9 words — < 1%
23	Sherifa Ahmed Hamed, Samir Kamal Abdulhamid, Ali Farrag El-Hadad, Mohamed Fawzy, Mohamed Ahmed Abd-Elhamed. "Restless leg syndrome in patients with chronic kidney disease: a hospital-based study from Upper Egypt", International Journal of Neuroscience, 2021 Crossref	9 words — < 1%
24	vdoc.pub Internet	9 words — < 1%

25 "Correlates of Depression, Anxiety and Stress among Female Students at Al-Jouf University, Saudi Arabia", The Egyptian Journal of Community Medicine, 2017

Crossref

8 words — < 1%

26 Georgina Henry, Phyllis Butow, Ilona Juraskova, Rebekah Laidsaar-Powell, Joanne Shaw.

"Measuring female breast cancer survivors' concerns about late effects of treatment: initial development of the Concerns about Late Effects in Oncology Questionnaire (CLEO)", Supportive Care in Cancer, 2022

Crossref

8 words — < 1%

27 Ismail Selvi, Muhammet İrfan Dönmez, Orhan Ziylan, Tayfun Oktar. " Urodynamically proven lower urinary tract dysfunction in children after - 19: A case series ", LUTS: Lower Urinary Tract Symptoms, 2022

Crossref

8 words — < 1%

28 link.springer.com

Internet

8 words — < 1%

29 Zoltan Rusznak. "Membrane Currents Influencing Action Potential Latency in Granule Neurons of the Rat Cochlear Nucleus", European Journal of Neuroscience, 11/1997

Crossref

7 words — < 1%

30 Maryam A Khouj, Samera Albasri, Anas A Albishri, Shadi M Softa, Alanoud S Almaslamani, Hanin M Ahmad. "Prevalence of Stress, Anxiety, and Depression Among Pregnant Women in Jeddah", Cureus, 2022

Crossref

6 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF