

84557_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 84557

Manuscript Type: CASE REPORT

Rapid progression of heart failure secondary to radioactive iodine treatment of hyperthyroidism: A case report

Zi-han Li, Lu-jia Ni, Yuan-qiao Liu, Dao-yuan Si

Abstract

BACKGROUND

The deterioration of thyroid health is involved in the progression of heart failure (HF). This is usually a lengthy process, so there are almost no reports on its rapid development. Here we report a case of a young male who rapidly developed hypothyroid cardiomyopathy secondary to radioactive iodine treatment, suggesting that severe HF might occur even after a short period of hypothyroidism.

CASE SUMMARY

A 26-year-old man was referred to our hospital for HF presenting with dyspnea on exertion and chest discomfort lasting for 1 mo. He received radioactive iodine treatment for hyperthyroidism 1 year ago and had an almost normal echocardiogram 6 mo ago. Admission echocardiogram and cardiac magnetic resonance (CMR) revealed left ventricle (LV) global hypokinesia and severely depressed systolic function. In addition, late gadolinium enhancement indicated no obvious changes in the myocardium. Thyroid function tests showed decreased serum levels of thyroid hormone (TH) and elevated thyroid-stimulating hormone. Based on an exclusionary examination, the patient was diagnosed with hypothyroid cardiomyopathy and was started on replacement therapy. His HF symptoms were completely relieved during the six-month

follow-up, and echocardiogram and CMR revealed recovered LV size and ejection fraction.

CONCLUSION

This report demonstrates that severe fluctuations in TH levels may lead to acute HF, which can completely recover with timely thyroid hormone replacement. In addition, our findings highlight the importance of routinely detecting cardiac function in patients treated with radioactive iodine.

INTRODUCTION

Thyroid hormones have important physiological effects. The lack of thyroid hormone (TH) can affect almost every organ system in the body, resulting in various clinical manifestations [1]. In the cardiovascular system, hypothyroidism may result in reduced cardiac output, arterial hypertension, sinus bradycardia, QT interval prolongation and pericardial effusion, and it can be associated with many diseases, especially heart failure (HF)[2-3]. Conversely, HF itself can lead to reduced serum levels of active TH[4]. The deterioration of thyroid health can be involved in the progression of HF, though this is usually a lengthy process, and there are almost no reports on its rapid development. Here we report a case of a young male who rapidly developed hypothyroid cardiomyopathy secondary to radioactive iodine treatment, suggesting that severe HF might occur even after a short period of hypothyroidism.

CASE PRESENTATION

Chief complaints

A 26-year-old man was admitted to our hospital due to dyspnea on exertion and chest discomfort, with no inducing factors, 1 mo ago.

History of present illness

The patient's symptoms gradually worsened within 1 mo. Additional symptoms included a weight gain of 20 kg in the prior 3 mo and paroxysmal nocturnal dyspnea. He denied fever, upper respiratory tract infection, chest pain, palpitations or a history of drug use.

History of past illness

His medical records were notable for radioactive iodine treatment for hyperthyroidism 1 year ago, and he had an almost normal echocardiogram 6 mo ago, which revealed a left ventricle (LV) with a diameter of 48.8 mm and a normal left ventricular ejection fraction (LVEF) of 65%. He did not frequently control his cardiac or thyroid function in the past 6 mo.

Personal and family history

He denied any family history of sudden cardiac death or primary cardiomyopathy.

Physical examination

On physical examination at admission, his blood pressure was 105/81 mmHg, and his pulse rate was 92 beats/min. Additionally, an early diastolic gallop rhythm could be noticed at the cardiac apex.

Laboratory examinations

His myocardial enzymes were in the normal range, which ruled out myocarditis. No obvious abnormalities were found in routine blood tests, routine coagulation tests, liver function tests, kidney function tests, etc. Notable laboratory findings on presentation included high total cholesterol and triglyceride levels of 5.81 mmol/L and 4.72 mmol/L, respectively, and an elevated brain natriuretic peptide level of 502 pg/mL. Thyroid function tests revealed thyroid-stimulating hormone of 41.08 mIU/L (0.37-4.94), free triiodothyronine (T3) of 2.17 pmol/L (3.1-6.8), free thyroxine (T4) of 8.0 pmol/L (12-22), T3 of 0.79 nmol/L (1.35-3.15), T4 of 46.9 nmol/L (70-156), anti-

thyroglobulin antibody of 758.3 IU/mL (0-115), antithyroid peroxidase autoantibody of 215.8 IU/mL, and antithyroid microsome antibody of 69.6% (0-25).

Imaging examinations

His echocardiogram (**Figure 1A**) showed significant enlargement of the LV (diameter of 54 mm), severe LV global hypokinesis, minimal pericardial effusion and severe systolic dysfunction, with LVEF of 35% (**Table 1**). We did not see any valve damage. The electrocardiogram (Figure 1D) revealed obvious Q waves in leads III, aVF, and V1~V3. As coronary computed tomography angiography detected no coronary artery stenosis, we ruled out coronary heart disease. Cardiac magnetic resonance (CMR) (**Figure 2A**) also indicated an enlarged LV and decreased LVEF without abnormal myocardial perfusion of the left ventricular wall. Late gadolinium enhancement showed no obvious changes in the myocardium. Based on the above results, certain special types of cardiomyopathies were ruled out. Thyroid ultrasonography showed no diffuse mass.

FINAL DIAGNOSIS

After ruling out other possibilities, the patient was diagnosed with hypothyroid cardiomyopathy.

TREATMENT

The patient was given TH replacement therapy with levothyroxine 25 µg/day. He was given anti-heart failure treatment with furosemide 20 mg/day, spironolactone 25 mg/day, metoprolol 47.5 mg/day, and Entresto 50 mg/day twice a day. One week later, he had responded well and was discharged because of the resolution of clinical symptoms. He continued the treatment with the oral drugs.

OUTCOME AND FOLLOW-UP

One month later, the patient showed no obvious clinical symptoms, and the heart did not have murmurs or extra heart sounds. Thyroid function had returned to normal.

Echocardiogram (**Figure 1B**) showed the improvement of LVEF from 35% to 44%, and pericardial effusion had resolved compared with that during admission. At the 6-month follow-up appointment, he remained asymptomatic. Physical examination showed no obvious positive signs. A repeat echocardiogram (**Figure 1C**) revealed a diminishment of the enlarged LV chamber and a continuous increase in LVEF to 54% (**Table 1**). The follow-up CMR (**Figure 2B**) also showed that the dilated LV decreased, the overall systolic function recovered, and pericardial effusion decreased compared with the CMR findings during admission. These objective data corresponded to significant symptom alleviation and return to normal functional status.

DISCUSSION

Our patient had rapidly progressive HF secondary to hypothyroidism after radioactive iodine therapy. LV global hypokinesia and severely depressed systolic function were found on his echocardiogram. Although hypothyroidism is common after iodine-131 treatment, it is rare to see such a drastic change in cardiac function that may be related to rapid changes in thyroid hormones. This suggests that severe fluctuations in TH levels may lead to acute changes in the myocardium, eventually resulting in HF. The patient responded well to TH replacement therapy and had a gradual shrinkage of his enlarged LV chamber and markedly improved LV systolic function after treatment. He was diagnosed with hypothyroid cardiomyopathy because of the strong temporal relationship between thyroid hormone supplementation and his reversed symptoms and other exclusionary examinations. ⁵ To the best of our knowledge, there are no similar reports in the literature. Hypothyroid cardiomyopathy has been reported in previous cases, most of which were secondary to Hashimoto's thyroiditis [5], and the remaining cases were complications of Sheehan's syndrome or hypopituitarism [6, 7]. In addition, they were often the result of a long period of untreated hypothyroidism. However, our patient's HF symptoms developed rapidly, in less than 6 mo after radioactive iodine therapy. It is worth mentioning that all these symptoms were improved or even completely recovered by timely supplementation with thyroid

hormones. The present results are consistent with previous reports concerning the potential reversibility of HF after levothyroxine replacement, emphasizing the potential relationship between rapid changes in thyroid hormones and the development of HF.

Hypothyroidism can lead to HF. Previous animal experiments suggested that chronic hypothyroidism alone could produce dilated HF, characterized by impaired coronary blood flow and increased myocyte length/width ratio [8]. Low TH levels have also been independently associated with a worse prognosis in HF patients without a history of hypothyroidism [9]. A growing body of evidence suggests that a mutual cause-and-effect relationship might exist between HF and the hypothyroid state. Several cross-sectional case-control studies have shown that approximately 30% of patients with CHF have low T3 Levels and that T3 in serum decreases proportionately to the severity of heart disease [10]. Although controversial, some HF patients may benefit from TH supplementation [11]. Our case shows that a sudden decrease in TH can cause HF in a short time and can be recovered by TH replacement, which may provide a new perspective to guide the treatment of hypothyroidism with HF.

CONCLUSION

Herein, we introduced a case of a young male who rapidly developed hypothyroid cardiomyopathy secondary to radioactive iodine treatment, thus demonstrating the importance of periodic monitoring of cardiac function after iodine-131 treatment. Moreover, we showed the importance of timely supplementation with thyroid hormones, which may completely reverse the disease process.

6%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

link.springer.com
Internet

30 words — 2%
- 2

A. Martin Gerdes, Michael A. Portman, Giorgio Iervasi, Alessandro Pingitore, David KC Cooper, Dimitri Novitzky. "Ignoring a basic pathophysiological mechanism of heart failure progression will not make it go away", American Journal of Physiology-Heart and Circulatory Physiology, 2021
Crossref

19 words — 1%
- 3

www.wjgnet.com
Internet

15 words — 1%
- 4

Tomás F. Cianiulli, Luis A. Morita, María C. Saccheri, Marcelo Zylberman. "Hypothyroid cardiomyopathy: A reversible phenocopy of hypertrophic cardiomyopathy", Echocardiography, 2021
Crossref

13 words — 1%
- 5

www.ncbi.nlm.nih.gov
Internet

13 words — 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

EXCLUDE MATCHES

OFF

< 12 WORDS