

87657_Auto_Edited.docx

*Observational Study***Efficacy of β_2 -Adrenergic Receptor Agonist Combined with Corticosteroid in the Treatment of Children with Cough Variant Asthma**

Treatment of Variant Asthma in Children

Jun-Yi Cao, Ying-Chun Wang, Xiao-Xia Deng

Abstract**BACKGROUND**

Cough variant asthma is one of the most common respiratory diseases in children, which has a serious impact on the quality of life and daily activities of children. For severe cough variant asthma, immunomodulatory drugs are needed.

AIM

The aim of this study was to evaluate the efficacy of salmeterol combined with budesonide in the treatment of pediatric cough variant asthma (CVA).

METHODS

130 children with CVA from January 2020 to December 2022 were prospectively selected and randomly divided into an observation group (salmeterol combined with budesonide) and a control group (budesonide combined with a placebo). Compare the clinical efficacy of two groups before and after intervention. The evaluation parameters include cough frequency score, nocturnal cough arousal, and

lung function indicators. Serum inflammatory markers, immune function markers and airway anatomical indicators were also measured.

RESULTS

After the intervention, the total effective rate of the observation group was significantly higher than that of the control group, and the cough frequency score and the night cough wake rate of the observation group were lower than that of the control group, with a statistically significant difference. In addition, the changes of lung function indicators, serum markers and immune function markers in the observation group were better than those in the control group.

CONCLUSION

The clinical efficacy of salmeterol combined with Budesonide in the treatment of CVA is better than that of Budesonide alone.

Key Words: Salmeterol; Budesonide; CVA; Pediatrics; Efficacy analysis

Cao JY, Wang YC, Deng XX. Efficacy of β_2 -Adrenergic Receptor Agonist Combined with Corticosteroid in the Treatment of Children with Cough Variant Asthma. *World J Clin Cases* 2023; In press

Core Tip: Cough variant asthma seriously affects children's daily activities and physical health. For severe cough variant asthma, immunomodulatory drugs are needed. The purpose of this study is to evaluate the efficacy of salmeterol combined with budesonide in the treatment of cough variant asthma (CVA) in children. The results showed that after the intervention, the clinical symptoms of patients in the observation group were lighter and the level of immune markers was higher. The results of this study show that the combined treatment of the two drugs is better than budesonide alone.

INTRODUCTION

Cough Variant Asthma (CVA) is one of the most common airway diseases in children and is characterized by cough as the main symptom while other typical asthma symptoms may not be obvious^[1, 2]. The prevalence of CVA in children is increasing year by year, which has a serious impact on the quality of life and daily activities of children^[3, 4]. Currently, drugs used to treat CVA include inhaled corticosteroids (ICS), bronchodilators, leukotriene receptor antagonists, long-acting anticholinergics, and immunomodulators^[5-9]. ICS drugs such as beclomethasone propionate, fluticasone and budesonide are used to relieve coughs and asthma attacks by reducing airway inflammation and controlling symptoms. Bronchodilators relieve airway spasms and dilate the airway by relaxing the smooth muscle of the airway to relieve coughing and breathing difficulties, and the main bronchodilators include short-acting beta2-agonists such as salbutamol and long-acting beta2-agonists such as salmeterol. Leukotriene receptor antagonists block leukotriene receptors, reduce airway inflammation and relieve asthma symptoms. Long-acting anticholinergic drugs relieve airway spasm and improve lung function by blocking cholinergic receptors. For severe cough variant asthma, immunomodulatory drugs may need to be considered to regulate the immune system's overresponse and reduce airway inflammation^[10, 11].

As a common medical treatment, ICS are commonly used in the treatment of asthma to reduce airway inflammation and control symptoms^[12, 13]. However, some ICS have limited efficacy in children with CVA and may need to be combined with other medications to achieve better control^[14]. In recent studies, ICS in combination with long-acting beta2-agonists (LABA) such as salmeterol can provide a more effective treatment strategy^[15, 16]. Salmeterol acts as a LABA to relieve airway spasms by dilating airway smooth muscle^[17]. Budesonide is a potent ICS that reduces airway inflammation and controls asthma symptoms^[18]. However, comprehensive studies and evaluations on the efficacy of salmeterol combined with budesonide in children with cough-variant asthma are lacking. ¹⁸ The aim of this study was to systematically evaluate the efficacy of the

salmeterol, a LABA, combined with budesonide, an ICS, in the treatment of children with CVA, and to explore its effect on clinical efficacy and recovery indicators of childhood asthma control.

MATERIALS AND METHODS

Participants

This study used a prospective approach to select 130 children with CVA who were diagnosed and treated at our hospital between January 2020 and December 2022. The patients were randomly assigned (1:1) to observation group and control group using random number table method. The observation group received salmeterol combined with budesonide, while the control group received the budesonide combined with placebo. The baseline characteristics were balanced between the 2 groups. The comparison of basic data between the two groups before intervention showed that the difference was not statistically significant and was comparable (Table1). This study protocol was approved by the Ethics Committee of the first People's Hospital of Jiangxia District, Wuhan, and all the children's families have voluntarily participated in the study and have signed informed consent forms.

Inclusion and Exclusion Criteria

All the children met the diagnostic criteria of pediatric CVA^[1]. The main eligibility criteria included: 1) Cough: persistent or periodic cough, lasting at least 4 wk, and frequent attacks; 2) Variability: Cough changes significantly in response to different triggers, such as exercise, cold air, viral infection, or allergies; 3) Wheezing sounds: wheezing sounds can be heard during an asthma attack or when airflow is blocked; 4) Doctors make a comprehensive assessment based on the child's medical history, symptoms and signs; 5) Voluntarily participate in this study and cooperate with relevant examinations. The main exclusion criteria were 1) Complicated with serious heart, liver, kidney and other dysfunction; 2) Persistent cough caused by other causes, such as respiratory infections, chronic bronchitis, *etc.*; 3) Chronic cough induced by

mycoplasma infection; 4) Half-treatment withdrawal or late follow-up lost contact; 5) Allergic to the drug in this study.

Therapeutic method

On the basis of conventional treatment, children in the observation group were treated with Budesonide (Shanghai Xinyi Pharmaceutical Co., LTD., China, SFDA Approval No. H20010552) combined with Salmeterol aerosol (Glaxo Wellcome Production, French, Registration No. H20140382) for atomization inhalation. And the children in the control group were treated with atomized inhalation of Budesonide combined placebo. In the course of atomizing inhalation therapy, In the course of treatment, both drugs are used 1 press at a time, the oxygen flow rate is 6-8 L/min and the treatment is performed twice a day for 2 mo. During the treatment period, all children received anti-asthmatic, anti-infection and other conventional symptomatic treatment.

Observation Indicators

Clinical efficacy: According to the BA diagnostic criteria^[1], the clinical efficacy of the two groups was evaluated, including the following aspects: 1) The number of daytime symptoms; 2) The number of emergency relievers used; 3) Number of awakenings caused by nighttime asthma symptoms; 4) Exercise restriction due to asthma. The evaluation criteria are as follows: Obvious effect: Children's cough, wheezing and other symptoms disappeared, and lung function was significantly improved; Effective: The clinical symptoms were relieved and the lung function was improved. Ineffective: The symptoms and signs of the child were basically unchanged. The formula for calculating the total effective rate is: (obvious cases + effective cases)/ total number of cases ×100%.

Cough score: According to the Guidelines for the Diagnosis and Treatment of Cough (2021) ^[19], the cough symptoms of the two groups of children were evaluated, 0 points: no cough during the day or night; 1 point: Intermittent coughing during the day or waking up coughing once during the night.; 2 points: Cough intermittently or singly during the day or waking up coughing twice during the night.; 3 points: Paroxysmal

coughing during the day or waking up coughing more than three times during the night. A higher score referred to more serious cough.

Pulmonary function assay: Before and after treatment, ⁸ forced vital capacity (FVC), forced expiratory volume in one second (FEV1), peak expiratory flow (PEF) and maximal voluntary ventilation (MVV) of the two groups were measured by a pulmonary function tester (Zhejiang Yiliankang Medical Equipment Co., LTD, China, size: PF286), and the ratio of FEV1 to FVC was calculated.

Immunofunction assay: Before and after treatment, 5 mL fasting blood samples were taken from the peripheral veins of all patients. Serum was obtained from the samples after centrifugation, and CD3+, CD4+ and CD8+ cells were analyzed during the detection by flow cytometry of CytoFocus421 of Beijing Zhizhen Biotechnology Co., LTD.

Immunoglobulin assay: The blood samples were centrifuged by an automatic centrifuge (Microfuge 20/20R) with a centrifuge parameter set at 3000 RPM, a centrifuge radius of 10 cm, and a centrifuge time of 15 minutes. After centrifugation, the supernatant is taken and stored at -80°C. Subsequently, serum expression concentrations of immunoglobulin A (IgA) and immunoglobulin M (IgM) were measured using the Beckman Coulter AU5800 biochemical analyzer. The kit used is from Shanghai Huzhen Industrial Co., LTD.

Serum inflammatory indexes were measured: The venous blood collection method was the same as described above. After the blood samples were collected, the serum ¹² interleukin-6 (IL-6), interleukin-4 (IL-4) and tumor necrosis factor- α (TNF- α) concentrations were detected by ELISA.

Airway anatomical index: Lung scans were performed using GE Lightspeed spiral CT and Barc displays, ranging from the autonomic arterial arch to the bottom layer of the lung. The scanned images were saved and analyzed. To ensure the accuracy of the results, two experienced radiologists took independent measurements using blind methods and averaged them. In the process of measurement, the window width of 1500 Hu and window position of -450 Hu were used to select 5 CT levels at the end

of inspiratory breath of children with asthma. The image was enlarged on the display and lumen diameters were measured on segmental, subsegmental bronchus and transectional images. At the same time, the outer diameter and inner diameter of the airway cavity were measured. Finally, the airway wall thickness and airway wall area are calculated automatically by a calculation program.

7 Statistical Analysis

SPSS 20.0 statistical software was used to analyze the data. The counting data were expressed as [example (%)], and (2 test or Fisher exact probability method. The measurement data conforming to the normal distribution were expressed as ($\bar{x} \pm s$), and the comparison between the two groups was performed by t-test, and $P < 0.05$ was considered statistically significant.

RESULTS

Clinical data

Basic data of children in observation group and control group were collected, including age, sex, weight, course of disease, vitamin D consumption and season of onset. The observation group consisted of 35 males and 30 females. The average age of the patients was (7.86 ± 1.56) years, and the average course of disease was (7.6 ± 2.5) months. In the control group, there were 36 males and 29 females. The average age of the patients was (8.36 ± 1.28) years, and the average course of disease was (6.9 ± 3.2) months. There was no significant difference in the general data between the two groups ($P > 0.05$). (Table 1)

Comparison of clinical efficacy

The total effective efficacy rate in the observation group (96.92%) was higher than the control group (90.77%), with statistical difference ($P < 0.05$), as shown in Table 2. Indicating that the efficacy of Salmeterol combined with Budesonide in the treatment of children with cough variant asthma was higher than that of Budesonide alone.

Comparison of clinical cough score

After treatment, the score of cough frequency and the number of cough wake times in the two groups were decreased, and the difference was statistically significant ($P<0.05$). At the same time, the reduction rate of salmeterol combined with budesonide was significantly higher than that of budesonide alone, and the difference was statistically significant ($P<0.05$). (Table 3)

Comparison of pulmonary function

The post-treatment MVV, PEF, FEV1 and FEV1/FVC were higher than those pretreatment in both groups ($P<0.05$). Moreover, the post-treatment MVV, PEF, FEV1 and FEV1/FVC in the observation group were higher than those in the control group ($P<0.05$) (Table 4). The results showed that salmeterol combined with budesonide improved lung function better than budesonide alone.

Comparison of inflammatory response

The serum levels of IL-4, IL-6 and TNF- α were compared between the two groups before and after treatment, and there was no statistical difference between the observation group and the control group before treatment ($P>0.05$). The post-treatment IL-6, IL-4 and TNF- α levels were lower than those pre-treatment in both groups ($P<0.05$), and post-treatment IL-6, IL-4 and TNF- α levels in the observation group were lower than those in the control group ($P<0.05$) (Table 5). These results indicated that Salmeterol combined with Budesonide was more effective in improving inflammatory response than Budesonide alone.

Comparison of T lymphocyte count

Before treatment, the levels of CD3+, CD4+ and CD8+ cells count between the observation group and the control group has no statistical difference ($P>0.05$). After treatment, the CD3+ and CD4+ cells count in both groups were higher than before treatment ($P<0.05$), while the CD8+ cells count were lower than before treatment ($P<$

0.05). Also, the post-treatment CD3⁺ and CD4⁺ cells count in observation group was higher, while CD8⁺ cells count was lower in the observation group than those in the control group ($P<0.05$) (Table 6).

Comparison of immunoglobulin degree

The post-treatment serum IgA and IgM levels were higher than those before treatment in both groups ($P<0.05$), and the post-treatment levels in the observation group were higher than those in the control group ($P<0.05$) (Table 7).

Comparison of Airway anatomical index

Airway wall thickness and total airway wall area were compared between the two groups before and after treatment, and there was no significant difference between the two group before treatment ($P>0.05$). After treatment, airway wall thickness and total area decreased in both groups ($P<0.05$), and the above parameters in the observation group were lower than those in the control group ($P<0.05$). (Table 8)

DISCUSSION

Cough variant asthma is a common type of childhood asthma characterized by a chronic cough with symptoms that fluctuate wildly across time and environment^[6]. Salmeterol, a long-acting beta-2 adrenergic receptor agonist, is a commonly used asthma medication that reduces airway spasms by dilating bronchial smooth muscles, thereby relieving breathing difficulties and coughing symptoms^[20]. The use of salmeterol has been widely discussed in children with cough variant asthma. Studies have shown that salmeterol is effective in reducing cough frequency and duration and improving respiratory symptoms in children. On the other hand, the corticosteroid budesonide is widely used as an anti-inflammatory drug in the treatment of asthma^[21]. It can inhibit airway inflammation, reduce airway edema and excessive mucus, and thus improve airway patency. The use of budesonide has also been extensively studied in children with cough-variant asthma. Studies have shown that budesonide can

significantly improve lung function and reduce inflammatory response in children^[22, 23]. In addition, budesonide improves immune function by increasing CD4⁺ cell count and immunoglobulin levels^[18].

In this study, we observed the efficacy of the β 2-receptor agonist salmeterol in combination with the corticosteroid budesonide in the treatment of children with cough-variant asthma. In terms of clinical efficacy, we found that the combination regimen significantly reduced cough frequency and duration, improved the overall response rate of treatment, and improved the life quality of children. In terms of lung function, the children's PEF and FEV1 were significantly improved. In addition, the serum inflammatory indicators IL-6, TNF- α , IL-4 were significantly reduced, and the improvement of immune function included the increase of the number of CD4⁺, CD3⁺ cells and the levels of immunoglobulin IgA and IgM.

CONCLUSION

In summary, the β 2-receptor agonist salmeterol combined with the corticosteroid budesonide is an effective regimen for the treatment of children with cough-variant asthma. The combination therapy can significantly improve clinical symptoms, increase lung function, reduce inflammation levels, and improve immune function.

These findings have important clinical implications for clinicians in the treatment of children with cough-variant asthma, but this study has some limitations, such as a small sample size and a short study period. Therefore, future studies can further expand the sample size and extend the observation time to further validate our findings.

ARTICLE HIGHLIGHTS

Research background

The combination of ICS and long-acting β 2 receptor agonist (LABA), such as salmeterol, can provide a more effective treatment strategy, but there is a lack of comprehensive research and evaluation on the efficacy of salmeterol combined with budesonide in the treatment of cough variant asthma in children.

Research motivation

To systematically explore the efficacy of LABA salmeterol combined with ICS budesonide in the treatment of cough variant asthma in children, and to provide some clinical treatment suggestions for the treatment and clinical recovery of cough variant asthma in children.

9

Research objectives

The purpose of this study is to systematically evaluate the efficacy of LABA salmeterol combined with ICS budesonide in the treatment of CVA in children, and to explore the influence of the combination of the two drugs on the clinical efficacy and recovery index of asthma control in children.

Research methods

17

Children with CVA were selected prospectively and randomly divided into observation group (salmeterol combined with budesonide) and control group (budesonide combined with placebo). The clinical efficacy and immune function markers of the two groups were compared.

16

Research results

After treatment, the effect and efficiency of the two groups were improved to some extent. The symptoms of CVA in the treatment group were relieved, and the lung function index and immune marker level were also better than those in the control group.

Research conclusions

Both salmeterol combined with budesonide and budesonide alone can treat CVA to a certain extent, but the therapeutic effect of the former is significantly higher than that of the latter in symptom improvement, immune marker level and lung function level.

Research perspectives

Salmeterol combined with budesonide is an effective method to treat CVA disease, which is worth popularizing in clinic.

ACKNOWLEDGEMENTS

we are grateful to all patients for their participation.

17%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|----------------|
| 1 | e-century.us
Internet | 146 words — 5% |
| 2 | www.ncbi.nlm.nih.gov
Internet | 89 words — 3% |
| 3 | downloads.hindawi.com
Internet | 34 words — 1% |
| 4 | journal07.magtech.org.cn
Internet | 34 words — 1% |
| 5 | www.ijpsonline.com
Internet | 29 words — 1% |
| 6 | Chaofan Zeng, Ying Huang, Longer Yu, Qingmei Zeng, Bijun Wang, Yi Xu. "Long-Term Assessment of Rehabilitation Treatment of Sports through Artificial Intelligence Research", Computational and Mathematical Methods in Medicine, 2021
Crossref | 22 words — 1% |
| 7 | Baolin Ma, Huping Song, Wuyong Lin. "Efficacy and prognosis of neuroendoscopy-assisted surgery for chronic subdural hematoma", Pakistan Journal of Medical Sciences, 2023
Crossref | 19 words — 1% |

8	nlist.inflibnet.ac.in Internet	19 words — 1%
9	www.wjgnet.com Internet	17 words — 1%
10	Shuang Zhou, Lai-Run Jin, Chiyi He. "Effects of imipenem combined with glutamine in the treatment of severe acute pancreatitis with abdominal infection in mainland China: a meta-analysis", Revista da Associação Médica Brasileira, 2022 Crossref	16 words — 1%
11	f6publishing.blob.core.windows.net Internet	15 words — < 1%
12	manu41.magtech.com.cn Internet	15 words — < 1%
13	www.ijo.cn Internet	15 words — < 1%
14	www.openaccessjournals.com Internet	15 words — < 1%
15	www.spandidos-publications.com Internet	14 words — < 1%
16	"Abstracts", Asia-Pacific Journal of Clinical Oncology, 2022 Crossref	13 words — < 1%
17	www.biolifesas.org Internet	13 words — < 1%
18	worldwidescience.org Internet	12 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS

EXCLUDE MATCHES < 12 WORDS