

87395_Auto_Edited.docx

Adult localized Langerhans cell histiocytosis: A case report

Yang *et al.* (Adult localized LCH)

Pan-Pan Yang, Su-Ye Hu, Xu-Ya Chai, Xiao-Meng Shi, Li-Xia Liu, Ling-E Li

Abstract**BACKGROUND**

Langerhans cell histiocytosis (LCH) is a rare clonal proliferative disease of Langerhans cells with unknown pathogenesis. An increasing number of clinicians recognize that LCH has a wide clinical spectrum and a highly varied course. Adults rarely develop LCH. Here, we report a case of adult localized LCH.

CASE SUMMARY

A 32-year-old woman presented with plaques and ulcers on the vulva and crissum, accompanied by pain that persisted for more than one year. Physical examination revealed a red-infiltrating plaque with ulcerations and exudates in the vulva and crissum. Pathological examination revealed a diffuse infiltration of lymphocytes, eosinophilic granulocytes, and histiocytoid cells in the superficial dermis. Proliferative histiocytoid cells showed mild atypia, partly with kidney-shaped nuclei. Immunohistochemical examination showed that the histiocytoid cells were positive for S100 protein and CD1 antigen and weakly positive for CD68 (20%+) antigen and Ki-67 index (30%+). Laboratory tests did not reveal any other systemic damages. The patient was diagnosed with adult localized LCH and was prescribed oral prednisone (20 mg) once daily. The skin lesions gradually improved and are still being followed-up.

CONCLUSION

Adult localized LCH is rare and must be differentiated from other common conditions.

Key Words: Langerhans cell histiocytosis; Adult; Vulva; Crissum; S100; Case report

Yang PP, Hu SY, Chai XY, Shi XM, Liu LX, Li LE. Adult localized Langerhans cell histiocytosis: a case report and review of literature. *World J Clin Cases* 2023; In press

Core Tip: Langerhans cell histiocytosis (LCH) is a rare clonal proliferative disease of Langerhans cells. Adult localized LCH presented with plaques and ulcers on the vulva and crissum is rare and must be differentiated from other common conditions.

INTRODUCTION

Langerhans cell histiocytosis (LCH) is a disease spectrum characterized by the proliferation of Langerhans cells that express an immunophenotype positive for S100 protein, CD1 antigen (CD1a), and Langerin (CD207) and contains cytoplasmic Birbeck granules^[1]. Several organs are involved in this process. The childhood and adult forms of LCH should be considered separately. Two-thirds or more children under one year of age and those aging 1–4 years have multi-system diseases, including those of the liver, lungs, or bone marrow. In adults, the peak age of presentation is between 20 years and 35 years, with multi-system diseases occurring in one-thirds to two-thirds of adults with LCH. The bones are the most commonly involved organs. Skin and mucosal involvement are the second most common manifestations in adults^[2]. Here, we report a case of adult LCH that presented solely with skin involvement, without systemic damage.

CASE PRESENTATION

Chief complaints

A 32-year-old female presented with plaques and ulcers on the vulva and crissum, accompanied by pain and exudates on the surface of the ulcers.

History of present illness

The patient reported a rash on the vulva as the first manifestation, which gradually enlarged and extended to the perianal area. She did not complain of diarrhea, constipation, melena, hematochezia, diabetes insipidus, changes in appetite, or weight loss.

History of past illness

The patient denied any previous chronic diseases, hepatitis, ² tuberculosis and other infectious disease history and history of close contact. Deny drug, food allergies. Vaccination history is unknown.

⁷

Personal and family history

The patient had no family history of cancer.

Physical examination

Physical examination revealed red erosive plaques and ulcers on the vulva and perianal areas, which were covered with exudates (Figures 1). No skull defects were observed.

Laboratory examinations

A punch biopsy was performed at the intersection between the normal skin and vulvar rash during the initial visit. ⁵ The tissue section was fixed in 10% neutral formalin, paraffin-embedded, sectioned, stained with hematoxylin and eosin staining and subjected to direct immunohistochemistry before observation under light microscopy. Each immunohistochemistry test group included a negative self-control. All primary and secondary antibodies were purchased from ZSGB-BIO, Beijing, China. The primary antibody incubation lasted 50 minutes at 37°C and 20 minutes at 37°C for the secondary

antibody. Visualization was performed using DAB(3,3'-diaminobiphenyl), and results were assessed by capturing images with a light microscope.

Pathology revealed histiocytoid cells in the superficial and middle dermal layers with mild nuclear atypia and a population of occasional eosinophils. The atypical histiocytoid cells were positive for S100 protein and CD1a, and weakly positive for CD68 (20%+) antigen, with a Ki-67 index of 30%. These cells were negative for creatine kinase (Figures 2,3).

Imaging examinations

Computed tomography of the sacroiliac joint showed that the cortical bone of the sacral surface of the right sacroiliac joint was discontinuous.

FINAL DIAGNOSIS

Based on these results, the patient was diagnosed with adult localized Langerhans cell histiocytosis.

TREATMENT

The patient was prescribed thalidomide 0.1 g orally twice daily and methotrexate (7.5 mg) once weekly.

OUTCOME AND FOLLOW-UP

Three months later, the patient reported a poor response and was prescribed prednisone 20 mg orally once daily; the lesions gradually improved and the patient is still being followed up (Figure 1).

DISCUSSION

LCH is a rare group of disorders in which mononuclear macrophages and dendritic cell systems proliferate, often affecting the skin and bones. It can also lead to multi-system diseases of the liver, spleen, lung, central nervous system, lymph nodes, thymus,

gastrointestinal tract, and bone marrow^[1-5]. Liver and spleen involvement causes abnormal liver function, hepatomegaly, and hypersplenism^[6]. Involvement of the lungs may cause chest pain and dyspnea^[7]. The pituitary gland may also be affected by LCH causing diabetes insipidus^[8-10]. The bones of the skull are the most common sites affected by LCH, but other bones, such as the femur, scapula, rib, mandible, and vertebrae, can also be affected^[11-13].

Depending on the number of systems involved in the patient, ¹ LCH is divided into single-system-LCH (SS-LCH) and multi-system-LCH (MS-LCH)^[14]. SS-LCH is divided into SS-S (a single site involving the bone, skin, or lymph nodes) and SS-M (involving multiple parts of the bone or lymph nodes). Patients with SS-LCH have a good prognosis, whereas those with MS-LCH, especially those with the liver and hematopoietic system involvement, have a poor prognosis and a higher mortality rate^[6, 15]. Generally, patients with LCH whose only skin is affected have a good prognosis and about 50% of the patients can be in remission within a few months; however, disease progression and persistence are more common, and long-term follow-up is recommended^[16].

The most common areas of skin involvement in LCH are the scalp, trunk, skin folds, and mucosa. Skin lesions vary and present as papules, blisters, pustules, purpura, plaques, or ulcers^[2]. The most characteristic presentations of adult LCH are groin, perianal, and vulvar involvements. Chen *et al.*^[17] reported a case of adult LCH with an eczematoid lesion in the vulva as the initial manifestation, whereas Wu *et al.*^[18] reported a case of LCH with perianal skin lesions as the first presentation. In the present case, the skin lesions were located on both the vulvar and perianal regions. Computed tomography of the sacroiliac joint revealed discontinuity in the cortical bone of the right sacroiliac joint. Changes in the sacroiliac joint necessitated further follow-up and progressed slowly with no apparent systemic involvement. In conclusion, adult LCH is a rare disease that must be distinguished from other common diseases, including Paget's disease, candidiasis, and malignant melanoma.

Treatment of LCH should be individualized according to the number and severity of organs involved. Therapeutic options should also prevent long-term side effects of medication. Topical and systemic corticosteroids, nitrogen mustard, methotrexate, psoralen plus ultraviolet-A radiation, narrow-band ultraviolet-B radiation (or excimer laser), thalidomide, interferon, and azathioprine can be considered for treatment depending on the extent of skin lesions^[2]. Therefore, surgical resection of a single lesion should be considered. No single treatment has been effective in all patients.

CONCLUSION

The rarity of LCH in adults may have resulted in its overlooked diagnosis. Therefore, dermatologists should consider this disease, along with its varied presentation and treatment options.

9%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.jsltr.org Internet	22 words — 2%
2	www.barnes-pump.com Internet	19 words — 1%
3	Grace Tan. "Langerhans cell histiocytosis of the cervical spine: a single institution experience in four patients", Journal of Pediatric Orthopaedics Part B, 03/2004 Crossref	14 words — 1%
4	f6publishing.blob.core.windows.net Internet	14 words — 1%
5	mafiadoc.com Internet	13 words — 1%
6	www.bib.irb.hr Internet	13 words — 1%
7	rcastoragev2.blob.core.windows.net Internet	12 words — 1%
8	www.wjgnet.com Internet	12 words — 1%

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 12 WORDS
EXCLUDE BIBLIOGRAPHY	ON	EXCLUDE MATCHES	< 12 WORDS