

81276_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 81276

Manuscript Type: CASE REPORT

Testicular pain originating from lumbar disc degeneration: A case report

discogenic testicular pain

Bing Wu, Xiujie Yan, Xin He, Zikai Tian, Bao-Gan Peng

Abstract

BACKGROUND

Testicular pain caused by lumbar disease is uncommon in clinic. Here we reported a case of discogenic low back pain with testicular pain which was successfully cured.

CASE SUMMARY

A 23-year-old male patient presented to our department with chronic low back pain. Based on his clinical symptoms, signs, and imaging, he was diagnosed with discogenic low back pain. Since conservative treatment for more than half a year did not significantly improve his low back pain, we decided to treat it with intradiscal methylene blue injection. During the course of surgery, we again identified the low back pain as originating from the degenerated lumbar disc by analgesic discography. Interestingly, the patient's low back pain disappeared along with the testicular pain that had been present for more than 3 mo. After the operation the patient's low back pain improved obviously and the testicular pain did not reappear anymore.

CONCLUSION

Intradiscal methylene blue injection is a convenient and effective surgical intervention for the treatment of discogenic low back pain; Lumbar disc degeneration may also be a possible clinical cause of testicular pain. With methylene blue working in the diseased disc, the symptom of low back pain was improved and the accompanying testicular pain was also successfully managed.

Key Words: Disc degeneration; Testicular pain; Low back pain

Wu B, Yan X, He X, Tian Z, Peng BG. Testicular pain originating from lumbar disc degeneration: A case report. *World J Clin Cases* 2023; In press

Core Tip: Herein, we present a case of a patient who was cured of testicular pain while treating discogenic low back pain, which means the lumbar disc degeneration may be a possible etiology of testicular pain.

INTRODUCTION

Chronic testicular pain is one of the common symptoms in urology ^[1]. Clinicians have found numerous causes that can evoke testicular pain, including infections, tumors, varicocele, and referred pain. However, lots of patients still suffer from testicular pain of unknown pathogenesis, and could not be effectively treated so far ^[2]. Lumbar disc degeneration has been proven to be a major source of low back pain, and some researchers have proposed that lumbar disc herniation could also lead to testicular pain ^[3]. Nevertheless, there has been no high-quality study on the testicular pain caused by lumbar disc degeneration. The relationship between lumbar disc degeneration and testicular pain or the pathogenesis of discogenic testicular pain remains elusive. In this paper, we report a young man whose testicular pain was cured while treating discogenic low back pain, and analyze the possible pathogenesis of testicular pain induced by lumbar disc degeneration.

1

CASE PRESENTATION

Chief complaints

A 23-year-old man had a 1-year history of severe low back pain, which seriously affected his quality of life.

History of present illness

Prior to our department, the patient had been on non-steroidal anti-inflammatory drugs (NSAIDs) and physical therapy for more than eight months. However, the low back pain did not improve but instead continued worsening.

History of past illness

The patient had also suffered from bilateral testicular pain for more than three months. After detailed physical examination and related ultrasonography, urologists thought that testicular pain does not seem to come from urinary system diseases. No other illnesses were documented and the patient had no relevant traumatic history.

Personal and family history

It was unremarkable.

Physical examination

The pressure pain of L_{4/5} intervertebral space was positive. No abnormal sensory or motor function of both lower extremities. The scrotum is normal, and testiculus are in the scrotum without enlargement. The right testis is slightly tender. Visual analogue scale (VAS) score of low back pain was 8 and the testicular pain was 7.

Laboratory examinations

It was unremarkable.

Imaging examinations

The height of L_{4/5} intervertebral space was lower in the radiographs of lumbar spine (Figure 1); Sagittal (left) and axial (right) T₂-weighted MRI of the lumbar spine images showed obvious degeneration of L_{4/5} disc (Figure 2).

FINAL DIAGNOSIS

Discogenic low back pain with testicular pain.

TREATMENT

The procedure was performed under fluoroscopy and we used a standard posterolateral approach and a double needle technique [4]. We conducted the discography for the L_{3/4} disc and the L_{4/5} disc respectively. About 0.2 mL of 2%

lidocaine hydrochloride was slowly injected into the nucleus pulposus after the needle was accurately inserted into the center of the disc. At the time of performing L_{3/4} disc, there was no change in low back pain or testicular pain. In contrast, when the L_{4/5} disc was performed, the patient clearly demonstrated a significant improvement in low back pain, at almost the same time, testicular pain disappeared. Subsequently, about 0.3 mL of 1% methylene blue was injected into the diseased disc. The patient was required to rest on bed for next 24 h after surgery.

OUTCOME AND FOLLOW-UP

The patient's bilateral testicular pain quickly disappeared after the analgesic discography. Testicular pain VAS score: 0; Low back pain VAS score: 2.

At 1 year follow-up, the patient's testicular pain did not recur and the patient's low back pain improved significantly.

DISCUSSION

Chronic testicular pain originating from lumbar disc degeneration is very rare in clinical. This may due to the insufficient awareness of this disease or the lack of a clear diagnostic method. In 2003, Kathryn *et al* reported a case who had central low back symptoms and right testicular pain with a small central disc protrusion between the T12 and L1 vertebrae. They successfully eased the symptoms by physical therapy 12 times during 3 mo [5]. Peng has also reported that degenerative lumbar spondylolisthesis can cause testicular pain, and suggested that this type of testicular pain is an involvement pain arising from lumbar spine disease [6].

Bogduk has showed the innervation of cervical and lumbar intervertebral discs, which receive ²innervation posteriorly from the sinuvetebral nerves, laterally from the vertebral nerve, and anteriorly from the sympathetic trunks [7]. Furthermore, the innervation of cervical and lumbar discs is multi-segmental [8,9]. For example, low back pain caused by L_{4/5} or L₅/S₁ disc degeneration is not transmitted upward *via* the L₅ or S₁ spinal nerves

as traditionally believed, but is transmitted through the paravertebral sympathetic chain to the dorsal root ganglia of L₁ and L₂ and then to the center [6]. In addition, the distribution of nerves in the lumbar disc is consistent with the distribution of nerves in the testis (genitofemoral nerve and genital branch of the ilioinguinal nerve), mainly originating from the L₁ and L₂ nerve roots [9]. Such neural pathways provide the basis for testicular pain induced by intervertebral disc degeneration.

Lumbar discography and methylene blue injection are common clinical methods for the diagnosis and treatment of discogenic lower back pain. The vascularized granulation tissue and the tissue repair process of injurious nerve fibers growing into the degenerated disc are important pathophysiological mechanisms of discogenic lower back pain [10]. Methylene blue as a nerve inactivator can inhibit the nociceptive nerves growing into the lumbar disc, eventually making the pain disappear [10]. In combination with the patient in this case, the imaging manifestations and positive signs were typical signs of discogenic low back pain, and the analgesic discography improved both low back pain and testicular pain. After intradiscal methylene blue injection, the low back pain improved again along with the disappeared testicular pain, suggesting that the testicular pain possible originated from the lumbar disc degeneration rather than the testis itself. The mechanism of discogenic testicular pain may therefore originate from the pain induced by the degenerated L_{4/5} disc non-segmentally transmitted through the paravertebral sympathetic chain to the L₁ and L₂ spinal nerve roots and then upward in conjunction with visceral sympathetic afferent nerves transmitted in the L₁ and L₂ spinal nerve roots [11].

In the present case, the patient's diagnosis and treatment were based on discogenic low back pain rather than specifically on testicular pain. Although lumbar analgesic discography suppressed both low back pain and testicular pain, this does not mean that lumbar analgesic discography is an effective method for diagnosed testicular pain. However, the disappearance of testicular pain in the patient after the procedure may further support that lumbar disc degeneration is a significantly clinical resource of

testicular

pain.

CONCLUSION

In this case, we found that disc degeneration, as the etiology of low back pain, can also evoke the concomitant testicular pain. Analgesic discography can be chosen as an effective diagnostic method, and intradiscal methylene blue injection can significantly improve low back pain and testicular pain.

ORIGINALITY REPORT

8%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.wjgnet.com Internet	52 words — 4%
2	www.ncbi.nlm.nih.gov Internet	17 words — 1%
3	academicjournals.org Internet	15 words — 1%
4	Baogan Peng, Duanming Li, Xiaodong Pang. "Degenerative Lumbar Spondylolisthesis with Testicular Pain", Pain Medicine, 2014 Crossref	13 words — 1%
5	Baogan Peng, Xiaodong Pang, Ye Wu, Changcheng Zhao, Xinghua Song. "A randomized placebo- controlled trial of intradiscal methylene blue injection for the treatment of chronic discogenic low back pain", Pain, 2010 Crossref	13 words — 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

EXCLUDE MATCHES

< 12 WORDS

< 12 WORDS