

90657_Auto_Edited.docx

Hematuria after nocturnal exercise of a man: A case report

Mingjian Bai, Songtao Yang, Xuekai Liu

Abstract**BACKGROUND**

A man experienced multiple episodes of macroscopic hematuria following nocturnal exercise. Urinary stones and tumors were considered the two most likely causes. The patient had two hobbies: consuming health care products in large quantities and engaging in late-night running.

CASE SUMMARY

Health care products contain a large amount of calcium phosphate, and we hypothesize that this could induce the formation of small phosphate stones. After exercise, the urinary system is abraded, resulting in bleeding. The patient was advised to stop using the health care products. Consequently, the aforementioned symptoms disappeared immediately. However, the patient resumed the above two habits one year later; correspondingly, the macroscopic hematuria reappeared.

CONCLUSION

This finding further confirmed the above inference and allowed for a new avenue to determine the cause of the patient's hematuria.

INTRODUCTION

Multiple studies have shown that clinical hematuria is common after exercise^[1], with 95-100% of the patients with clinical hematuria having it after exercise^[2]. The degree of hematuria is related to the intensity of the exercise; for instance, contact sports can increase the risk of macroscopic hematuria. Exercise-related urological trauma is regarded as the leading cause of macroscopic hematuria, of which renal trauma accounts for 80% of cases. Here, we present the case of a patient with macroscopic hematuria who consumed a significant amount of health care products orally and favored exercise.

CASE PRESENTATION

Chief complaints

2

A 42-year-old man was admitted to the Emergency Department of our hospital on June 14, 2020 due to macroscopic hematuria after nocturnal exercise (Figure 1).

History of present illness

Upon reviewing the medical history, the patient experienced macroscopic hematuria three times after nocturnal exercise in the previous four years, and the hematuria was accompanied by lower abdominal pain. However, the actual cause of the disease had not been determined. Following a multidisciplinary team consultation, it was speculated that the nutcracker phenomenon, the very small stone traumatizing the mucosa, and urinary tract tumors were the three most likely causes of the patient's hematuria.

1

History of past illness

none

Personal and family history

none

Physical examination

none

Laboratory examinations

We conducted routine urine tests, Doppler ultrasound, CT urography, and endoscopic examination of the urinary system. The results indicated that only the routine urine test was abnormal, revealing urine protein (+), occult blood (++), and a red blood cell count of 21,587.0 cells/ μ l. Additionally, a significant number of amorphous phosphate crystals were observed under the microscope (Figure 1).

Imaging examinations

No obvious compression changes were observed in the left renal vein. The angle between the superior mesenteric artery and abdominal aorta was 41 degrees (Figure 2). At this point, hematuria caused by the nutcracker phenomenon and urologic neoplasms were essentially ruled out.

MULTIDISCIPLINARY EXPERT CONSULTATION

We re-inquired the patient again, with a focus on his lifestyle habits. The patient particularly enjoyed doing two things: taking large doses of health care products and exercising late at night. Upon reviewing the labels of the health care products, we identified calcium phosphate as one of the main components.

FINAL DIAGNOSIS

A hypothesis was formulated: the patient had orally consumed excessive amounts of health care products containing calcium phosphate, leading to the formation of tiny stones in the urine. These stones are not easily detected by imaging or endoscopy. Following nocturnal exercise, these tiny stones scratch the mucosa of the urinary system, resulting in bleeding and abdominal pain.

TREATMENT

Based on the above assumptions, the patient was advised to stop using the health care products after discharge.

OUTCOME AND FOLLOW-UP

Subsequently, the patient's hematuria disappeared. However, one year later, the patient consumed the aforementioned health care products in large quantities again, and his hematuria recurred after late-night exercise. It was then confirmed that the patient's hematuria was caused by the consumption of health care products containing calcium phosphate. The patient was again advised to stop using those health care products, at which point the patient's hematuria then disappeared.

DISCUSSION

Hematuria is a common phenomenon following exercise^[1]. A previous study provided evidence of the relationship between hypercalciuria and postglomerular hematuria in children^[3]. Escribano *et al* also discovered that hypercalciuria can lead to recurrent macroscopic or microscopic hematuria^[4]. Through follow-up, it was observed that the occurrence of hematuria after exercise was clearly associated with the patient's use of the aforementioned health care products. Consequently, we deduced the pathogenesis of this patient: the abundance of amorphous phosphate leads to the formation of crystals in the urine. These crystals may scratch the urethra during exercise, ultimately resulting in hematuria.

CONCLUSION

In conclusion, if patients consume a significant quantity of health care products containing calcium phosphate, amorphous phosphate crystals may appear in the urine. During exercise, these crystals may scratch the urethra, leading to hematuria. Therefore,

the present case provides a new avenue for determining the cause of hematuria in clinical practice.

5%

SIMILARITY INDEX

PRIMARY SOURCES

1	f6publishing.blob.core.windows.net Internet	27 words — 3%
2	doctiktak.com Internet	14 words — 2%