

Name of Journal: *World Journal of Clinical Cases*

ESPS Manuscript NO: 16875

Manuscript Type: Original Article

Retrospective Study

Improved bowel preparation increases polyp detection and unmasks significant polyp miss rate

Ioannis S Papanikolaou, Athanasios D Sioulas, Nektarios Magdalinos, Iosif Beintaris, Lazaros-Dimitrios Lazaridis, Dimitrios Polymeros, Chrysoula Malli, George D Dimitriadis, Konstantinos Triantafyllou

Match Overview

1	CrossCheck 126 words Hong, Sung Noh, In Kyung Sung, Jeong Hwan Kim, Won Hyeok Choe, Byung Keek Kim, Soon Young Ko, Jung Hyun Lee	3%
2	CrossCheck 111 words Sang Bong Ahn, "The Miss Rate for Colorectal Adenoma Determined by Quality-Adjusted, Back-to-Back Colonoscopies"	2%
3	CrossCheck 46 words Boroff, Erika S, Suryakanth R Gurudu, Joseph G Hertz, Jonathan A Leighton, and Francisco C Ramirez, "Polyp and Adei	1%
4	CrossCheck 30 words "UEG Week 2013 Poster Presentations", United European Gastroenterology Journal, 2013.	1%
5	Internet 20 words crawled on 28-Sep-2012 www.labdominguez.com.ar	<1%
6	Internet 10 words crawled on 17-Sep-2013 www.toibg.com	<1%

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 66,600 条结果 (用时 0.88 秒)

Google 学术: Improved bowel preparation increases polyp detection and unmasks significant polyp miss rate

Recognition and characterization of small colonic ... - Hoffman - 被引用次数: 90

... for adenoma and hyperplastic polyp detection with ... - East - 被引用次数: 76

Advances in endoscopic imaging of colorectal ... - Wallace - 被引用次数: 75

Inadequate Bowel Preparation Increases Missed Polyps

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: HW Kim - 2012 - 相关文章

2012年11月30日 - Inadequate Bowel Preparation Increases Missed Polyps ... polyps and that colonoscopic polypectomy significantly reduces the ... having reached the current target adenoma detection rate (42% and 27%, 47.9% and 18%, respectively). ... associated with an increased risk of missed polyp, missed adenoma, ...

A simple method to improve adenoma detection rate during ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: AŞ Köksal - 2013 - 被引用次数: 12 - 相关文章

However, tandem colonoscopy studies reveal a pooled polyp miss rate of 22%. ... The increase in the number of adenomas detected was statistically significant in ... increasing the adequacy of the bowel preparation with the use of split-dose ...

The Miss Rate for Colorectal Adenoma Determined by ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: SB Ahn - 2012 - 被引用次数: 37 - 相关文章

2012年1月12日 - The smaller polyps and increased number of polyps detected during the first ... There was a significant miss rate in the detection of colonic ... The colonoscopic withdrawal time, quality of bowel preparation, and optimal

[网页](#) [图片](#) [新闻](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 66,800 条结果 (用时 0.68 秒)

Google 学术 : Improved bowel preparation increases polyp detection and unmasks significant polyp miss rate

Recognition and characterization of small colonic ... - Hoffman - 被引用次数 : 90

... for adenoma and hyperplastic polyp detection with ... - East - 被引用次数 : 76

Advances in endoscopic imaging of colorectal ... - Wallace - 被引用次数 : 75

Inadequate Bowel Preparation Increases Missed Polyps

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)作者 : HW Kim - 2012 - [相关文章](#)

2012年11月30日 - Inadequate Bowel Preparation Increases Missed Polyps ... having reached the current target adenoma detection rate (42% and 27%, 47.9% and 18%, respectively). ... polyps and adenomas during screening colonoscopy is significantly ... associated with an increased risk of missed polyp, missed adenoma, ...

A simple method to improve adenoma detection rate during ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)作者 : AŞ Köksal - 2013 - 被引用次数 : 12 - [相关文章](#)

However, tandem colonoscopy studies reveal a pooled polyp miss rate of 22%. ... during colonoscopy withdrawal increases the adenoma detection rate (ADR). ... The increase in the number of adenomas detected was statistically significant in the ... All patients received a standard bowel preparation of 4 L of polyethylene ...

The Miss Rate for Colorectal Adenoma Determined by ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)作者 : SB Ahn - 2012 - 被引用次数 : 37 - [相关文章](#)

2012年1月12日 - The smaller polyps and increased number of polyps detected during the first ... There was a significant miss rate in the detection of colonic ... The colonoscopic withdrawal time, quality of bowel preparation, and optimal