

Match Overview

Name of Journal: World Journal of Clinical Cases

ESPS Manuscript NO: 15924

Column: CASE REPORT

Intestinal type gastric adenocarcinoma with unusual synchronous metastases to the colorectum and bladder

Isaac Seow-En, Francis Seow-Choen

Abstract

A 75 year old male presented with difficult defecation and increasing urinary frequency over a few months. He had a significant history of previous partial gastrectomy for gastric carcinoma 20 years prior. CT of the abdomen and pelvis showed extensive lymphadenopathy, a gastric mass and rectal as well as bladder wall thickening with bilateral ureterohydronephrosis. Normal looking serosal surfaces of the bladder and bowel were seen on laparoscopy and a defunctioning



Intestinal type gastric adenocarcinoma with unusual synchronous metas



网页

图片

新闻

购物

视频

更多

搜索工具

找到约 609,000 条结果 (用时 0.64 秒)

您是不是要找: Intestinal type gastric adenocarcinoma with unusual synchronous metastases to the *colorectal* and bladder

Google 学术: Intestinal type gastric adenocarcinoma with unusual synchronous metastases to the colorectum and bladder

... and specific marker of adenocarcinomas of intestinal ... - Werling - 被引用次数: 415

... of microsatellite instability in colorectal cancer - Boland - 被引用次数: 3319

Occurrence of other malignancies in patients with ... - Agaimy - 被引用次数: 145

Synchronous gist, colon and breast adenocarcinoma with ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: S Jafferbhoy - 2014 - 相关文章

2014年6月4日 - Invasive lobular cancer is the most common histological type of breast

... to a colonic polyp and synchronous colonic adenocarcinoma and a gastric GIST. ...

colorectal clinic with per rectal bleed and underwent a colonoscopy which ... and bladder

cancer.⁶ The most frequent sites of GI tract metastasis are ...

Case report and literature review: Metastatic lobular ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) - 翻译此页

作者: R Arrangoiz - 2011 - 被引用次数: 24 - 相关文章

2011年10月12日 - Invasive lobular carcinoma is the second most common type of

invasive breast ... The incidence of extrahepatic gastrointestinal (GI) tract metastases

observed in ... to the GI tract, along with melanoma, ovarian and bladder cancer. ...

breast cancer presenting as a synchronous rectal and gastric tumors.

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 583,000 条结果 (用时 0.79 秒)

您是不是要找: **Intestinal type gastric adenocarcinoma with unusual synchronous metastases to the *colorectal* and bladder**

Google 学术: **Intestinal type gastric adenocarcinoma with unusual synchronous metastases to the colorectum and bladder**

... and specific marker of adenocarcinomas of intestinal ... - Werling - 被引用次数: 414

... of microsatellite instability in colorectal cancer - Boland - 被引用次数: 3324

Occurrence of other malignancies in patients with ... - Agaimy - 被引用次数: 145

Case report and literature review: Metastatic lobular ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: R Arrangoiz - 2011 - 被引用次数: 24 - 相关文章

2011年10月12日 - Invasive lobular carcinoma is the second most common type of invasive breast ... The incidence of extrahepatic gastrointestinal (GI) tract metastases observed in ... to the GI tract, along with melanoma, ovarian and bladder cancer. ... breast cancer presenting as a synchronous rectal and gastric tumors.

Metachronous Colon Metastases from Gastric ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: JJ Pace - 2009 - 被引用次数: 4 - 相关文章