

Match Overview

1	Internet 58 words crawled on 15-Sep-2013 www.gastrores.org	4%
2	CrossCheck 10 words "Toxicological Profile for Mirex and Chlordane", ATSDR's Toxicological Profiles Web Version, 2002.	1%

Name of journal: *World Journal of Clinical Cases*

ESPS Manuscript NO: 14163

Columns: Case Report

Conservatively managed case of type 2 gallbladder perforation in a child

Dikshit Vishesh, Gupta Rahul, Kothari Paras, Gupta Abhaya, Kamble Ravikiran, Kesan Krushnakumar

Abstract

Gallbladder perforation (GBP) is a rare but serious complication of cholecystitis and needs to be managed promptly. Acalculus cholecystitis leading to GBP is frequently associated with enteric fever and critically ill patients and a surgical approach is not always feasible in such patients. Use of percutaneous tube cholecystostomy (PTC) in such patients is a known entity but it is usually followed by interval cholecystectomy.



Conservatively managed case of type 2 gallbladder perforation in a child



网页

图片

视频

新闻

更多 -

搜索工具

找到约 211,000 条结果 (用时 0.50 秒)

Google 学术: Conservatively managed case of type 2 gallbladder perforation in a child

Sonographic findings in perforation of the gallbladder - Madrazo - 被引用次数: 73

Extrahepatic biliary tract injury: operative management ... - POSNER - 被引用次数: 68

... management guideline based on perforation type - Kwon - 被引用次数: 9

Proposal of an endoscopic retrograde ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: W Kwon - 2012 - 被引用次数: 9 - 相关文章

2012年9月25日 - A retrospective review of 53 patients with ERCP-related perforation ...

Type II injuries were conservatively manageable and demonstrated ... There were 3 cases each of gallbladder cancer, advanced gastric cancer, and benign biliary stricture. The outcomes of conservatively managed type I injuries were ...

Outcomes of Rurally Treated Small-Bowel Perforations ...

<https://ispub.com/IJS/28/2/13921> ▾ 翻译此页

作者: T Allsopp - 2012 - 相关文章

The overall mortality rate was 8.33% (n=2), one was treated conservatively and one surgically. ... This was accomplished by performing a retrospective analysis of all cases of ... type of management (operative and conservative), length of operation, site of ... and

[网页](#)[图片](#)[视频](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 176,000 条结果 (用时 0.72 秒)

Google 学术: Conservatively managed case of type 2 gallbladder perforation in a child

Sonographic findings in perforation of the gallbladder - Madrazo - 被引用次数: 73

Extrahepatic biliary tract injury: operative management ... - POSNER - 被引用次数: 68

... management guideline based on perforation type - Kwon - 被引用次数: 10

Management of Gallstones - American Family Physician

www.aafp.org > ... > [afp](#) > Vol. 72/No. 4(August 15, 2005) ▾ [翻译此页](#)

2005年8月15日 - Many patients with gallstones can be managed expectantly. ... Biliary pain is alleviated by cholecystectomy in the majority of cases. ... greater than 30 kg per m²) as strong risk factors for symptomatic gallstone disease with a more conservative approach is warranted in patients with Child class C cirrhosis ...

[PDF] Gall Bladder complications of Typhoid - JIOM

jiom.com.np/index.php?journal=jiomjournal&page... ▾ [翻译此页](#)

perforation of gall bladder is extremely rare with dreadful ... cholecystitis due to enteric fever, patient was managed conservatively. ... Case 2. A 40 year old male presented with history of fever for 8 days and sudden severe pain ... Widal test was strongly positive for type O and ... during the course of typhoid fever in children.