



20390-Review

BY JIA-REN LIU

Quotes Excluded
Bibliography Excluded1%
SIMILAR

Name of journal: *World Journal of Clinical Cases*

ESPS Manuscript NO: 20390

Columns: Original Article

Basic Study

Leukocyte function-associated antigen-1 deficiency impairs responses to polymicrobial sepsis

Jia-Ren Liu, Xiaohui Han, Sulpicio G Soriano, Koichi Yuki

Abstract

AIM: To determine the role of leukocyte function-associated antigen-1 (LFA-1) in polymicrobial sepsis model in mice.

METHODS: Coecal ligation and puncture model was used to study polymicrobial

Match Overview

1

CrossCheck 54 words

Liu, Jia-Ren, Xiaohui Han, Sulpicio G. Soriano, and Koichi Yuki. "The Role of Macrophage-1 Antigen in Polymicrobial Sepsis"

1%

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 15,900 条结果 (用时 0.73 秒)

Google 学术: Leukocyte function-associated antigen-1 deficiency impairs responses to polymicrobial sepsis

Septic shock - Annane - 被引用次数: 1089

The role of neutrophils in severe sepsis - Alves-Filho - 被引用次数: 110

Neutrophil recruitment and function in health and ... - Kolaczowska - 被引用次数: 355

Increased Susceptibility of ST2-Deficient Mice to ...

www.jimmunol.org/content/187/8/4293.full.pdf ▾ 翻译此页

作者: JM Buckley - 2011 - 被引用次数: 12 - 相关文章

Polymicrobial Sepsis Is Associated with an Impaired Bactericidal Function ... osome maturation and NOX2 function. Thus ... Fit-1, or T1) was originally identified as a serum-induced late response ... the IL-1 family, has been recently identified as the natural and Peritoneal leukocyte accumulation and phagocytic receptor.

[PDF] Neutrophil Migration Under Normal and Sepsis Conditions

www.researchgate.net/...Sepsis.../54cab10a0cf2517b755fbf8b.... ▾ 翻译此页

作者: YV Lerman

inflammatory response with associated tissue damage and subsequent ... neutrophils' inability to contain and control infection and impaired wound healing. to the bone marrow, overall clearance of CXCR4-deficient ... LFA-1 and Mac-1 act through distinct

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 11,800 条结果 (用时 0.88 秒)

Google 学术 : Leukocyte function-associated antigen-1 deficiency impairs responses to polymicrobial sepsis

Septic shock - Annane - 被引用次数 : 1090

Leukocyte apoptosis and its significance in sepsis and ... - Wesche - 被引用次数 : 282

..., and the regulation of Toll-like receptor responses - Murphy - 被引用次数 : 113

Increased Susceptibility of ST2-Deficient Mice to ...

www.jimmunol.org/content/187/8/4293.full.pdf ▾ 翻译此页

作者 : JM Buckley - 2011 - 被引用次数 : 12 - 相关文章

Polymicrobial Sepsis Is Associated with an Impaired Bactericidal Function ... osome maturation and NOX2 function. Thus ... Fit-1, or T1) was originally identified as a serum-induced late response ... the IL-1 family, has been recently identified as the natural and Peritoneal leukocyte accumulation and phagocytic receptor.

[PDF] Neutrophil Migration Under Normal and Sepsis Conditions

www.researchgate.net/...Sepsis.../54cab10a0cf2517b755fbf8b.... ▾ 翻译此页

作者 : YV Lerman

inflammatory response with associated tissue damage and subsequent ... neutrophils' inability to contain and control infection and impaired wound healing. to the bone