



28161-Review

Quotes Excluded
Bibliography Excluded4%
SIMILAR

1

Name of Journal: *World Journal of Clinical Cases*

ESPS Manuscript NO: 28161

Manuscript Type: Case Report

Neurostimulation for fecal incontinence after correction of repair of imperforate anus

Alexandre Bougie, Nathalie McFadden, Sandeep Mayer, Michel Lebel, Ghislain Devroede

Abstract

We are reporting the case of a 32-year-old female who had suffered from fecal incontinence. She was born with an imperforate anus and a recto-vaginal fistula; she underwent repair at 6 months of age. At 29 years of age, she was still fecally incontinent despite extensive pelvic floor reeducation. A magnetic resonance imaging and an anal electromyography were performed. Because her symptoms were considered to be probably due to extra-sphincteric implantation of the neo-anus, a redo was performed of the recto-neo-anal intra-sphincteric anastomosis. A neurostimulator device was subsequently implanted for persistent incontinence. Solid and liquid fecal incontinence resolved, and her quality of life improved markedly. Combining surgery to correct the position of the neo-anus

Match Overview

- 1** **Crossref** 81 words
Leblanc, Dominique, Nathalie McFadden, Michel Lebel, and Ghislain Devroede. "Fecal continence can be restored by sa" **4%**
- 2** **Crossref** 12 words
Tang, Yu-rong, Wei-wei Yang, Yi-Lin Wang, and Lin Lin. "Sex differences in the symptoms and psychological factors that in" **1%**

[全部](#)[图片](#)[视频](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找不到和您查询的"[Sacral Neurostimulation helps establish fecal continence after erroneous repair of ...](#)"相符的内容或信息。

建议：

- 请检查输入字词有无错误。
- 请尝试其他查询词。
- 请改用较常见的字词。
- 请减少查询字词的数量。

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 3,850 条结果 (用时 0.64 秒)

Google 学术: Neurostimulation for fecal incontinence after correction of repair of imperforate anus

... for fecal incontinence in patients with imperforate anus - da Silva - 被引用次数: 42

... of short term sacral nerve stimulation on anal and ... - Vaizey - 被引用次数: 249

Outcomes from the correction of anorectal ... - Levitt - 被引用次数: 126

[PDF] Surgery for Faecal Incontinence - International Continence Society

www.ics.org/publications/ici_3/v2.pdf/chap25.pdf ▾ 翻译此页

作者: RD MADOFF - 被引用次数: 28 - 相关文章

SACRAL NERVE STIMULATION (LEVEL C). 8. ... Therapy for fecal incontinence has traditionally been divided into three ... teen months (median) after a repair, and 60% were incontinent. ... gutter deformity was corrected or adequate symmetrical anal ... reported 7 patients (5 with imperforate anus) aged 6.5 to 19 years ...

Anorectal malformations - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: AN Gangopadhyay - 2015 - 被引用次数: 6 - 相关文章

But still the fecal and urinary incontinence can occur even with an excellent anatomic repair ... KEY WORDS: Anorectal malformations, cloaca, continence, imperforate anus ... Definitive repair of persistent cloaca is performed through PSARP. ... The functional results after surgical correction of ARMs are often assessed by a ...

Anorectal Disease: Fecal Incontinence: Etiology, Evaluation, and ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: DM Hayden - 2011 - 被引用次数: 50 - 相关文章

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 3,910 条结果 (用时 0.62 秒)

您是不是要找: **Neurostimulator** for fecal incontinence after correction of repair of imperforate anus

Google 学术: **Neurostimulation for fecal incontinence after correction of repair of imperforate anus**

... for fecal incontinence in patients with imperforate anus - da Silva - 被引用次数: 42

... of short term sacral nerve stimulation on anal and ... - Vaizey - 被引用次数: 249

Outcomes from the correction of anorectal ... - Levitt - 被引用次数: 126

^[PDF] **Surgery for Faecal Incontinence - International Continence Society**

www.ics.org/publications/ici_3/v2.pdf/chap25.pdf ▾ 翻译此页

作者: RD MADOFF - 被引用次数: 28 - 相关文章

SACRAL NERVE STIMULATION (LEVEL C). 8. ... Therapy for fecal incontinence has traditionally been divided into three teen months (median) after a repair, and 60% were incontinent. gutter deformity was corrected or adequate symme- trical anal reported 7 patients (5 with imperforate anus) aged. 6.5 to 19 years ...

Anorectal malformations - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: AN Gangopadhyay - 2015 - 被引用次数: 6 - 相关文章

But still the fecal and urinary incontinence can occur even with an excellent anatomic repair ... KEY WORDS: Anorectal malformations, cloaca, continence, imperforate anus Definitive repair of persistent cloaca is performed through PSARP. The functional results after surgical correction of ARMs are often assessed by a ...