

**Name of journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 40034

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Cohort Study*

**Diagnostic value of elevated serum carbohydrate antigen 19-9 level in acute cholangitis secondary to choledocholithiasis**

Mei Y *et al.* Diagnostic value of CA19-9 level in secondary acute cholangitis

Yong Mei, Li Chen, Ci-Jun Peng, Jun Wang, Peng-Fei Zeng, Guo-Xing Wang,

## Match Overview

1	<b>Internet</b> 30 words crawled on 15-Nov-2017 <a href="http://hub.hku.hk">hub.hku.hk</a>	1%
2	<b>Internet</b> 25 words crawled on 28-Nov-2017 <a href="http://s-space.snu.ac.kr">s-space.snu.ac.kr</a>	1%
3	<b>Crossref</b> 24 words "Monday 29 August 2016", <i>European Heart Journal</i> , 2016	1%
4	<b>Internet</b> 14 words crawled on 29-Nov-2016 <a href="http://www.basic.medscimonit.com">www.basic.medscimonit.com</a>	1%
5	<b>Crossref</b> 12 words "Abstracts from the Third Chinese Congress on Gerontology and Health Industry", <i>Journal of the American Geriatrics</i>	<1%
6	<b>Internet</b> 11 words crawled on 14-Jun-2018 <a href="http://www.birpublications.org">www.birpublications.org</a>	<1%
7	<b>Crossref</b> 11 words Xiao-Dan Ju, Peng-Fei Li, Peng-Bin Zhang. "Value of serum CA19-9 levels in diagnosis of secondary acute cholangitis"	<1%
8	<b>Internet</b> 11 words crawled on 23-Jun-2017 <a href="http://easpub.com">easpub.com</a>	<1%

[全部](#)[图片](#)[新闻](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 4,050 条结果 (用时 0.56 秒)

### [Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> ▼ [翻译此页](#)

作者: MS Lin - 2014 - 被引用次数: 22 - [相关文章](#)

2014年3月15日 - Although carbohydrate antigen (CA19-9) level is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, .... The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high** serum levels of CA19-9 [22].

缺少字词: 199

### [Highly elevated serum levels of CA 19-9 in choledocholithiasis: a case ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2740064/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2740064/> ▼ [翻译此页](#)

作者: G Marcouizos - 2009 - 被引用次数: 21 - [相关文章](#)

2009年7月30日 - However, high levels of serum CA 19-9 is occasionally found in benign disease of ... stone with **acute cholangitis** presenting with extraordinarily **high** serum CA 19-9 levels, ... A CA 19-9 **value** of more than 1.000 U/ml usually indicates a .... The clinical utility of serum CA 19-9 in the **diagnosis**, prognosis and ...

### [Benign Hydronephrosis and Elevated of Serum Levels of ... - NCBI - NIH](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4917065/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4917065/> ▼ [翻译此页](#)

作者: B Filipovic - 2016 - 被引用次数: 4 - [相关文章](#)

2016年6月11日 - There is no doubt that CA 19-9 level is occasionally elevated in ... in the differential **diagnosis** of elevated concentrations of CA 19-9 marker. ... track such as pancreatitis, **cholangitis**, and **choledocholithiasis** [5-7]. .... Serum **carbohydrate antigen** 19-9 levels in patients with unilateral hydronephrosis **due** to ...

### [An extremely increased CA19-9 level due to common bile duct stone ...](http://amj.amegroups.com/article/view/3666/4397)

[amj.amegroups.com/article/view/3666/4397](http://amj.amegroups.com/article/view/3666/4397) ▼ [翻译此页](#)

作者: X Wang - 2017 - [相关文章](#)

2017年2月9日 - Abstract: **Serum carbohydrate antigen** 19-9 (CA19-9), a common tumor ... that CA19-9 level was more than 1,000.0 U/mL (normal **range**: 0-37 U/mL), ... Serum carbohydrate antigen 19-9 (CA19-9) is mainly used for the **diagnosis** and ... **acute cholangitis**, acute pancreatitis, diabetes, and liver cirrhosis (3).

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 3,420 条结果 (用时 0.50 秒)

### Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> ▼ [翻译此页](#)

作者 : MS Lin - 2014 - 被引用次数 : 22 - [相关文章](#)

2014年3月15日 - Although **carbohydrate antigen (CA19-9) level** is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, ..... The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high serum levels** of CA19-9 [22].

缺少字词 : 499

### Highly elevated serum levels of CA 19-9 in choledocholithiasis: a case ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2740064/> ▼ [翻译此页](#)

作者 : G Marcouizos - 2009 - 被引用次数 : 21 - [相关文章](#)

2009年7月30日 - However, **high levels** of **serum CA 19-9** is occasionally found in benign disease of ... stone with **acute cholangitis** presenting with extraordinarily **high serum CA 19-9 levels**, ... A CA 19-9 **value** of more than 1,000 U/ml usually indicates a .... The clinical utility of **serum CA 19-9** in the **diagnosis**, prognosis and ...

### An extremely increased CA19-9 level due to common bile duct stone ...

[amj.amegroups.com/article/view/3666/4397](http://amj.amegroups.com/article/view/3666/4397) ▼ [翻译此页](#)

作者 : X Wang - 2017 - [相关文章](#)

2017年2月9日 - Abstract: **Serum carbohydrate antigen 19-9 (CA19-9)**, a common tumor ... that CA19-9 **level** was more than 1,000.0 U/mL (normal **range**: 0–37 U/mL), ... **Serum carbohydrate antigen 19-9 (CA19-9)** is mainly used for the **diagnosis** and ... **acute cholangitis**, acute pancreatitis, diabetes, and liver cirrhosis (3).

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 2,970 条结果 (用时 0.42 秒)

小提示：仅限搜索简体中文结果。您可以在设置中指定搜索语言

### [Elevated serum level of carbohydrate antigen 19-9 in benign biliary ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3992417/> [▼ 翻译此页](#)

作者：MS Lin - 2014 - 被引用次数：23 - [相关文章](#)

2014年3月15日 - Although **carbohydrate antigen (CA19-9) level** is frequently upregulated in ... The latter, including inflammatory stricture **secondary to choledocholithiasis**, Mirizzi syndrome, ... The **diagnostic value** of CA19-9 might be improved by using a more ... **Cholangitis** may result in **high serum levels** of CA19-9 [22].

### [\[PDF\] Extraordinarily elevated serum levels of CA 19-9 and rapid decrease ...](http://www.turkjgastroenterol.org/sayilar/220/buyuk/8641.pdf)

[www.turkjgastroenterol.org/sayilar/220/buyuk/8641.pdf](http://www.turkjgastroenterol.org/sayilar/220/buyuk/8641.pdf) [▼ 翻译此页](#)

2010年1月15日 - drate antigen 19-9 **value** of >1,000 U/ml usually indicates a di- gestive cancer and has been ... sis and **choledocholithiasis** with **acute cholangitis** who had very **high serum levels** of **carbohydrate antigen 19-9** (9586 IU/ml). The rapid decrease in ... for the **diagnosis** of pancreatic cancer. A CA 19-9 **value** of ...

### [Oncotarget | Significant increased CA199 levels in acute pancreatitis ...](http://www.oncotarget.com/fulltext/23993)

[www.oncotarget.com/fulltext/23993](http://www.oncotarget.com/fulltext/23993) [▼ 翻译此页](#)

2018年2月27日 - Background and study aims: **Carbohydrate antigen 19-9 (CA199)** has ... Patients with **high CA199 levels** were older and had predominantly biliary ... The mean **CA199 value** in 8 patients with pancreatic cancer was 338.2 ... **levels** [25], and this rate was even higher when **acute cholangitis** was included.