

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,250 条结果 (用时 0.29 秒)

### Retrograde intrarenal surgery versus percutaneous nephrolithotomy in ...

<https://www.ncbi.nlm.nih.gov/pubmed/21657824> ▾ 翻译此页

作者 : OF Bozkurt - 2011 - 被引用次数 : 134 - 相关文章

2011年6月9日 - Retrograde intrarenal surgery versus percutaneous nephrolithotomy in the management of lower-pole renal stones with a diameter of 15 to 20 mm. ... (necessitated transfusion) in three patients who were treated with PCNL.

### Miniaturized percutaneous nephrolithotomy versus retrograde ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5907325/> - 翻译此页

作者 : S Mhaske - 2018 - 相关文章

Keywords: Mini-perc, retrograde intrarenal surgery, small renal calculi ... [2] Owing to this role of miniaturized percutaneous nephrolithotomy (PCNL) (mini-perc) ..... Yttrium-aluminium-garnet laser lithotripsy in the treatment of lower pole renal calculi. ... surgery in the management of renal stones more than 2 cm in diameter.

### Miniaturised percutaneous nephrolithotomy: Its role in the treatment of ...

<https://www.sciencedirect.com/science/article/pii/S2214388218300328> - 翻译此页

作者 : G Zeng - 2018

2018年5月14日 - Common approaches in active treatment of renal stone disease include ... intra-renal surgery (RIRS), and percutaneous nephrolithotomy (PCNL). ... treatment modality in the management of renal stones that are larger than 2 cm [2]. .... useful in the treatment of stones <20 mm in diameter in lower pole calyx.

### [PDF] Comparison of percutaneous nephrolithotomy and retrograde flexibl...

[www.imop.gr/sites/default/files/bjui-imop.pdf](http://www.imop.gr/sites/default/files/bjui-imop.pdf) ▾ 翻译此页

作者 : T Akman - 2011 - 被引用次数 : 154 - 相关文章

retrograde intrarenal surgery as the second-line therapy for the treatment of kidney stones ... treatment of 2 – 4 cm kidney stones. ... ranging from 2 to 4 cm in diameter were .... or 273 μ m laser fibre were used for treatment. We used a holmium laser ..... nephrostolithotomy for lower pole ... benefit from

**Name of Journal:** World Journal of Clinical Cases

**Manuscript NO:** 40696

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Retrograde Intrarenal Surgery versus Miniaturized Percutaneous  
Nephrolithotomy to Treat Lower Pole Renal Stones 1.5-2.5 cm in Diameter**

Mao-Mao Li, Hou-Meng Yang, Xiao-Ming Liu, Hong-Gang Qi, Guo-Bing  
Weng

**Abstract**

*AIM*

Match Overview

1	<b>Crossref</b> 55 words Kirac, Mustafa, Ömer Faruk Bozkurt, Lütfi Tu ne, Cagri Güneri, Ali Ünal, and Hasan Biri	2%
2	<b>Internet</b> 39 words crawled on 17-Jul-2018 <a href="http://www.science.gov">www.science.gov</a>	2%
3	<b>Crossref</b> 17 words Tolga Akman. "Comparison of percutaneous nephrolithotomy and retrograde flexible nep	1%
4	<b>Crossref</b> 16 words Amr S. Fayad, Mohamed G. Elsheikh, Walee d Chameima. "Tubaloss mini percutaneous	1%



[全部](#)[图片](#)[视频](#)[新闻](#)[更多](#)[设置](#)[工具](#)

找到约 3,370 条结果 (用时 0.73 秒)

### Retrograde intrarenal surgery versus percutaneous nephrolithotomy in ...

<https://www.ncbi.nlm.nih.gov/pubmed/21657824> ▼ [翻译此页](#)

作者 : OF Bozkurt - 2011 - 被引用次数 : 138 - [相关文章](#)

2011年6月9日 - Retrograde intrarenal surgery versus percutaneous nephrolithotomy in the management of lower-pole renal stones with a diameter of 15 to 20 mm. ... (necessitated transfusion) in three patients who were treated with PCNL.

### Miniaturized percutaneous nephrolithotomy versus retrograde ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5907325/> - [翻译此页](#)

作者 : S Mhaske - 2018 - [相关文章](#)

The treatment options for small renal calculi (<1.5 cm) include extracorporeal shock wave lithotripsy (ESWL), PCNL, and retrograde intrarenal surgery (RIRS).[3,4,5] However, the limitation of ESWL includes relatively lower stone clearance rates and the need for repeated sessions, especially in lower polar and harder ...

[Results](#) · [INTRODUCTION](#) · [MATERIALS AND METHODS](#)

[PDF]

### Comparison of percutaneous nephrolithotomy and retrograde flexible ...

[www.imop.gr/sites/default/files/bjui-imop.pdf](http://www.imop.gr/sites/default/files/bjui-imop.pdf) ▼ [翻译此页](#)

作者 : T Akman - 2011 - 被引用次数 : 159 - [相关文章](#)

retrograde intrarenal surgery as the second-line therapy for the treatment of kidney stones ... treatment of 2 – 4 cm kidney stones. ... ranging from 2 to 4 cm in diameter were .... or 273 μ m laser fibre were used for treatment. We used a holmium laser ..... nephrostolithotomy for lower pole ... benefit from miniaturized tubeless.

### Role of Minimally Invasive Percutaneous Nephrolithotomy Techniques ...