

找到约 76,200 条结果 (用时 0.70 秒)

Comparative efficacy of combination of 1 L polyethylene glycol, castor ...

https://journals.lww.com/.../Comparative_efficacy_of_combination_of_1_L.... - 翻译此页

作者: X Tian - 2018 - 相关文章

Colonoscopy has been regarded as an important **method** of early ... It remains a challenge to optimize the **bowel preparation efficacy** prior to ... Moreover, the potential of 2 L PEG plus ascorbic acid is equal to that of 3 L PEG, and which is better ... More importantly, 2 trials found that combination of 1 L PEG, bisacodyl, and ...

Efficacy and Acceptability of 1 Liter of Polyethylene Glycol with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5795919/> - 翻译此页

作者: M Kamei - 2018 - 被引用次数: 1 - 相关文章

2018年1月26日 - 2 Liters of Polyethylene Glycol Plus Mosapride and Sennoside for Colonoscopy Preparation ... A low-volume PEG administered with ascorbic acid solution (PEG-Asc) was ... There are many **methods** and preparations for **bowel cleansing**. ... We decided not to prescribe more than 1 L of PEG-Asc solution.

Comparable Efficacy of a 1-L PEG and Ascorbic Acid Solution ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5010253/> - 翻译此页

作者: JE Kwon - 2016 - 被引用次数: 2 - 相关文章

2016年9月2日 - Bowel preparation with Bis and a 1-L PEG/Asc solution is as ... with other lower-volume **methods** leads to **poor** compliance and low tolerability. ... [12,13] The **efficacy** of **bowel cleansing** with 15 mg of Bis plus 2 L of PEG was ...

The use of low-volume polyethylene glycol containing ascorbic acid ...

https://www.researchgate.net/.../274781596_The_use_of_low-volume_polye... - 翻译此页

acid versus 2 L of polyethylene glycol plus bisacodyl as bowel preparation for ... the efficacy and

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 42374

Manuscript Type: ORIGINAL ARTICLE

Clinical Trials Study

Efficacy of 0.5-L vs 1-L of polyethylene glycol containing ascorbic acid as additional colon cleansing methods for inadequate bowel preparation as expected by last stool examination before colonoscopy

Abstract

AIM

To compare 0.5-L and 1-L polyethylene-glycol containing ascorbic acid (PEG+Asc) as additional bowel cleansing methods after a 2-L split-dose PEG+Asc regimen in patients with expected inadequate bowel preparation before colonoscopy.

METHODS

Match Overview

| | | |
|---|---|----|
| 1 | Internet 102 words crawled on 06-Jan-2017 www.e-sciencecentral.org | 3% |
| 2 | Crossref 62 words Jun Lee. "Bowel Preparation Formulations and Salvage Options for Inadequate Preparation", The Korean Journal of Gastroenterology | 2% |
| 3 | Internet 60 words crawled on 12-May-2018 www.kddw.org | 2% |
| 4 | Crossref 44 words Norman Sohn. "Management of the Poorly Prepared Colonoscopy Patient: Colonoscopic Colon Enemas as a Pre-emptive Strategy" | 1% |
| 5 | Internet 38 words crawled on 17-Jun-2017 f1000.com | 1% |
| 6 | Crossref 37 words Mercedes Ibáñez. "Usefulness of an Intensive Bowel Cleansing Strategy for Repeat Colonoscopy After Preparation" | 1% |
| 7 | Crossref 37 words Kjetil Garborg, Håvard Wiig, Audun Hasund, Jon Matre et al. "Gradual stiffness versus magnetic imaging-guided v" | 1% |
| 8 | Internet 32 words crawled on 04-Jul-2016 www.wjgnet.com | 1% |

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

已启用安全搜索

找到约 31,100 条结果 (用时 0.68 秒)

Efficacy and Acceptability of 1 Liter of Polyethylene Glycol with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5795919/> - 翻译此页

作者: M Kamei - 2018 - 被引用次数: 1 - 相关文章

2018年1月26日 - A low-volume PEG administered with ascorbic acid solution (PEG-Asc) was ... To perform a successful colonoscopy, optimal bowel preparation is required. There are many methods and preparations for bowel cleansing. ... 1 L/h. Patients were advised to drink 0.5 L of additional clear fluid after completion.

A Randomized Controlled Trial Evaluating a Low-Volume PEG ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4662975/> - 翻译此页

作者: M Tajika - 2015 - 被引用次数: 10 - 相关文章

跳到Patients and Methods - Comparisons between the 1.5 L PEG+ASC group and the 2 L ... 2.2. Bowel Preparation Methods (Figure 1) ... glycol electrolyte lavage solution containing ascorbic acid. The day before colonoscopy, all patients were admitted to our ... In addition, overall cleansing of the colon was scored by ...

Comparative efficacy of combination of 1 L polyethylene glycol, castor ...

https://journals.lww.com/.../Comparative_efficacy_of_combination_of_1_L.... - 翻译此页

作者: X Tian - 2018 - 相关文章

The aims of this study are to test the hypotheses that low dose castor oil (30 mL) may ... More importantly, 2 trials found that combination of 1 L PEG, bisacodyl, and ... 0.5 L of PEG, 30 mL castor oil and 5 g ascorbic acid and extra 0.5 L clean ... If the inadequate bowel preparation is not suitable for colonoscopy, patients will ...

Optimal intake of clear liquids during preparation for ... - Thieme Connect

<https://www.thieme-connect.com/products/ejournals/.../s-0043-106185.pdf> - 翻译此页

作者: M Tajika - 2017 - 被引用次数: 1 - 相关文章

Adequate bowel preparation is essential for detection of colonic polyps and ... aqueous solution containing