

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 942 条结果 (用时 0.67 秒)

A case of leukemoid reaction in a patient with ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3304658/> - 翻译此页

作者: HP Shin - 2011 - 被引用次数: 7 - 相关文章

2011年9月30日 - Leukemoid reaction can also be a paraneoplastic manifestation of several ...

Leukemoid reaction in patients with sarcomatoid hepatocellular carcinoma was fatal in our case. ...

Microscopy features of hepatocellular carcinoma revealing a typical ... Due to the limited data and case reports, we are not able to ...

(PDF) A case of leukemoid reaction in a patient with sarcomatous ...

https://www.researchgate.net/.../51816533_A_case_of_leukemoid_reaction_i... - 翻译此页

PDF | A leukemoid reaction is defined as reactive leukocytosis exceeding ... neutrophil precursors, and can be a paraneoplastic manifestation of various malignant tumors. ... of the mass confirmed hepatocellular carcinoma (HCC) with a sarcomatoid component. Due to the limited data and case reports, we are not able to.

(PDF) Paraneoplastic Leukemoid Reaction in a Patient with Urothelial ...

https://www.researchgate.net/.../272654570_Paraneoplastic_Leukemoid_Rea... - 翻译此页

Paraneoplastic Leukemoid Reaction in a Patient with Urothelial Carcinoma: A Case Report. Article (PDF Available) in Case Reports in Clinical Medicine 03(05):249-252 high-grade urothelial carcinoma with sarcomatoid differentiation. ... 109/l and a repeat CT scan revealed rapidly progressive disease affecting the liver.

Paraneoplastic Leukemoid Reaction in a Patient with Urothelial ...

file.scirp.org/Html/1-2770365_45758.htm - 翻译此页

Cystoscopy revealed a high-grade urothelial carcinoma with sarcomatoid ... a repeat CT scan revealed rapidly progressive disease affecting the liver and bowel. ... Other case reports of paraneoplastic leukemoid reactions in different cancer ...

A case of leukemoid reaction in a patient with ... - Semantic Scholar

<https://www.semanticscholar.org/...case...leukemoid-reaction-in-a-patient.../2...> - 翻译此页

Liver dynamic computed tomography revealed a hypervascular mass, and an ... of the mass confirmed hepatocellular carcinoma (HCC) with a sarcomatoid component. ... This is a rare case of leukemoid reaction in a patient with sarcomatous HCC. ... and can be a paraneoplastic manifestation of various malignant tumors.

1 **Name of Journal:** *World Journal of Clinical Cases*

2 **Manuscript NO:** 45502

3 **Manuscript Type:** CASE REPORT

4

5 **Paraneoplastic leukemoid reaction in a patient with sarcomatoid hepatocellular**

6 **carcinoma: A case report**

7

8 Hu B *et al.* Liver cancer patient with a complication

9

10 Bo Hu, Xin-Ting Sang, Xiao-Bo Yang

11

12 **Abstract**

13 **BACKGROUND**

14 Sarcomatoid hepatocellular carcinoma (SHC) combined with paraneoplastic leukemoid

15 reaction (PLR), which is associated with poor prognosis, is rarely seen in the clinic. Here,

16 we report the case of a patient in the above situation.

17

Match Overview

1	Crossref 29 words Hyun Phil Shin. "A case of leukemoid reaction in a patient with sarcomatous hepatocellular carcinoma", The Korean	1%
2	Internet 22 words crawled on 07-Sep-2017 hbsn.amegroups.com	1%
3	Crossref 16 words Pranoti Mandrekar, Vincenzo Cardinale. "Periostin and mesothelin: Potential predictors of malignant progression in l ...	1%
4	Internet 16 words crawled on 11-May-2011 casesjournal.com	1%
5	Internet 15 words crawled on 29-Aug-2015 www.wjso.com	1%
6	Internet 14 words crawled on 09-Mar-2019 www.nejm.org	1%
7	Internet 12 words crawled on 10-Jan-2017 e-sciencecentral.org	1%
8	Internet 12 words crawled on 10-Jan-2019 www.cjcrn.org	1%

[国内版](#)[国际版](#)

Paraneoplastic leukemoid reaction in a patient with sarcomatoid hepatocellular ca

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[关闭取词](#)

2,910 Results

Any time ▼

A case of leukemoid reaction in a patient with sarcomatous ...

[www.ncbi.nlm.nih.gov](#) › ... › [Korean J Hepatol](#) › v.17(3); 2011 Sep

INTRODUCTION. **Leukemoid reaction** can also be a **paraneoplastic** manifestation of several **cancers** such as lung, gastrointestinal, genitourinary, ovarian, head and neck **cancers**, 1 and **hepatocellular carcinoma (HCC)**. 2 **Sarcomatoid hepatocellular carcinoma** likely forms a larger tumor than ordinary HCC but prognosis of this disease is as yet unclear.

Cited by: 4

Author: Hyun Phil Shin, Jung Won Jeon, Jae Jun ...

Publish Year: 2011

Paraneoplastic leukemoid reaction in pancreatic cancer: A ...

[www.ncbi.nlm.nih.gov](#) › ... › v.7(10); 2015 Oct 15

Oct 15, 2015 · **CASE REPORT**. We report the case of a 68-year-old man with pancreatic **carcinoma**, who was diagnosed with **paraneoplastic leukemoid reaction** in the absence of plausible differential diagnoses. Our **patient** was diagnosed with pT2N0M0 **carcinoma** of the head of the pancreas, discovered by jaundice, and operated by cephalic duodenopancreatectomy.

Cited by: 3

Author: Mélanie Dos Santos, Karine Bouhier, Ma...

Publish Year: 2015

[PDF] Paraneoplastic Leukemoid Reaction in a Patient with ...

[file.scirp.org/pdf/CRCM_2014051213253520.pdf](#)

A **paraneoplastic leukemoid reaction** has a **cancer -induced cause** in the absence of **leukemia** and infection. The mechanism of this **paraneoplastic** syndrome is thought to result from tumour secretion of granulocyte -colony stimulating factor (G-CSF) [3] -[9] .

Published in: **Case Reports in Clinical Medicine** · 2014

Authors: **Aleksi Suo · Tahir Abbas**

About: **Leukemoid reaction**

[PDF] Leukemoid reaction in sarcomatoid renal cell carcinoma: ...

[citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.812.4337&rep=rep...](#)

CASE REPORT Open Access **Leukemoid reaction** in **sarcomatoid renal cell carcinoma**: a two-case report Weiping Huang†, Feng Wang†, Yeping Li, Feifei Duan and Zhixian Yu* Abstract **Leukemoid reaction** is defined as reactive leukocytosis exceeding 40×10⁹/l, with a significant increase in early neutrophil precursors, and can be a **paraneoplastic** manifestation of various malignant tumors.

[国内版](#)[国际版](#)

Paraneoplastic leukemoid reaction in a patient with sarcomatoid hepatocellular ca

[All](#)[Images](#)[Videos](#)[翻译成中文](#)[开启取词](#)

2,880 Results

Any time ▾

[A case of leukemoid reaction in a patient with sarcomatous ...](#)

[www.ncbi.nlm.nih.gov](#) > ... > [Korean J Hepatol](#) > v.17(3); 2011 Sep

INTRODUCTION. Leukemoid reaction can also be a paraneoplastic manifestation of several cancers such as lung, gastrointestinal, genitourinary, ovarian, head and neck cancers, 1 and hepatocellular carcinoma (HCC). 2 Sarcomatoid hepatocellular carcinoma likely forms a larger tumor than ordinary HCC but prognosis of this disease is as yet unclear.

Cited by: 4

Author: Hyun Phil Shin, Jung Won Jeon, Jae Jun ...

Publish Year: 2011

[Paraneoplastic leukemoid reaction in pancreatic cancer: A ...](#)

[www.ncbi.nlm.nih.gov](#) > ... > v.7(10); 2015 Oct 15

Oct 15, 2015 · CASE REPORT. We report the case of a 68-year-old man with pancreatic carcinoma, who was diagnosed with paraneoplastic leukemoid reaction in the absence of plausible differential diagnoses. Our patient was diagnosed with pT2N0M0 carcinoma of the head of the pancreas, discovered by jaundice, and operated by cephalic duodenopancreatectomy.

Cited by: 3

Author: Mélanie Dos Santos, Karine Bouhier, Ma...

Publish Year: 2015

[\[PDF\] Paraneoplastic Leukemoid Reaction in a Patient with ...](#)

[file.scirp.org/pdf/CRCM_2014051213253520.pdf](#)

A paraneoplastic leukemoid reaction has a cancer -induced cause in the absence of leukemia and infection. The mechanism of this paraneoplastic syndrome is thought to result from tumour secretion of granulocyte -colony stimulating factor (G-CSF) [3] -[9] .

Published in: [Case Reports in Clinical Medicine](#) · 2014

Authors: [Aleksi Suo](#) · [Tahir Abbas](#)

About: [Leukemoid reaction](#)

[\[PDF\] A case of leukemoid reaction in a patient with ...](#)

[www.e-cmh.org/upload/pdf/kjhep-17-226.pdf](#)

Hyun Phil Shin, et al. A case of leukemoid reaction in a patient with sarcomatous hepatocellular carcinoma 227 Figure 1. Computed tomography (CT) findings. (A) 7 months prior to the patient's admission. (B) CT scan obtained on admission showing a hypervascular mass of increased size with central necrosis in the right hepatic lobe. Figure 3.

[Paraneoplastic leukemoid reaction – poor prognostic marker ...](#)

[https://www.mdedge.com/.../paraneoplastic-leukemoid-reaction-poor](#)

Mar 14, 2016 · He H, Zhang Z, Ge J, Zhou W. Leukemoid reaction associated with transitional cell carcinoma: a case report and literature review. Niger J Clin Pract. 2014;17(3):391-394. Certain cancers have been observed to cause symptoms called paraneoplastic syndromes that are not directly attributed to tumor invasion or compression.