

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 17,400 条结果 (用时 0.72 秒)

Basal cell adenoma in the deep portion of the parotid gland: a case ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4699939/>  [翻译此页](#)作者：WY Chung - 2015 - [相关文章](#)

跳到II. **Case Report** - **Based** on the history and physical exam, a suspected diagnosis of ... After the superficial portion of the **parotid gland** was carefully dissected ... the mass was positive for p63 and **high-molecular weight cytokeratine**.

缺少字词： **extending ipsilateral skull mastoid right**

Article: Imaging of salivary gland tumours - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1866314/>  [翻译此页](#)作者：HC Thoeny - 2007 - 被引用次数：112 - [相关文章](#)

Imaging of **salivary gland** tumours is a major challenge for radiologists due to the ... The **large** range of differential diagnoses influences not only prognosis but also determination of perineural **spread** along the **cranial** nerve VII (stylomastoid myoepitheliomas, monomorphic adenomas and **basal cell adenomas**, are ...

Parotid Tumors Extended to the Skull Base | Request PDF

https://www.researchgate.net/.../314626009_Parotid_Tumors_Extended_to_t... - [翻译此页](#)

The **spread** of the **parotid** tumor to the **skull base** and intracranial may happen ... from book Management Of **Extended Parotid** Tumors (pp.65-74) The differential diagnosis included polymorphous low-grade adenocarcinoma, **basal cell adenoma** and adenoid ... A **case report** of adenoid cystic carcinoma of lacrimal **gland**.

Perineural Tumor Spread Along the Auriculotemporal Nerve ...

www.ajnr.org/content/23/2/303  [翻译此页](#)作者：IM Schmalfluss - 2002 - 被引用次数：64 - [相关文章](#)

2002年2月1日 - Adenoid cystic carcinoma of the minor and major **salivary glands** is well ... However, in the **large** percentage of patients with perineural tumor **spread** in the head and neck region, it arises from primary squamous **cell** carcinomas of ... division of the trigeminal nerve was recorded); 4) **skull base** erosion; ...

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 45079

Manuscript Type: CASE REPORT

A large basal cell adenoma extending to the ipsilateral skull base and mastoid in the right parotid gland: A case report

Du LY *et al.* Basal cell adenoma invading skull base

Lu-Yang Du, Xiu-Hong Weng, Zhen-Yu Shen, Bo Cheng

Match Overview

Match Number	Source	Words	Similarity
1	Crossref	42 words	2%
	Woo-Yeol Chung, Chul-Hwan Kim. "Basal cell adenoma in the deep portion of the parotid gland: a case report", Jour ...		
2	Internet	25 words	1%
	crawled on 22-Apr-2010 www.medicinaoral.com		
3	Crossref	18 words	1%
	Chung, Woo-Yeol, and Chul-Hwan Kim. "Basal cell adenoma in the deep portion of the parotid gland: a case report ...		
4	Crossref	14 words	1%
	Lambade, Pravin N., Dilpreet Rajkhokar, and Dipti Lambade. "Basal Cell Adenoma of Submandibular Salivary Gland ...		
5	Internet	13 words	1%
	crawled on 10-Oct-2018 clinmedjournals.org		
6	Crossref	12 words	1%
	Hwang, Seung-Bae, Gyung-Ho Chung, and Chong-Soo Kim. "Basal Cell Adenoma Presenting as a Parapharyngeal ...		
7	Internet	12 words	1%
	crawled on 07-Feb-2017 www.ajnr.org		

A large basal cell adenoma extending to the ipsilateral skull base and mastoid in tl



All

Images

Videos

翻译成中文

关闭取词

22,100 Results

Any time ▾

Basal cell adenoma: A diagnostic dilemma on fine needle ...

https://www.researchgate.net/publication/227811272_Basal_cell...

Taking into account that **basal cell** adenomas represent 1% of all salivary **gland** tumors, being the majority of cases in the **parotid** glands, the goal of this paper is to **report a case of basal cell** ...

Neuro CAQ Flashcards | Quizlet

<https://quizlet.com/164204750/neuro-caq-flash-cards> ▾

Neuro CAQ. "** In addition to the presence of necrosis, the size criterion for pathological cervical **lymph nodes** is a long axis measurement of more than 15 mm for jugulodigastric **lymph nodes** and more than 10 mm for other nodes, except retropharyngeal **lymph nodes**, which are considered **pathologic** is larger ...

Parotid and Temporal Bone Resection for Skull Base ...

www.ncbi.nlm.nih.gov › ... › **Skull Base** › v.20(3); 2010 May

Advanced **lateral skull base malignancies** requiring this approach typically arise from auricular and periauricular skin, external **auditory canal** (EAC) or **parotid gland**. When the **temporal bone** is involved by direct **extension**, **temporal bone resection** is performed to clear the disease.

Cited by: 2

Author: Zoukaa B Sargi, Robert Gerring, Simon I ...

Publish Year: 2010

(PDF) Basal Cell Carcinoma Metastatic to Parotid Gland

https://www.researchgate.net/publication/258529655_Basal_Cell...

Basal Cell Carcinoma Metastatic to Parotid Gland. ... Metastatic Basal Cell Carcinoma: A Case Report and Review of the Literature ... initial involvement of the **skull base** and/or dura by a **basal** ...

Imaging of salivary gland tumours - SlideShare

<https://www.slideshare.net/sg8779/imaging-of-salivary-gland-tumours> ▾

Imaging of salivary gland tumours. • These lymphomas are considered as extranodal marginal zone B-cell lymphomas. • Incidence: 1 and 8% of the patients with lymphomas • All forms of Hodgkin's and non-Hodgkin's lymphomas have been reported. • In the case of secondary lymphomatous involvement of the salivary glands,...



国内版

国际版

A large basal cell adenoma extending to the ipsilateral skull base and mastoid in t



Chat with Bing

Sign in



All

Images

Videos

翻译成中文

关闭取词

Make Bing your search engine

21,800 Results

Any time

Basal cell adenoma: A diagnostic dilemma on fine needle ...

https://www.researchgate.net/publication/227811272_Basal_cell...

Taking into account that **basal cell** adenomas represent 1% of all salivary **gland** tumors, being the majority of cases in the **parotid** glands, the goal of this paper is to **report a case of basal cell** ...

Parotid and Temporal Bone Resection for Skull Base ...

www.ncbi.nlm.nih.gov > ... > **Skull Base** > v.20(3); 2010 May

Advanced **lateral skull base malignancies** requiring this approach typically arise from auricular and periauricular skin, external **auditory canal** (EAC) or **parotid gland**. When the **temporal bone** is involved by direct **extension**, **temporal bone resection** is performed to clear the disease.

Cited by: 2

Author: Zoukaa B Sargi, Robert Gerring, Simon I ...

Publish Year: 2010

Neuro CAQ Flashcards | Quizlet

<https://quizlet.com/164204750/neuro-caq-flash-cards>

Neuro CAQ. ** In addition to the presence of necrosis, the size criterion for pathological cervical **lymph nodes** is a long axis measurement of more than 15 mm for jugulodigastric **lymph nodes** and more than 10 mm for other nodes, except retropharyngeal **lymph nodes**, which are considered **pathologic** is larger ...

Imaging of salivary gland tumours - SlideShare

<https://www.slideshare.net/sg8779/imaging-of-salivary-gland-tumours>

Imaging of salivary gland tumours. • These lymphomas are considered as extranodal marginal zone B-cell lymphomas. • Incidence: 1 and 8% of the patients with lymphomas • All forms of Hodgkin's and non-Hodgkin's lymphomas have been reported. • In the case of secondary lymphomatous involvement of the salivary glands,...

(PDF) Basal Cell Carcinoma Metastatic to Parotid Gland

https://www.researchgate.net/publication/258529655_Basal_Cell...