



11,000 Results

Any time ▾

Effect of Parecoxib as an Adjunct to Patient-Controlled ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5021366>

The present study investigated the **effect of parecoxib** as an adjuvant to a **multimodal PCEA** approach in **patients undergoing gynecological surgery**, with a hypothesis that **parecoxib** might **reduce epidural morphine consumption** (presented as a reduction in PCEA boluses).

Author: Wei-Feng Liu, Hai-Hua Shu, Guo-Don... Publish Year: 2016

Effect of parecoxib combined with thoracic epidural ...

jtd.amegroups.com/article/view/7081/6721 ▾

first carried out a study to find out the **effect of parecoxib** combined with **thoracic paravertebral ropivacaine infusion** in **thoracotomy**. Patients receiving **parecoxib** (40 mg intravenously used 20 min before **extubation** and 12 hrs after the procedure) had less **pain** at rest and during movement compared with those with **placebo** at 12, 24, and 48 hours after **surgery**.

Published in: **Journal of Thoracic Disease** · 2016

Authors: Xiaomin Ling · Fang Fang · Xiaoguang Zhang · Ming Ding · Qiuaxue Liu · Jing Cang

About: **Text mining** · **Bioinformatics**

(PDF) Combined parecoxib and I.V. paracetamol provides ...

https://www.researchgate.net/publication/51203078_Combined_parecoxib_and_IV...

Conclusion: **Intra-articular bupivacaine** in combination with **intravenous parecoxib** may improve **pain relief** and reduce the demand for **rescue analgesics** in patients undergoing **total knee arthroplasty**.

A 96-hour continuous wound infiltration with ropivacaine ...

<https://onlinelibrary.wiley.com/doi/10.1002/jso.25280>

Patients were excluded in case of preplanned laparoscopic or combined procedure, **contraindication** to wound catheter, **abnormal liver function** (**primary liver cancer**, **hepatocellular carcinoma**, and **hepatic cirrhosis**), or suffering from cardiovascular problems (>2 New York Heart Association [NYHA]), or when chronically treated by **morphine** or with a pre-existing abdominal pain, or a known hypersensitivity to **morphine**, **acetaminophen**...

Analgesic effect of preoperative dezocine-based local ...

<https://journals.sagepub.com/doi/full/10.1177/0300060518781708>

Objective This study was performed to investigate the **analgesic effects** of **intramuscular injection** of

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 49413

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Analgesic effect of parecoxib combined with ropivacaine in patients undergoing a laparoscopic hepatectomy

Shan-Shan Huang, Wei-Wei Lv, Yan-Feng Liu, Shao-Zhong Yang

Abstract

BACKGROUND

Currently, there is no uniform standard for analgesia during laparoscopic

Match Overview

5 matches

1	Internet 83 words crawled on 10-Aug-2019 www.wjgnet.com	4%
2	Internet 30 words crawled on 17-Feb-2018 www.spandidos-publications.com	1%
3	Crossref 25 words Cheng Wang, Guoqun Zheng, Wenlong Zhang, Fabiao Zhang, Shangdong Lv, Aidong Wang, Zheping Fang. "Enhanc ...	1%
4	Internet 11 words crawled on 13-Apr-2016 www.ijcem.com	<1%



11,500 Results

Any time ▾

Effect of Parecoxib as an Adjunct to Patient-Controlled ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5021366>

The present study investigated the effect of parecoxib as an adjuvant to a multimodal PCEA approach in patients undergoing gynecological surgery, with a hypothesis that parecoxib might reduce epidural morphine consumption (presented as a reduction in PCEA boluses).

Author: Wei-Feng Liu, Hai-Hua Shu, Guo-Don... Publish Year: 2016

Effect of parecoxib combined with thoracic epidural ...

jtd.amegroups.com/article/view/7081/6721 ▾

first carried out a study to find out the effect of parecoxib combined with thoracic paravertebral ropivacaine infusion in thoracotomy. Patients receiving parecoxib (40 mg intravenously used 20 min before extubation and 12 hrs after the procedure) had less pain at rest and during movement compared with those with placebo at 12, 24, and 48 hours after surgery.

Published in: Journal of Thoracic Disease · 2016

Authors: Xiaomin Ling · Fang Fang · Xiaoguang Zhang · Ming Ding · Qiuaxue Liu · Jing Cang

About: Text mining · Bioinformatics

Comparison of bupivacaine and parecoxib for postoperative ...

<https://docksci.com/comparison-of-bupivacaine-and-parecoxib-for-postoperative-pain...> ▾

Effect of bupivacaine soaked gauze in postoperative pain relief in laparoscopic cholecystectomy: a prospective observational controlled trial in 120 patients. Comparison between lornoxicam quick-release and parecoxib for post-operative analgesia after laparoscopic cholecystectomy: A prospective randomized, placebo-controlled trial.

Enhanced Recovery Program Versus Traditional Care in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4779010>

Mar 03, 2016 · DISCUSSION. Meanwhile, laparoscopic hepatectomy is a minimally invasive surgery that causes less stress and trauma. It can improve patients' recovery and shorten their hospital stay and cost. Therefore, laparoscopic liver resection is an important part of the ERAS program in patients undergoing liver resection.

Cited by: 24

Author: Xiao Liang, Hanning Ying, Hongwei Wang...

Publish Year: 2016