



All

Images

Videos

翻译成中文

关闭取词

30,900 Results

Any time ▾

## The Complications of Polyacrylamide Hydrogel ...

<https://journals.sagepub.com/doi/full/10.1177/2513826X17693821>

The largest case series of PAAAG **breast augmentation**-related complications was described by Luo et al in their experience with 235 patients. 6 The population ranged in age from 20 to 38 years, and the time from injection to **complication** presentation ranged from 6 months to 10 years (39 months). They found the most common **complication** to be ...

**Author:** Jessica Winter, Sarah Shiga, Avinash ... **Publish Year:** 2017

## Does the Use of Intraoperative Breast Sizers Increase ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4820630>

Mar 22, 2016 · Background. Is the use of intraoperative **breast** sizers beneficial for plastic surgeons or do they result in higher **complication** rates? Methods. This is a retrospective study of 416 consecutive cases of primary **breast augmentation** with silicone **implants** at the Plastic Surgery Service of Professor Ivo Pitanguy at the 38th Infirmary Santa Casa Misericórdia Hospital, Rio De Janeiro, from January ...

**Cited by:** 2

**Author:** Lee Seng Khoo, Henrique N. Radwanski, ...

**Publish Year:** 2016

## Complication of AQUAfilling ® Gel Injection for Breast ...

<https://link.springer.com/article/10.1007/s00266-018-1107-0>

This is a **case report** describing a patient that suffered complications after AQUAfilling ® gel injection for **breast augmentation**. Results A 32-year-old female patient was referred to our plastic surgery department with a chief complaint of **repeated** wound dehiscence and ...

**Cited by:** 1

**Author:** Bok Ki Jung, In Sik Yun, Young Seok Ki...

**Publish Year:** 2018

**Location:** Seoul

## FULL TEXT - A case report of breast implant-associated ...

[www.ijcasereportsandimages.com/archive/2016/008-2016-ijcri/CR-10683-08-2016-binmahfouz/...](http://www.ijcasereportsandimages.com/archive/2016/008-2016-ijcri/CR-10683-08-2016-binmahfouz/)

**Case Report.** The differential **diagnosis** was **malignancy** versus aggressive **infection**, therefore a contrast-enhanced staging CT scan of the chest, abdomen and pelvis was performed, which re-demonstrated an irregular left **breast** soft tissue mass posterosuperior to the **implant** extending into the adjacent intercostal spaces,...

**Cited by:** 1

**Author:** Alya Binmahfouz, Karin Steinke

**Publish Year:** 2016

# Repeated lumps and infection: A case report of breast augmentation complication

Ming-Xuan Zhang, Shi-Yan Li, Li-Long Xu, Bo-Wen Zhao, Xiao-Yan Cai, Guang-Lan Wang

## Abstract

### BACKGROUND

Polyacrylamide hydrogel (PAAG) injection was once prevalent in breast augmentation, and <sup>1</sup>had been prohibited for augmentation mammoplasty in China, since a great quantity of patients whose breasts were augmented with the PAAG injection have continued to seek medical advice as a result of its complications. Among all these complications, distant migration is relatively rare.

### CASE SUMMARY

A 49-year-old female presented to the hospital with one-year history of vulva lump. The lump sonography showed several subcutaneous fluid regions from left vulva to pubic symphysis, and considered it as the possibility of fat liquefaction. An enhanced magnetic resonance imaging revealed a cystic area, which was considered as benign lesion. Intraoperative observation showed

## Match Overview

1	<b>Crossref</b> 64 words Sheng-Kang Luo, Gunang-Ping Chen, Zhong-Shen Sun, Ning-Xin Cheng. "Our strategy in complication management	3%
2	<b>Crossref</b> 19 words Dmytro Unukovych, Vasyl Khrapach, Marie Wickman, Annelie Liljegren, Volodymyr Mishalov, Gennadiy Patlazhan, Kersti	1%
3	<b>Crossref</b> 13 words Rami Mossad Ibrahim, Elisabeth Lauritzen, Caspar Weel Kramer. "Breastfeeding difficulty after Polyacrylamid Hydr	1%



All

Images

Videos

关闭取词

28,800 Results

Any time ▾

## Case Report: Percutaneous breast implant herniation: a ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4289805>

This case details the first report of percutaneous breast implant herniation resulting from periprosthetic TB infection. While the majority of non-mycobacterial and NTM-associated breast implant infections arise from intraoperative contamination, in this case it is likely that the infection resulted from haematogenous mycobacterial seeding.

Author: Adam P Dale, Martin J Dedicoat, Taus... Publish Year: 2015

## FULL TEXT - A case report of breast implant-associated ...

[www.ijcasereportsandimages.com/archive/2016/008...](http://www.ijcasereportsandimages.com/archive/2016/008...) ▾

Case Report: We report an extremely uncommon presentation of BIA-ALCL in a 55-year-old lady who presented 23 years post-bilateral cosmetic breast implants with a locally aggressive mass in the left breast. The mass invaded the chest wall and was associated with ...

Cited by: 1 Author: Alya Binmahfouz, Karin Steinke

Publish Year: 2016

## Recurrent complications of PAAG implants during lactation ...

<https://casereports.bmj.com/content/2018/bcr-2017-219688> ▾

Nov 05, 2018 · Complications range from breast induration and lumps, recurrent infections, gel leakage and implant migration<sup>8</sup> to haematoma, galactocoele, potential misdiagnosis in malignancy screening<sup>2</sup> and even possible breast cancer.<sup>9</sup> The mean time from injection to complications was 6.1 years in a review of 106 patients in Ukraine who had operations for ...

Author: Anita Geraldine Bourke, Chaitra Jose Publish Year: 2018

Online: ISSN 1757-790X

## Late seroma during pregnancy, a rare complication in ...

<https://www.sciencedirect.com/science/article/abs/pii/S1748681511006395>

Case report Late seroma during pregnancy, a rare complication in prosthetic breast augmentation: Case report Author links open overlay panel Forrest S. Roth a d Daniel J. Gould b c d Chuma J. Chike-Obi a Jamal M. Bullocks a

Cited by: 5 Author: Forrest S. Roth, Daniel J. Gould, Daniel J.