

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 51045

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

Human podocyte injury in the early course of hypertensive renal injury

Sun D *et al.* Human podocyte in hypertensive renal injury

Da Sun, Jiao-Jiao Wang, Wei Wang, Juan Wang, Li-Ning Wang, Li Yao, Ying-Hui

Sun, Zi-Long Li

Abstract

BACKGROUND

Match Overview

1	Internet 87 words crawled on 12-Aug-2018 academic.oup.com	3%
2	Internet 42 words crawled on 25-Jul-2019 www.physiology.org	1%
3	Internet 37 words crawled on 29-Mar-2019 journals.plos.org	1%
4	Crossref 33 words Masanori Hara, Toshio Yanagihara, Itaru Kihara, Kazuhiro Higashi, Kotarou Fujimoto, Tadahiro Kajita. "Apical Cell Me	1%
5	Crossref 29 words Gunter Wolf, Rolf AK Stahl. "CD2-associated protein and glomerular disease", The Lancet, 2003	1%
6	Crossref 28 words Arndt Petermann, Jürgen Floege. "Podocyte Damage Resulting in Podocyturia: A Potential Diagnostic Marker to Asses	1%
7	Crossref 26 words Xuezhu Li, Peter Y. Chuang, Vivette D. D'Agati, Yan Dai et al. "Nephrin Preserves Podocyte Viability and Glomerular Str	1%



All

Images

Videos

关闭取词

972,000 Results

Any time ▾

Podocyte Loss in Human Hypertensive Nephrosclerosis ...

<https://academic.oup.com/ajh/article/22/3/300/131366> ▾

Mar 01, 2009 - Podocyte injury probably plays a crucial role in the renal damage of hypertensive nephrosclerosis. 11–13 In fact, podocyte loss due to severe podocyte injury has been proved to be directly related to glomerulosclerosis in other kidney diseases such as diabetic nephropathy and minimal change disease. 14,15 Recent studies also showed that ...

Cited by: 83

Author: Gang Wang, Fernand Mac-Moune Lai, Bo...

Publish Year: 2009

Author: Wang, Gang

[PDF] Early development of podocyte injury independently of ...

<https://www.physiology.org/doi/pdf/10.1152/ajprenal.00590.2015>

ment of renal injury in the absence of hypertension and diabetes remain unclear. The early stages of renal disease associated with obesity are renal vasodilation, hypertrophy, and increased renal blood flow (RBF) and glomerular filtration rate (GFR) (hyperfiltration). There are several components of

Cited by: 1

Author: Kasi C McPherson, Lateia Taylor, Ashley ...

Publish Year: 2016

Podocyte injury in diabetic nephropathy: implications of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4674698>

Dec 10, 2015 - Injury to podocytes is considered a major contributor to diabetic kidney disease: their loss causes proteinuria and progressive glomerulosclerosis. Podocyte depletion may result from improper calcium handling due to abnormal activation of the calcium permeant TRPC (Transient Receptor Potential Canonical) channels.

Cited by: 35

Author: Daria V. Ilatovskaya, Vladislav Levchenko...

Publish Year: 2015

Podocyte Injury Underlies the Glomerulopathy of Dahl Salt ...

<https://www.ahajournals.org/doi/full/10.1161/01.HYP.0000222003.28517.99>

Podocyte Injury Commences at Early Stage in DSH. Time course analysis of podocyte injury revealed that desmin, an early injury marker, was already expressed in some glomeruli of DSHs at 2 weeks of salt loading (Figure 3A and 3B). We further assessed podocyte damage by transmission electron microscopy (Figure 3C). Podocyte foot processes, which line the outer surface of ...

Cited by: 302

Author: Miki Nagase, Shigeru Shibata, Shigetaka...

Publish Year: 2006



Human podocyte injury in the early course of hypertensive rer



ALL

IMAGES

VIDEOS

关闭取词

971,000 Results

Any time ▾

Podocyte Loss in Human Hypertensive Nephrosclerosis ...

<https://academic.oup.com/ajh/article/22/3/300/131366> ▾

Mar 01, 2009 - Podocyte injury probably plays a crucial role in the renal damage of hypertensive nephrosclerosis. 11–13 In fact, podocyte loss due to severe podocyte injury has been proved to be directly related to glomerulosclerosis in other kidney diseases such as diabetic nephropathy and minimal change disease. 14,15 Recent studies also showed that ...

Cited by: 83

Author: Gang Wang, Fernand Mac-Moune Lai, Bo...

Publish Year: 2009

Author: Wang, Gang

[PDF] Early development of podocyte injury independently of ...

<https://www.physiology.org/doi/pdf/10.1152/ajprenal.00590.2015>

ment of renal injury in the absence of hypertension and diabetes remain unclear. The early stages of renal disease associated with obesity are renal vasodilation, hypertrophy, and increased renal blood flow (RBF) and glomerular filtration rate (GFR) (hyperfiltration). There are several components of

Cited by: 1

Author: Kasi C McPherson, Lateia Taylor, Ashley ...

Publish Year: 2016

Podocyte injury in diabetic nephropathy: implications of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4674698>

Dec 10, 2015 - Injury to podocytes is considered a major contributor to diabetic kidney disease: their loss causes proteinuria and progressive glomerulosclerosis. Podocyte depletion may result from improper calcium handling due to abnormal activation of the calcium permeant TRPC (Transient Receptor Potential Canonical) channels.

Cited by: 35

Author: Daria V. Ilatovskaya, Vladislav Levchenko...

Publish Year: 2015

Podocyte Injury Underlies the Glomerulopathy of Dahl Salt ...

<https://www.ahajournals.org/doi/full/10.1161/01.HYP.0000222003.28517.99>

Podocyte Injury Commences at Early Stage in DSH. Time course analysis of podocyte injury revealed that desmin, an early injury marker, was already expressed in some glomeruli of DSHs at 2 weeks of salt loading (Figure 3A and 3B). We further assessed podocyte damage by transmission electron microscopy (Figure 3C). Podocyte foot processes, which line the outer surface of ...

Cited by: 302

Author: Miki Nagase, Shigeru Shibata, Shigetaka...

Publish Year: 2006