

国内版 国际版

Successful treatment of warfarin-induced skin necrosis using c



All

Images

Videos

关闭取词

24,800 Results

Any time ▾

Warfarin-Induced Skin Necrosis - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4319805>

Dec 11, 2014 · Following the diagnosis of warfarin-induced skin necrosis, the patient was administered **vitamin K intravenously** and **fresh frozen plasma (FFP)** to reverse the effects of warfarin and promote **protein C and S synthesis**. Once the **patient's INR** was no longer supratherapeutic, **subcutaneous enoxaparin** was re-started as treatment for the known recent DVT.

Cited by: 7

Author: Neda Pourdeyhimi, Zackery Bullard

Publish Year: 2014

[PDF] Rivaroxaban dose adjustment using thrombin generation ...

www.bloodadvances.org/content/bloodoa/2/2/142.full.pdf?sso-checked=true

that can cause skin necrosis at treatment outset.³ Warfarin-induced skin necrosis (WISN) is rare but more likely to occur in severe PC deficiency.⁴ Rivaroxaban reversibly inhibits free and bound factor Xa and prevents the thrombin burst during the amplification phase of coagulation. It has predictable pharmacokinetic (PK) and pharmacodynamic (PD)

Heparin-induced thrombocytopenia and warfarin-induced ...

<https://onlinelibrary.wiley.com/doi/full/10.1046/j.1538-7836.2003.00057.x>

Feb 07, 2003 · Heparin-induced thrombocytopenia and warfarin-induced skin necrosis in a child with severe protein C deficiency: successful treatment with dermatan sulfate and protein C concentrate. After 10 days, DS was shifted from the intravenous to intramuscular route (11 mg kg⁻¹ day⁻¹, by single daily injection).

Cited by: 17

Author: L. Gatti, V. Carnelli, R. Rusconi, M. Moia

Publish Year: 2003

Rivaroxaban dose adjustment using thrombin generation in ...

<https://ashpublications.org/.../2/2/142/16287/Rivaroxaban-dose-adjustment-using-thrombin> ▾

Rivaroxaban reversibly inhibits free and bound factor Xa and prevents the thrombin burst during the amplification phase of coagulation. It has predictable pharmacokinetic (PK) and

Cited by: 1

Author: Neethu Menon, Ravindra Sarode, Ayesha...

Publish Year: 2018

(PDF) Warfarin Induced Necrosis of The Breast: A Case Report

Warfarin

Common brand names

This medication may cause bleeding from forming blood clots.

Precautions



Side Effects

Brand names

Drug classes

Pregnancy

Availability

May treat: S

People also ask

Heparin

Rivaroxaban

Clopidogrel

Aspirin

Enoxaparin

See more ▾

Data from First

Consult a medi

Suggest an ed



25,300 Results

Any time ▼

Warfarin-Induced Skin Necrosis - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4319805>

Dec 11, 2014 - Following the diagnosis of warfarin-induced skin necrosis, the patient was administered **vitamin K intravenously** and **fresh frozen plasma (FFP)** to reverse the effects of warfarin and promote **protein C** and **S** synthesis. Once the **patient's INR** was no longer supratherapeutic, **subcutaneous enoxaparin** was re-started as treatment for the known recent DVT.

Cited by: 7

Author: Neda Pourdeyhimi, Zackery Bullard

Publish Year: 2014

Rivaroxaban dose adjustment using thrombin generation in ...

<https://ashpublications.org/.../2/2/142/16287/Rivaroxaban-dose-adjustment-using-thrombin> ▼

Jan 19, 2018 - A **rivaroxaban** dose of 40 mg divided 4 times daily was necessary to **prevent recurrence of skin necrosis** in a child with **PS deficiency** and **WISN**. 7 In contrast, our patient did not experience similar **recurrences of skin necrosis** at lower doses. The PK profile of the PS-deficient patient demonstrated **rivaroxaban levels** <25 µg/L 8 hours after a 10-mg dose, whereas our patient had a ...

Cited by: 1

Author: Neethu Menon, Ravindra Sarode, Ayesha...

Publish Year: 2018

Heparin-induced thrombocytopenia and warfarin-induced ...

<https://onlinelibrary.wiley.com/doi/full/10.1046/j.1538-7836.2003.00057.x>

In summary, this is a case report of an uncommon clinical presentation in a **pediatric** patient with a rare congenital prothrombotic state. **Anticoagulant treatment with DS** and **replacement therapy with PC** concentrate were safe and effective measures for **treating** two serious complications of **anticoagulant therapy**: HIT and **warfarin-induced skin necrosis**.

Cited by: 17

Author: L. Gatti, V. Carnelli, R. Rusconi, M. Moia

Publish Year: 2003

Anticoagulation therapy for thromboembolism prevention: a ...

<https://casereports.bmj.com/content/2017/bcr-2016-218015> ▼

May 12, 2017 - **Non-vitamin K antagonist anticoagulant** options should be considered in patients with **warfarin-induced skin necrosis (WISN)** in the setting of protein C. We report a 41-year-old African American male patient with **WISN** and protein C deficiency who was **treated with rivaroxaban** ...

Cited by: 3

Author: Jonathan Lai, Daryl Ramai, Ramiz Alchi, ...

Publish Year: 2017

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 51210

Manuscript Type: CASE REPORT

Successful treatment of warfarin-induced skin necrosis using oral rivaroxaban: A case report

Momoka Kamada, Tsuneaki Kenzaka

Abstract

BACKGROUND

Heparin is commonly recommended for warfarin-induced skin necrosis; however, there is currently no established therapy for this disease. We present a serious case of warfarin-induced skin necrosis that was successfully treated with oral rivaroxaban, a factor Xa inhibitor.

Match Overview

1	Crossref 13 words Makimoto, Go, Michiko Asano, Nobukazu Fujimoto, Yasuko Fuchimoto, Katsuichiro Ono, Shinji Ozaki, Koji Taguchi, and	1%
2	Crossref 12 words "Society of general internal medicine 23rd annual meetin ... Boston, Massachusetts May 4-6, 2000 abstracts", Journal	1%
3	Internet 11 words crawled on 18-Mar-2016 www.deckerip.com	1%
4	Internet 11 words crawled on 24-Mar-2016 austinpublishinggroup.com	1%
5	Crossref 10 words Sumba Harrison, Hamza Lamchahab, Jacquet Djamal, Yousuness Jabbour et al. "Cystitis Glandularis: A Rare Benign..."	1%



All

Images

Videos

关闭取词

25,300 Results

Any time ▾

Warfarin-Induced Skin Necrosis - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4319805>

Dec 11, 2014 · Following the diagnosis of warfarin-induced skin necrosis, the patient was administered **vitamin K intravenously** and **fresh frozen plasma (FFP)** to reverse the effects of warfarin and promote **protein C** and S synthesis. Once the **patient's INR** was no longer supratherapeutic, **subcutaneous enoxaparin** was re-started as treatment for the known recent DVT.

Cited by: 7

Author: Neda Pourdeyhimi, Zackery Bullard

Publish Year: 2014

Rivaroxaban dose adjustment using thrombin generation in ...

<https://ashpublications.org/.../2/2/142/16287/Rivaroxaban-dose-adjustment-using-thrombin> ▾

Jan 19, 2018 · A **rivaroxaban** dose of 40 mg divided 4 times daily was necessary to **prevent recurrence of skin necrosis** in a child with **PS deficiency** and WISN. 7 In contrast, our patient did not experience similar **recurrences of skin necrosis** at lower doses. The PK profile of the PS-deficient patient demonstrated **rivaroxaban levels** <25 µg/L 8 hours after a 10-mg dose, whereas our patient had a ...

Cited by: 2

Author: Neethu Menon, Ravindra Sarode, Ayesha...

Publish Year: 2018

(PDF) Warfarin Induced Necrosis of The Breast: A Case Report

https://www.researchgate.net/publication/329835214_Warfarin_Induced_Necrosis_of_The...

Warfarin-induced **skin necrosis** is an infrequent and well-recognized complication of warfarin **treatment**. The incidence was estimated between 0.01% and 0.1% whereas a paradoxal prothrombotic state ...

Heparin-induced thrombocytopenia and warfarin-induced ...

<https://onlinelibrary.wiley.com/doi/full/10.1046/j.1538-7836.2003.00057.x>

Heparin-induced thrombocytopenia and warfarin-induced **skin necrosis** in a child with **severe protein C deficiency**: **successful treatment with dermatan sulfate** and protein C concentrate. After 10 days, DS was shifted from the intravenous to intramuscular route (11 ...