



Isolated peritoneal lymphomatosis defined as post-transplant



ALL

IMAGES

VIDEOS

关闭取词

12,100 Results

Any time ▾

Posttransplantation lymphoproliferative disorder involving ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3304639>

Posttransplantation lymphoproliferative disorder (PTLD) is a serious complication following solid organ transplantation. Most of PTLD is related to an interaction of Epstein-Barr virus (EBV) infection and immune suppression. 1, 2 PTLD occurs most common **after intestine transplantation** and the least common **after renal transplantation**. 3, 4 PTLD can involve any organs.

Cited by: 1

Author: Yong Moon Shin

Publish Year: 2011

Isolated Peritoneal Donor-Related Plasmacytoma 3 Years ...

https://www.researchgate.net/publication/259489522_Isolated_Peritoneal_Donor-Related...

Post-transplant lymphoproliferative disorder (PTLD) is a serious complication after organ transplantation with a cumulative incidence of 1.1% at 18 months and 4.7% at 15 years.

Plasmacytoma-like post-kidney-transplant ...

<https://www.sciencedirect.com/science/article/pii/S1769725518305637>

Post-transplantation lymphoproliferative disorder (PTLD) is a well-know complication **after organ transplantation**. We report a case of a patient who developed an **extramedullary plasmacytoma-like PTLD** around his **transplanted kidney** treated with standard multiple myeloma chemotherapy.

Cited by: 2

Author: Vincent Thibaud, Joseph Rivalan, Francis...

Publish Year: 2018

Posttransplantation Lymphoproliferative Disease: Proposed ...

<https://pubs.rsna.org/doi/full/10.1148/rg.347130130>

Nov 10, 2014 - The liver is the most common site of **disease when lymphoma occurs after liver transplantation**, and the lungs are the most common site of **disease when lymphoma occurs after lung or heart-lung transplantation** (35).

Cited by: 14

Author: Juan C. Camacho, Courtney Coursey Mor...

Publish Year: 2014

Primary central nervous system post-transplant ...

<https://insightsimaging.springeropen.com/articles/10.1186/s13244-019-0726-6> ▾

Apr 11, 2019 - Central nervous system post-transplant lymphoproliferative disorder (CNS-PTLD) is a rare **disorder due to immunosuppression secondary to solid organ, stem cell, or bone marrow transplantation** [1, 2]. Cases begin to appear as early as 6 weeks in the first year **after transplantation** but most cases take years to present **after transplantation** and the attendant

Match Overview

1	Internet 80 words crawled on 31-Oct-2019 www.wjgnet.com	4%
2	Internet 27 words crawled on 09-Jan-2014 www.tumoronline.it	1%
3	Internet 19 words crawled on 08-Sep-2017 preview-virologyj.biomedcentral.com	1%
4	Internet 15 words crawled on 21-Mar-2019 mjhid.org	1%
5	Internet 12 words crawled on 23-Jun-2017 www.jove.com	1%
6	Internet 12 words crawled on 02-Apr-2019 f6publishing.blob.core.windows.net	1%
7	Internet 11 words crawled on 26-Nov-2018 jmedicalcasereports.biomedcentral.com	<1%

6
Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 51868

Manuscript Type: CASE REPORT

Isolated peritoneal lymphomatosis defined as post-transplant lymphoproliferative disorder after a liver transplant: A case report

Hong Beum Kim, Ran Hong, Yung Sub Na, Woo Young Choi, Sang Gon Park, Hee Jeong Lee

Abstract

BACKGROUND

Post-transplant lymphoproliferative disorder (PTLD) is a fatal complication of solid organ transplantation or allogenic hematopoietic stem cell transplantation that is



国内版 国际版

Isolated peritoneal lymphomatosis defined as post-transplant



All

Images

Videos

关闭取词

12,100 Results

Any time ▾

Posttransplantation lymphoproliferative disorder involving ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3304639>

Posttransplantation lymphoproliferative disorder (PTLD) is a serious complication following solid organ transplantation. Most of PTLD is related to an interaction of Epstein-Barr virus (EBV) infection and immune suppression. 1, 2 PTLD occurs most common after intestine transplantation and the least common after renal transplantation. 3, 4 PTLD can involve any organs.

Cited by: 1

Author: Yong Moon Shin

Publish Year: 2011

Plasmacytoma-like post-kidney-transplant ...

<https://www.sciencedirect.com/science/article/pii/S1769725518305637>

Post-transplantation lymphoproliferative disorder (PTLD) is a well-know complication after organ transplantation. We report a case of a patient who developed an extramedullary plasmacytoma-like PTLD around his transplanted kidney treated with standard multiple myeloma chemotherapy.

Cited by: 2

Author: Vincent Thibaud, Joseph Rivalan, Francis...

Publish Year: 2018

Isolated Peritoneal Donor-Related Plasmacytoma 3 Years ...

https://www.researchgate.net/publication/259489522_Isolated_Peritoneal_Donor-Related...

Post-transplant lymphoproliferative disorder (PTLD) is a serious complication after organ transplantation with a cumulative incidence of 1.1% at 18 months and 4.7% at 15 years.

Posttransplantation Lymphoproliferative Disease: Proposed ...

<https://pubs.rsna.org/doi/full/10.1148/rg.347130130>

Nov 10, 2014 - The liver is the most common site of disease when lymphoma occurs after liver transplantation, and the lungs are the most common site of disease when lymphoma occurs after lung or heart-lung transplantation (35).

Cited by: 14

Author: Juan C. Camacho, Courtney Coursey Mor...

Publish Year: 2014

Post-liver-transplant anemia: Etiology and management ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/lt.20031>

Jan 30, 2004 - Anemia is common in patients with chronic liver disease. 1-3 A high prevalence of anemia has been reported in recipients of kidney, heart, and lung transplantation. 4-7 Although the