



Comparative evaluation of standard prophylaxis versus splitted-dose NS



全部

图片

新闻

地图

更多

设置

工具

找到约 5,580 条结果 (用时 0.60 秒)

Nonsteroidal anti-inflammatory drugs for prevention of post-ERCP ...

<https://www.ncbi.nlm.nih.gov/pubmed/23164513> ▼ 翻译此页

作者: X Ding - 2012 - 被引用次数: 103 - 相关文章

INTERVENTIONS: NSAIDs use for the prevention of PEP. ... Meta-analysis showed that NSAID use decreased the overall incidence of PEP (risk ratio [RR], ... However, after removing the main source of heterogeneity, the prophylactic efficacy was ... of the NSAIDs, study heterogeneity, inconsistent use of pancreatic stenting.

Effectiveness of nonsteroidal anti-inflammatory drugs in prevention of ...

<https://www.ncbi.nlm.nih.gov/pubmed/25232268> ▼ 翻译此页

作者: X Li - 2014 - 被引用次数: 21 - 相关文章

2014年9月14日 - The pooled RR of PEP incidence with prophylactic NSAID administration ... which demonstrates that NSAID administration after ERCP significantly reduced the incidence of PEP when compared to the placebo group ($P < 0.0001$). ... Moreover, the administration of NSAIDs as a rectal suppository (RR = 0.35, ...

Rectal nonsteroidal anti-inflammatory drugs administration is effective ...

<https://www.ncbi.nlm.nih.gov/pubmed/28734720> - 翻译此页

作者: C Yang - 2017 - 被引用次数: 6 - 相关文章

2017年7月17日 - ... is effective for the prevention of post-ERCP pancreatitis: An updated ... Whether the prophylactic administration of rectal non-steroidal ... the risk reduction for different NSAID types, for general vs. high-risk patients, ... CONCLUSIONS: A single rectal dose of NSAIDs is effective in preventing PEP both in ...

Role of nonsteroidal anti-inflammatory drugs in the prevention of post ...

<https://www.ncbi.nlm.nih.gov/pubmed/19208508> ▼ 翻译此页

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 43221

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Split-dose or hybrid NSAIDs and N-acetylcysteine therapy for prevention of post-ERCP pancreatitis

Pavel L⁹ *et al.* Pharmacological prophylaxis of post-ERCP pancreatitis

Laura Pavel, Gheorghe Gh Bălan, Alexandra Nicorescu, Georgiana Emmanuela Gîlcă-Blănariu, Cătălin Sfarti, Ștefan Chiriac, Smaranda Diaconescu, Vasile Liviu Drug, Gheorghe Bălan, Gabriela Ștefănescu

Abstract

AIM

Match Overview

1	Internet 67 words crawled on 12-May-2018 www.wjgnet.com	2%
2	Internet 57 words crawled on 24-Nov-2017 bmcgastroenterol.biomedcentral.com	1%
3	Internet 42 words crawled on 03-Dec-2018 onlinelibrary.wiley.com	1%
4	Crossref 29 words Lan Liu, Chenghao Li, Yuan Huang, Haiyan Jin. "Nonsteroidal Anti-inflammatory Drugs for Endoscopic Retrograd	1%
5	Internet 25 words crawled on 13-Apr-2015 www.ncbi.nlm.nih.gov	1%
6	Internet 22 words crawled on 13-Nov-2018 annalsofintensivecare.springeropen.com	1%
7	Crossref 17 words "P060: Incidence of hepatocellular carcinoma in chronic hepatitis B patients responders to nucleoside analogue	<1%
8	Internet 12 words crawled on 28-Oct-2017 thorax.bmj.com	<1%

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 9 条结果 (用时 0.37 秒)

Pharmacological Prophylaxis of Post- Endoscopic ... - ClinicalTrials.gov

<https://clinicaltrials.gov/ct2/show/NCT03708458> ▼ [翻译此页](#)

2018年10月17日 - ... Prophylaxis Versus Divided-dose NSAIDs or Hybrid NSAID and N-acetylcysteine Therapy for the Prevention of Post-ERCP Pancreatitis.

Prevention of post-ERCP pancreatitis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4024515/> ▼ [翻译此页](#)

作者 : LL Wong - 2014 - 被引用次数 : 34 - [相关文章](#)

2014年2月15日 - Severity of post-ERCP pancreatitis is graded based on length of Many advanced endoscopists use a hybrid of the two techniques (wire The pooled number needed to treat with NSAIDs to prevent one episode of pancreatitis was 15; N-acetylcysteine and sodium selenite have both been studied in ...

Prevention of Post-ERCP Pancreatitis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3154520/> - [翻译此页](#)

作者 : F Donnellan - 2012 - 被引用次数 : 56 - [相关文章](#)

2011年6月13日 - Post-ERCP pancreatitis is defined as acute pancreatitis that has developed de ... patients undergoing unselected ERCP (both diagnostic and therapeutic). subcutaneously in three divided doses) 24 hours prior to ERCP and placebo N-acetylcysteine is a free radical scavenger and has been shown to ...

(PDF) Prevention of Post-ERCP pancreatitis - ResearchGate

https://www.researchgate.net/.../51574861_Prevention_of_Post-ERCP_pancr... - [翻译此页](#)

evidence to support their use in the prevention of post-ERCP pancreatitis. 1. unselected ERCP (both diagnostic and therapeutic). It was NSAIDs are potent inhibitors of a number of inflammatory

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 7,580 条结果 (用时 0.42 秒)

Pharmacological Prophylaxis of Post- Endoscopic Retrograde ...

<https://clinicaltrials.gov/ct2/show/NCT03708458> - 翻译此页

2018年10月17日 - ... Prophylaxis Versus Divided-dose NSAIDs or Hybrid NSAID and N-acetylcysteine Therapy for the Prevention of Post-ERCP Pancreatitis.

(PDF) Prevention of Post-ERCP pancreatitis - ResearchGate

https://www.researchgate.net/.../51574861_Prevention_of_Post-ERCP_pancr... - 翻译此页

evidence to support their use in the prevention of post-ERCP pancreatitis. 1. ... unselected ERCP (both diagnostic and therapeutic). It was NSAIDs are potent inhibitors of a number of inflammatory (0.6 mg subcutaneously in three divided doses) 24 hours randomized to N-acetylcysteine or placebo, respectively.

Prevention of post-ERCP pancreatitis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4024515/> - 翻译此页

作者: LL Wong - 2014 - 被引用次数: 34 - 相关文章

2014年2月15日 - Severity of post-ERCP pancreatitis is graded based on length of Many advanced endoscopists use a hybrid of the two techniques (wire The pooled number needed to treat with NSAIDs to prevent one episode of pancreatitis was 15; N-acetylcysteine and sodium selenite have both been studied in ...

Prevention of Post-ERCP Pancreatitis - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3154520/> - 翻译此页

作者: F Donnellan - 2012 - 被引用次数: 56 - 相关文章

2011年6月13日 - Post-ERCP pancreatitis is defined as acute pancreatitis that has developed de ... patients undergoing unselected ERCP (both diagnostic and therapeutic). subcutaneously in three divided doses) 24 hours prior to ERCP and placebo N-acetylcysteine is a free radical scavenger and has been shown to ...

Medications and Methods for the Prevention of Post-ERCP Pancreatitis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5439140/> - 翻译此页

作者: AY Wang - 2017 - 被引用次数: 3 - 相关文章

However, in patients who are at elevated risk for post-ERCP pancreatitis, the ... Prevention of post-endoscopic retrograde cholangiopancreatography ... AW Nonsteroidal anti-inflammatory drugs (NSAIDs) work by inhibiting ... G&H What other therapies are available for post-ERCP pancreatitis prevention? N Engl J Med.