



国内版

国际版

Mechanical intestinal obstruction due to isolated diffuse veno



ALL

IMAGES

VIDEOS

关闭取词

23,700 Results

Any time ▾

Solitary intestinal neurofibroma with no associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3679427>

The clinical presentation of **isolated** neurofibromatous lesions of the **intestines** are myriad and are dependent upon the focal or **diffuse** nature of the lesions, their location, their effect on **gastrointestinal motility** and their possible impingement on adjacent structures, **13 resulting** in abdominal pain, palpable masses, **bleeding due** to necrosis or ulceration of the mucosa, **obstruction due** to intussusception or ...

Cited by: 4

Author: Ali Al-Harake, Mohomad Chour, Osama S...

Publish Year: 2013

(PDF) Mechanical bowel obstruction due to colonic ...

https://www.academia.edu/18418381/Mechanical_bowel_obstruction_due_to_colonic... ▾

The **Mechanical** bowel **obstruction due** to colonic hemangioma 307 incidence of vascular **malformations** of the gastro- vomiting and constipation. Physical examination **intestinal tract** is reported as 0.06% in the litera- revealed minimal abdominal distension and bowel ture (7).

Muco-submucosal elongated polyps of the gastrointestinal ...

https://www.researchgate.net/publication/236106042_Muco-submucosal_elongated_polyps_of...

Muco-submucosal elongated polyps of the **gastrointestinal tract**: A case series and a review of the literature Article · Literature Review in World Journal of Gastroenterology 19(11):1845-9 ...

Malformations, choristomas, and hamartomas of the ...

<https://www.sciencedirect.com/science/article/pii/S074025701830114X>

Intestinal atresias and stenoses are often accompanied by other **gastrointestinal** and **extra-intestinal** anomalies. 16 The most common association is that of **duodenal atresia** with trisomy 21. 14 Familial atresias **due** to 2p21 mutations affecting TTC7A are inherited in an autosomal recessive fashion and are often lethal in infancy. 17 The diagnosis of **intestinal atresia** is usually suspected following detection of polyhydramnios and **dilated intestinal** ...

Author: Melanie Johncilla, Rhonda K. Yantiss

Publish Year: 2019

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 51811

Manuscript Type: CASE REPORT

Mechanical intestinal obstruction due to isolated diffuse venous malformations in the gastrointestinal tract: A case report and review of literature

Li HB *et al.* Diffuse venous malformations in GI tract

Han-Bo Li, Jing-Fang Lv, Ning Lu, Zong-Shun Lv

Abstract

Match Overview

1	Crossref 118 words Han Tao Wang, Xian Hua Gao, Chuan Gang Fu, Lie Wang, Rong Gui Meng, Lian Jie Liu. "Diagnosis and Treatm ...	3%
2	Internet 61 words crawled on 01-Jul-2016 www.wjgnet.com	1%
3	Crossref 46 words Patrícia Andrade, Susana Lopes, Guilherme Macedo. "D iffuse cavernous hemangioma of the rectum: case rep ...	1%
4	Internet 25 words crawled on 24-Dec-2017 www.karger.com	1%
5	Crossref 24 words Xian-rui Wu, Wei-wen Liang, Xing-wei Zhang, Liang Kang, Ping Lan. "Transanal total mesorectal excision as a s	1%
6	Crossref 23 words J. El-Sheikha, M. W. Little, M. Bratby. "Rectal Venous Ma... ormation Treated by Superior Rectal Artery Embolizatio	1%
7	Internet 18 words crawled on 19-Oct-2018 www.cureus.com	<1%
8	Crossref 17 words Shigeru Kimura. "Cavernous hemangioma in the asce ... ding colon treated by endoscopic mucosal resection", Jo	<1%
9	Crossref 17 words A M Kaiser, C Spanos. "An unusual cause of rectal bleed ing", The British Journal of Radiology, 2005	<1%
10	Internet 15 words crawled on 05-Mar-2016 pediatrics.aappublications.org	<1%
11	Internet 14 words crawled on 19-Sep-2017 scielo.isciii.es	<1%



All

Images

Videos

关闭取词

23,400 Results

Any time ▼

Solitary intestinal neurofibroma with no associated ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3679427>

The clinical presentation of **isolated** neurofibromatous lesions of the **intestines** are myriad and are dependent upon the focal or **diffuse** nature of the lesions, their location, their effect on **gastrointestinal motility** and their possible impingement on adjacent structures, **13 resulting** in abdominal pain, palpable masses, **bleeding due** to necrosis or ulceration of the mucosa, **obstruction due** to intussusception or ...

Cited by: 4

Author: Ali Al-Harake, Mohomad Chour, Osama S...

Publish Year: 2013

Multiple venous malformations in the left colon and rectum ...

<https://casereports.bmj.com/content/12/3/e227700> ▼

Mar 01, 2019 · **Venous** or cavernous malformations of the **colon** or rectum are a rare cause of lower **gastrointestinal bleeds**. It has been previously described as a **diffuse** cavernous haemangioma which was thought to be a benign **vascular tumour**. It mainly affects the rectosigmoid area of the **gastrointestinal** tract and is most common in children and young adults.

Author: Sashiananthan Ganesananthan, Jonat... Publish Year: 2019

Diffuse infantile hemangiomatosis of the ileum presenting ...

<https://www.sciencedirect.com/science/article/pii/S0022346810004264>

Diffuse involvement of the **gastrointestinal tract** is rare. Anomalies in the **vascular tumor** class grow mostly by endothelial hyperplasia, whereas **malformations** have no such endothelial **dysfunction**. **Malformations** are caused by dysregulated vasculogenesis and endure ...

Cited by: 6

Author: Anita B. Rao, Jeffrey C. Pence, David L. ...

Publish Year: 2010

Muco-submucosal elongated polyps of the gastrointestinal ...

https://www.researchgate.net/publication/236106042_Muco-submucosal_elongated_polyps_of...

Muco-submucosal elongated polyps of the **gastrointestinal tract**: A case series and a review of the literature Article · Literature Review in World Journal of Gastroenterology 19(11):1845-9 ...

Abdominal ultrasonography of the pediatric ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4965350>

Jul 28, 2016 · **Gastrointestinal duplication** cysts demonstrate a connection with the **gastrointestinal**