



All

Images

Videos

关闭取词

5,610 Results

Any time ▾

Uterine Cystic Adenomyosis: A Disease of Younger Women ...

https://www.researchgate.net/publication/262690832_Uterine_Cystic_Adenomyosis_A...

Uterine Cystic Adenomyosis: A Disease of Younger Women. Few data on the diagnostic accuracy of hysteroscopy in the diagnosis of adenomyosis are available; however, office hysteroscopy may add information in young patients with irregular bleeding [62], and it is indicated in patients undergoing in vitro fertilization (IVF).

(PDF) Laparoscopic excision of uterine adenomyomas ...

https://www.academia.edu/18775312/Laparoscopic_excision_of_uterine_adenomyomas ▾

Ultrasound examination Hysterectomy remains the mainstay of treatment for dif- identified at least one uterine lesion with characteristics of fuse or focal adenomyosis, whereas the use of levonorges- an adenomyoma in each case (mean preoperative size, 52 trel intrauterine device is the preferable medical treatment mm; range, 30–84 mm) (Table 1).

Laparoscopic excision of uterine adenomyomas - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S0015028207010059>

In women who want to preserve their uteri, the current approach for adenomyosis and/or adenomyomas is medical treatment or conservative surgery. Medical treatment involves the use of a levonorgestrel-releasing intrauterine system 37, 38, 39, the use of danazol-containing intrauterine device, or a 3- to 6-month course of GnRH-analogs 40, 41.

Cited by: 22

Author: Grigoris F. Grimbizis, Themistoklis Mikos...

Publish Year: 2008

Laparoscopic excision of uterine adenomyomas - Fertility ...

[https://www.fertstert.org/article/S0015-0282\(07\)01005-9/fulltext](https://www.fertstert.org/article/S0015-0282(07)01005-9/fulltext)

Hysterectomy remains the mainstay of treatment for diffuse or focal adenomyosis, whereas the use of levonorgestrel intrauterine device is the preferable medical treatment (1 x 1 Farquhar, C. and Brosens, I. Medical and surgical management of adenomyosis.

Cited by: 22

Author: Grigoris F. Grimbizis, Themistoklis Mikos...

Publish Year: 2008

ClinicalKey Latest Issue: Journal of Minimally Invasive ...

<cdn.clinicalkey.com/rss/issue/15534650.xml>

Journal of Minimally Invasive Gynecology, The, Volume 26, Issue 7, Supplement, 2019 Nov-Dec, Pages

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 51175

Manuscript Type: CASE REPORT

Giant exophytic cystic adenomyosis with a levonorgestrel containing intrauterine device out of the uterine cavity after uterine myomectomy: A case report

Yong Zhou, Zheng-Yun Chen, Xin-Mei Zhang

Abstract

BACKGROUND

A cystic lesion arising from the myometrium of the uterus, termed as cystic adenomyosis, has chocolate-like, thick viscous contents and contains various amounts of endometrial stroma below the glandular epithelium. It is an extremely rare type of adenomyosis

Match Overview

1	Crossref 36 words Brosens, Ivo, Stephan Gordts, Marwan Habiba, and Giuseppe Benagiano. "Uterine Cystic Adenomyosis: A Disease (...)	2%
2	Internet 30 words crawled on 14-Mar-2019 www.wjgnet.com	2%
3	Crossref 16 words Hiroshi Nabeshima, Takashi Murakami, Yukihiro Terada, Takahiro Noda, Nobuo Yaegashi, Kunihiro Okamura. "Total	1%
4	Crossref 12 words Malcolm G. Munro. "Uterine polyps, adenomyosis, leiomyomas, and endometrial receptivity", <i>Fertility and Sterility</i> , 201	1%



All

Images

Videos

关闭取词

5,650 Results

Any time ▾

Uterine Cystic Adenomyosis: A Disease of Younger Women ...

https://www.researchgate.net/publication/262690832_Uterine_Cystic_Adenomyosis_A...

Uterine Cystic Adenomyosis: A Disease of Younger Women. Few data on the diagnostic accuracy of hysteroscopy in the **diagnosis of adenomyosis** are available; however, office hysteroscopy may add information in young patients with irregular bleeding [62] , and it is indicated in patients undergoing in vitro fertilization (IVF).

Tuboovarian abscess mimicking intraligamentar uterine ...

https://www.researchgate.net/publication/7449948_Tuboovarian_abscess_mimicking...

Case 2: A 46-year-old woman with an **intrauterine device** had a preoperative diagnosis of **uterine myoma** with degeneration. At **laparotomy**, an **omentum cake** with **dense pelvic adhesions** was noted.

Laparoscopic excision of uterine adenomyomas - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S0015028207010059>

In women who want to preserve their uteri, the current approach for adenomyosis and/or adenomyomas is medical treatment or conservative surgery. Medical treatment involves the use of a **levonorgestrel-releasing intrauterine system** 37, 38, 39, the use of **danazol-containing intrauterine device** , or a 3- to 6-month course of **GnRH-analogs** 40, 41.

Cited by: 22

Author: Grigoris F. Grimbizis, Themistoklis Mikos...

Publish Year: 2008

UWIS Complete QBank Flashcards | Quizlet

<https://quizlet.com/210591457/uwice-complete-qbank-flash-cards> ▾

The **levonorgestrel intrauterine device** has lower failure rates within the first year of use than does the **copper containing intrauterine device**. It causes more disruption in menstrual bleeding, especially during the first few months of use, although the overall volume of bleeding is decreased long-term and many women become amenorrheic.

Ob Gyn Flashcards | Quizlet

<https://quizlet.com/87316218/ob-gyn-flash-cards> ▾

Intrauterine Growth Restricted. Nausea and vomiting are the most common **side effects** of the **oral hormonal drugs**. Single- and **split-dose levonorgestrel regimens** have similar rates of nausea and