

1 / 19

7

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 55403

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Combination of endoscopic submucosal dissection and laparoscopic sentinel lymph node dissection in early mucinous gastric cancer: Role of lymph node metastasis

Hua Li, Li-Li Zhao, Xiao-Chong Zhang, Deng-Xiang Liu, Gui-Ying Wang, Zhi-Bin Huo

Match Overview

| | | |
|---|--|----|
| 1 | Internet 93 words crawled on 26-Jan-2020 www.scilit.net | 3% |
| 2 | Internet 64 words crawled on 20-Jan-2020 link.springer.com | 2% |
| 3 | Crossref 52 words Hua Li, Lin Chen, Zhibin Huo, Hongqing Xi, Jianxin Cui, Xudong Zhao. "Defining a Subgroup Treatable for Laparoscopic | 2% |
| 4 | Crossref 41 words Weon Jin Ko, Yoo Min Kim, In kyung Yoo, Joo Young Cho. "Clinical outcomes of minimally invasive treatment for earl ... | 1% |
| 5 | Crossref 33 words Huo, Zhi-Bin. "Risk clinicopathological factors for lymph node metastasis in poorly differentiated early gastric cancer a ... | 1% |
| 6 | Crossref 29 words Hua Li, Ping Lu, Yang Lu, Cai-Gang Liu, Hui-Mian Xu, Shu-Bao Wang, Jun-Qing Chen. "Predictive factors for lymph nod | 1% |
| 7 | Internet 23 words crawled on 19-Oct-2019 f6publishing.blob.core.windows.net | |
| 8 | Crossref 22 words Yuko Kitagawa, Seigo Kitano, Tetsuro Kubota, Koichiro Ku mai, Yoshihide Otani, Yoshiro Saikawa, Masashi Yoshida, M | 1% |





The combination of endoscopic submucosal dissection a



ALL

IMAGES

VIDEOS

26,300 Results

Any time ▾

[Long-term outcomes of combination of endoscopic ...](#)

<https://www.sciencedirect.com/science/article/pii/S0016510711017834>

The combination of endoscopic submucosal dissection and laparoscopic lymph node dissection can be an effective, minimally invasive treatment that maintains long-term quality of life for selected early gastric cancer patients who have a potential risk of nodal metastasis.

Cited by: 68

Author: Nobutsugu Abe, Hirohisa Takeuchi, Atsuk...

Publish Year: 2011

[Sentinel Lymph Node Navigation Surgery for Early Gastric ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4834616>

Abe N, Takeuchi H, Ohki A, Yanagida O, Masaki T, Mori T, et al. Long-term outcomes of combination of endoscopic submucosal dissection and laparoscopic lymph node dissection without gastrectomy for early gastric cancer patients who have a potential risk of lymph node metastasis. *Gastrointest Endosc.* 2011; 74:792–797.

Cited by: 2

Author: Guilherme Pinto Bravo Neto, Elizabeth G...

Publish Year: 2016

[Prospective Multicenter Feasibility Study of Laparoscopic ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6589427>

Apr 15, 2019 · Prospective Multicenter Feasibility Study of Laparoscopic Sentinel Basin Dissection after Endoscopic Submucosal Dissection for Early Gastric Cancer: SENORITA 2 Trial Protocol ... residual tumor and lymph node metastasis have not been identified after surgery. The aim of this study is to evaluate the feasibility of sentinel node navigation ...

Cited by: 2

Author: Bang Wool Eom, Hong Man Yoon, Jae S...

Publish Year: 2019

[Laparoscopic lymph node dissection after endoscopic ...](#)

<https://www.sciencedirect.com/science/article/pii/S0002961005005040>

Endoscopic submucosal dissection (ESD), a newly developed endoscopic mucosal resection (EMR) technique, is a valuable alternative to gastrectomy for treating early-stage gastric cancer (EGC)The accurate assessment of the potential presence of lymph node metastasis (LNM) is required for the appropriate treatment (endoscopic treatment or gastrectomy with lymph node dissection).

Cited by: 81

Author: Nobutsugu Abe, Toshiyuki Mori, Hirohisa ...



ALL

IMAGES

VIDEOS

16,800 Results

Any time ▾

Sentinel Lymph Nodes in Endometrial Cancer Update 2018

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6743227>

Aug 29, 2019 · INTRODUCTION. Endometrial **cancer** (EC) is the most common gynecological **cancer** in Europe.[] **Lymph node** evaluation is the key point in EC staging and prognosis.[]The International Federation of Gynecology and Obstetrics (FIGO) staging of ECs revised in 2009 to include the pelvic and para-aortic lymphadenectomy (PPLND) as one of the most important prognostic factors in ECs.[]

Cited by: 2

Author: Ibrahim A Abdelazim, Mohannad Abu-Fa...

Publish Year: 2019

Long-term outcomes of combination of endoscopic ...

<https://www.researchgate.net/publication/51666343...>

Long-term outcomes of **combination of endoscopic submucosal dissection and laparoscopic lymph node dissection** without gastrectomy for **early gastric cancer** patients who have a potential risk of ...

Sentinel lymph node biopsy for gastric cancer: Where do we ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3192223>

Sep 27, 2011 · Keywords: **Sentinel lymph node** biopsy, **Gastric cancer**, Laparoscopy, **Lymph node dissection**, Lymphatic **metastasis**, Staining and labeling INTRODUCTION Over the past two decades, **sentinel lymph node** (SLN) biopsy in surgical oncology has been so successful that it is now considered as the gold standard procedure in breast **cancer** surgery.


Cited by: 15

Author: Mehmet Fatih Can, Gokhan Yagci, Sadett...

Publish Year: 2011

Precision surgical approach with lymph-node dissection in ...

<https://www.researchgate.net/publication/332410034>



IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

17,100 Results

Any time ▾

[Sentinel Lymph Node Navigation Surgery for Early Gastric ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4834616>

Abe N, Takeuchi H, Ohki A, Yanagida O, Masaki T, Mori T, et al. Long-term outcomes of **combination of endoscopic submucosal dissection and laparoscopic lymph node dissection** without gastrectomy for **early gastric cancer** patients who have a potential risk of **lymph node metastasis**. *Gastrointest Endosc.* 2011; 74:792–797.

Cited by: 3

Author: Guilherme Pinto Bravo Neto, Elizabeth G...

Publish Year: 2016

Precision surgical approach with lymph-node dissection in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6465935>

Apr 14, 2019 · The gravest prognostic factor in **early gastric cancer** is **lymph-node metastasis**, with an incidence of about 10% overall. About two-thirds of **early gastric cancer** patients can be diagnosed as **node-negative** prior to treatment based on clinic-pathological data. Thus, the tumor can be resected by **endoscopic submucosal dissection**.

Cited by: 3

Author: Shinichi Kinami, Naohiko Nakamura, Yas...

Publish Year: 2019

Current role of minimally invasive approaches in the ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3983444>

Apr 14, 2014 · Abe N, Takeuchi H, Ohki A, Yanagida O, Masaki T, Mori T, Sugiyama M. Long-term outcomes of **combination of endoscopic submucosal dissection and laparoscopic lymph node dissection** without gastrectomy for **early gastric cancer** patients who have a potential risk of **lymph node metastasis**. *Gastrointest Endosc.* 2011; 74:792–797.

Cited by: 11

Author: Abraham El-Sedfy, Savtaj S Brar, Natalie ...

Publish Year: 2014

Prospective Multicenter Feasibility Study of Laparoscopic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6589427>

~~A 15 0010 B C M H I E J L O U V W X Y Z~~