

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 55800

Manuscript Type: CASE REPORT

**Diagnosis and treatment of mixed infection of hepatic cystic and alveolar
echinococcosis: Four case reports**

Ji-De A, Jin-Ping Chai, Hao Wang, Wei Gao, Zhe Peng, Shun-Yun Zhao,
Xiang-Ren A

Abstract

BACKGROUND

Match Overview

There are no matching sources for this report.



ALL

IMAGES

VIDEOS

34,100 Results

Any time ▾

Diagnosis and treatment of a case of hepatic mixed ...

<https://www.ncbi.nlm.nih.gov/pubmed/31156003>

Jun 02, 2019 · Cystic echinococcosis, which is one of the most common types of echinococcosis, is described as swelling and growth of cystic lesions. Alveolar echinococcosis, which is less common, is invasive. Cases of mixed echinococcosis infection accompanied by ...

Cited by: 1

Author: Xiaolei Xu, Cancan Gao, Haiwen Ye, Zhixi...

Publish Year: 2019

Diagnosis, treatment, and management of echinococcosis ...

<https://www.bmj.com/content/344/bmj.e3866> ▾

Jun 11, 2012 · Summary points. Echinococcosis is a parasitic zoonosis caused by Echinococcus cestode worms. The two major species of medical importance are Echinococcus granulosus and E. multilocularis, which cause cystic echinococcosis (CE) and alveolar echinococcosis (AE), respectively. CE and especially AE are life threatening chronic diseases with a high fatality rate and ...

Cited by: 182

Author: Donald P McManus, Darren J Gray, Wen...

Publish Year: 2012

CDC - Echinococcosis - Resources for Health Professionals

https://www.cdc.gov/parasites/echinococcosis/health_professionals ▾

Diagnosis

Disease

Treatment

The diagnosis of E. granulosus infection (cystic hydatid disease) is suggested by identification of a cyst-like mass in a person with a history of exposure to sheepdogs in areas where the parasite is endemic. Cystic echinococcosis must be differentiated from benign cysts, cavitary tuberculosis, mycoses, abscesses, and benign or malignant neoplasms. Noninvasive imaging techniques such as CT scans, MRI, and ultrasound imaging are all used for detecting and defining the extent and condition of a...

[See more on cdc.gov](#)

Published: Aug 28, 2019

Echinococcosis: Symptoms, Causes, Treatment and Prevention

<https://iytmed.com/echinococcosis> ▾

Overview

Pathophysiology

Prognosis

Signs and symptoms





Diagnosis and treatment of mixed infection of hepatic cy  

Sign in



ALL IMAGES VIDEOS

12,700 Results Any time ▾

Diagnosis and treatment of a case of hepatic mixed ...

<https://journals.sagepub.com/doi/full/10.1177/0300060519851651>

The **liver** section and the **hepatic** and intestinal anastomosis were placed in the drainage tube. The operation took 300 minutes, intraoperative blood loss was approximately 3000 mL, and the blood transfusion volume was 2000 mL. The postoperative pathological **diagnosis** was **hepatic alveolar echinococcosis** and **hepatic cystic echinococcosis** .

Diagnosis and treatment of a case of hepatic mixed ...

<https://www.ncbi.nlm.nih.gov/pubmed/31156003>

Jun 02, 2019 · Cases of **mixed echinococcosis infection** accompanied by extrahepatic organ metastasis are extremely rare. **Treatment** of these cases is complicated and the prognosis is extremely poor. We report a **case of hepatic echinococcosis** in a 40-year-old Tibetan man who was treated with the **hepatic** right tricuspidectomy + left **hepatic** duct jejunostomy (Roux ...

Cited by: 1 **Author:** Xiaolei Xu, Cancan Gao, Haiwen Ye, Zhix...

Publish Year: 2019

[PDF] Diagnosis and treatment The Author(s) 2019 of a case of ...

<https://journals.sagepub.com/doi/pdf/10.1177/0300060519851651>

Search Tools

Turn off Hover Translation (关闭取词)



12,700 Results Any time ▾

[Diagnosis and treatment of a case of hepatic mixed ...](#)

<https://journals.sagepub.com/doi/full/10.1177/0300060519851651>

The **liver** section and the **hepatic** and intestinal anastomosis were placed in the drainage tube. The operation took 300 minutes, intraoperative blood loss was approximately 3000 mL, and the blood transfusion volume was 2000 mL. The postoperative pathological **diagnosis** was **hepatic alveolar echinococcosis** and **hepatic cystic echinococcosis** .

[PDF] [Diagnosis and treatment The Author\(s\) 2019 of a case of ...](#)

<https://journals.sagepub.com/doi/pdf/10.1177/0300060519851651>

multilocularis. **Cystic echinococcosis**, which is one of the most common types of **echinococcosis**, is described as swelling and growth of **cystic** lesions. **Alveolar echinococcosis**, which is less common, is invasive. Cases of **mixed echinococcosis infection** accompanied by ...

[Pitfalls in diagnosis and treatment of alveolar ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4599161>

Alveolar echinococcosis (AE) is a neglected zoonosis presenting with focal **liver** lesions (FLL). 1 In endemic regions, awareness of this disease is high due to the known transmission risk (figure 1). 2 AE should, however, always be included in the differential **diagnosis** of FLLs due to the high mobility of patients. The clinical problem is thus ...

Cited by: 16 **Author:** M. Stojkovic, C. Mickan, Tf F. Weber, T. J...

Publish Year: 2015

[A new treatment strategy for end-stage hepatic alveolar ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6602961>

The **diagnosis** of **alveolar echinococcosis** (AE) mainly depends on the following clues: epidemiological evidence (contact history in endemic area), signs and **symptoms**, serological findings (human echinococcus IgG antibody by ELISA), imaging findings (computed tomography [CT], magnetic resonance imaging [MRI], ultrasound), intraoperative probing ...

Cited by: 1 **Author:** Qiancheng Du, Yanyan Wang, Yanyan W...