

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 56737

Manuscript Type: CASE REPORT

Low-grade fibromyxoid sarcoma of the liver: A case report

Dugalic V¹⁹ *et al.* Low-grade fibromyxoid sarcoma of the liver

Vladimir Dugalic, Igor I Ignjatovic, Jelena Djokic Kovac, Nikola Ilic, Jelena Sopta, Slavenko R Ostojic, Dragan Vasin, Marko D Bogdanovic, Igor Dumic, Tamara Milovanovic

⁸
Abstract

BACKGROUND

Low grade fibromyxoid sarcoma (LGFMS) is a rare and benign mesenchymal tumor with indolent course, most commonly found in young or middle-aged men. The majority of the LGFMSs are located in the trunk and deep soft tissue of the lower

Match Overview

1	Crossref 51 words Eriko Maeda, Satoshi Ohta, Takeyuki Watadani, Akiteru Goto, Atsushi Nakajima, Kuni Ohtomo. "Imaging findings ..."	2%
2	Internet 48 words crawled on 25-Oct-2015 www.researchgate.net	2%
3	Internet 41 words crawled on 18-Aug-2009 www.biomedcentral.com	2%
4	Internet 33 words crawled on 16-Jun-2020 www.karger.com	1%
5	Internet 30 words crawled on 28-Mar-2019 synapse.koreamed.org	1%
6	Crossref 25 words Atlas of Upper Gastrointestinal and Hepato-Pancreato-Biliary Surgery, 2016.	1%
7	Crossref 22 words Bharat Rekhi, Mahesh Deshmukh, Nirmala A. Jambhekar. "Low-grade fibromyxoid sarcoma: a clinicopathologic stud	1%
8	Internet 20 words crawled on 28-Nov-2019 www.hematology.fi	1%



ALL

IMAGES

VIDEOS

38,000 Results

Any time ▾

Including results for [low-grade fibromyxoid sarcoma of the liver a case report review of the literature](#).

Do you want results only for [LOW-GRADE FIBROMYXOID SARCOMA OF THE LIVER: A CASE REPORT AND REVIEW OF THE LITERATURE?](#)

A Huge Low-Grade Fibromyxoid Sarcoma of Small Bowel Mesentery Simulating Hyper Immune Splenomegaly Syndrome: A Case Report and Review of Literature - PubMed LGFMS may **cause a diagnostic dilemma**, especially in a third world setting where preoperative diagnosis is hampered by lack of facilities and poverty.

[A Huge Low-Grade Fibromyxoid Sarcoma of Small Bowel ...](#)
pubmed.ncbi.nlm.nih.gov/24250315/

Was this helpful?

[Low grade fibromyxoid sarcoma: a case report and review of ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2920234>

Jul 29, 2010 · Background. Low grade **fibromyxoid sarcoma** (LGFMS) is a distinctive variant of fibrosarcoma. According to Evans [], who first described this pathologic entity, LGFMS presents a rare soft-tissue tumour with a high metastasizing potential, despite the benign histologic appearance. The sometimes long interval between tumour presentation and metastasis poses ...

Cited by: 18

Author: Christina Arnaoutoglou, Marios G. Lykis...

Publish Year: 2010

[MRI findings of low-grade fibromyxoid sarcoma: a case ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6389061>

Mastoraki A, Strigkos T, Tatakis FP, Christophi A, Smyrniotis V. Recurrent low-grade **Fibromyxoid sarcoma** of the neck: **report of a case and review of the literature**. Indian Journal of Surgical

Low grade fibromyxoid sarcoma of the falciform ligament: a ...

Dec 01, 2002 · We report a 37-year old man with a giant **intra-abdominal low grade fibromyxoid sarcoma** measuring **21 cm** in size; which was mistaken for a possible hemangioma of the **liver** on CT scan. **Case report** A 37 year old man noticed a mass in the epigastric region since four months and dull non radiating pain in the epigastrium since a month and a half.

Low grade fibromyxoid sarcoma of the falciform ligament: a ...

Sep 24, 2003 · BACKGROUND: **Low grade fibromyxoid sarcomas (LGFMS)** are very rarely seen. They commonly arise from deep soft tissues of the lower extremities. Very few cases of intra-abdominal location have been reported. **CASE PRESENTATION:** We report a 37 year old man who presented with an abdominal mass and dragging pain.

Cited by: 56 Author: K Harish, AC Ashok, NK Alva
Publish Year: 2003

Huge mesenteric low-grade fibromyxoid sarcoma: A case ...

low-grade fibromyxoid sarcoma (LGFMS) or **Evans tumor** has been described in 1987 by Evans as an indolent soft tissue tumor and potentially metastasizing tumor. 1 This tumor has distinctive histopathologic findings with bland morphology and an aggressive behaviour. 2 The most common location of this **soft tissue tumor** is lower extremities of young patients, especially thigh.

Cited by: 3 Author: Bitu Geramizadeh, Zahra Zare, Amir Reza D...
Publish Year: 2018

Case Report: An extremely rare occurrence of recurrent ...

Introduction. **Low-grade fibromyxoid sarcoma (LGFMS)** is a rare sarcoma subtype, first described by Evans in 1987 1. The most common tumor locations are the deep soft tissue of extremities or trunks 2. Paratesticular LGFMS are rare with few cases published in the literature 3, 4. The etiology still unknown and the incidence is 0.18 per million, representing 0.6% of all soft tissue sarcomas 5.

LOW-GRADE FIBROMYXOID SARCOMA OF THE MAXILLA. ...

Jan 01, 2020 · **REPORT OF A CASE.** **Low-grade fibromyxoid sarcoma (LGFS)** is a variant of **fibrosarcoma** that commonly arises in the deep soft tissues of the lower extremities in young or middle-aged adults. It is characterized by high rates of local recurrence and late metastases, despite its ...

Author: René Martínez Flores, Bernardo Venega... Publish Year: 2020

Diffuse intra-abdominal low grade fibromyxoid sarcoma with ...

Jan 01, 2015 · **Low grade fibromyxoid sarcoma (LGFMS)** is a rare entity, first described by Evans in 1987 as a **malignant tumor** having a deceptively benign appearance , , . It is most commonly found in the **trunk and proximal extremities** but can occur almost anywhere in the body , .

Cited by: 8 Author: Jana Konecna, Gabriel Liberale, Johnny Ha...
Publish Year: 2015

Low-grade fibromyxoid sarcoma of the external anal ...

May 30, 2017 · **Low-grade fibromyxoid sarcoma (LGFMS)** is a rare soft tissue tumor that has a tendency to grow in the deep soft tissue of the **trunk** and extremities. Despite its benign appearance, the tumor has a high recurrence rate and metastatic potential. LGFMS in the perineal space is rare, and only a few cases have been reported.

Cited by: 2 Author: Lee-Kiat Ban, Ailun Heather Tseng, Shih-Hu...
Publish Year: 2017

Low-grade fibromyxoid sarcoma of the hand: a case report

Low-grade fibromyxoid sarcomas (LGFMS) are very rare soft tissue tumors. Evans first described the tumor in 1987 [1]. Most patients range in age from 25 to 46 years, but cases of this tumor in children have been reported [2–4]. The tumor typically occurs in the lower extremity, groin, head, and neck area [1, 3–5], but rare cases of LGFMS have been found in the chest wall, shoulder, arm [6], forearm [7], ...

Low Grade Fibromyxoid Sarcoma of the Neck: A Case Report

Low-grade fibromyxoid sarcoma (LGFMS) is a deceptively bland malignancy with potential for late recurrence and metastasis. It is a rare tumor but probably under-diagnosed because it is often confused with other entities. It is mainly localized in the deep soft ...

MRI and CT of Low-Grade Fibromyxoid Sarcoma in Children: A ...

Low-grade fibromyxoid sarcoma is a rare soft-tissue sarcoma characterized by a relatively benign histologic profile, with fibrous paucicellular and myxoid hypervascular zones . In the short term (< 5 years), **low-grade fibromyxoid sarcoma** has an indolent clinical behavior but also a tendency for late local recurrence and metastasis . It is more common in young and middle-aged adults but has increasingly ...

Low-grade fibromyxoid sarcoma

Rare Type

Low-grade fibromyxoid sarcoma is a rare type of low grade sarcoma first described by Harry Evans in 1987. It is characterized by a long and indolent clinical course and the possibility of local recurrence or distant metastases in a subset of patients. Unlike many other types of cancer, low grade fibromyxoid sarcoma can metastasize after many years, sometimes decades after the initial presentation of the tumor.

 Wikipedia

Data from: Wikipedia

[Suggest an edit](#)

Search Tools

[Turn off Hover Translation \(关闭取词\)](#)

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

162,000 Results

Any time ▾

[PDF] [Low-Grade Fibromyxoid Sarcoma: A Rare Distinctive Soft ...](#)

<https://www.bahrainmedicalbulletin.com/december...>

Low-grade fibromyxoid sarcoma (LGFMS) is a variant of fibrosarcoma with distinctive histopathological features. **Immunohistochemical staining (IHC)** is usually positive for vimentin while it is negative for a variety of antibodies such as **desmin, keratin, S100 protein, epithelial membrane antigen ...**

[Low grade fibromyxoid sarcoma of the falciform ligament: a ...](#)

europepmc.org/articles/PMC222986

Dec 01, 2002 · We **report** a 37-year old man with a giant **intra-abdominal low grade fibromyxoid sarcoma** measuring **21 cm** in size; which was mistaken for a possible hemangioma of the **liver** on CT scan. **Case report** A 37 year old man noticed a mass in the epigastric region since four months and dull non radiating pain in the epigastrium since a month and a half.

[Low grade fibromyxoid sarcoma of the falciform ligament: a ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/14507419>

Sep 24, 2003 · **BACKGROUND:** **Low grade fibromyxoid sarcomas (LGFMS)** are very rarely seen. They commonly arise from deep soft tissues of the lower extremities. Very few cases of intra-abdominal location have been **reported**. **CASE PRESENTATION:** We **report** a 37 year old man who presented with an abdominal mass and dragging pain.

Cited by: 56**Author:** K Harish, AC Ashok, NK Alva**Publish Year:** 2003

[Huge mesenteric low-grade fibromyxoid sarcoma: A case ...](#)

<https://journals.sagepub.com/doi/10.1177/2036361318777031>

low-grade fibromyxoid sarcoma (LGFMS) or **Evans tumor** has been described in 1987 by Evans as an indolent soft tissue tumor and potentially metastasizing tumor. 1 This tumor has distinctive histopathologic findings with bland morphology and an aggressive behaviour. 2 The most common location of this **soft tissue tumor** is lower extremities of young patients, especially thigh.

Cited by: 3**Author:** Bita Geramizadeh, Zahra Zare, Amir Reza D...**Publish Year:** 2018

Low-grade fibromyxoid sarcoma

Rare Type

Low-grade fibromyxoid sarcoma is a rare type of low grade sarcoma first described by Harry Evans in 1987. It is characterized by a long and indolent clinical course and the possibility of local recurrence or distant metastases in a subset of patients. Unlike many other types of cancer, low grade fibromyxoid sarcoma can metastasize after many years, sometimes decades after the initial presentation of the tumor.

Wikipedia

Data from: Wikipedia

[Suggest an edit](#)