

Microsoft Bing

国内版

国际版

Change in Neoadjuvant Chemotherapy could alter the Prognosis of

Sign in

Chat with Bing

ALLIMAGESVIDEOS

58,900 ResultsAny time

Chemotherapy Change Improves Pancreatic Cancer Outcomes ...

<https://www.cancer.gov/news-events/cancer-currents...>

Altering Chemotherapy Improves Outcomes in Early Stage Pancreatic Cancer. In stage IIA pancreatic cancer, the cancer has not spread and the tumor can be removed surgically. Results from two clinical trials are expected to improve the outlook for people diagnosed with early-stage pancreatic cancer that can be treated with surgery.

Advances of pathological complete response after ...

<https://journals.lww.com/pancreatology/Fulltext/...>

Pancreatic ductal adenocarcinoma (PDAC) is an aggressive malignancy with a poor prognosis. Only 15% to 20% of patients present with a primarily resectable tumor at the time of diagnosis. There has been an increasing interest in the use of neoadjuvant chemotherapy alone or combination with radiotherapy in patients with resectable, borderline resectable, and locally advanced pancreatic cancer.

PEOPLE ALSO ASK

Can chemotherapy shrink pancreatic cancer?

What is the prognosis for pancreatic cancer?

Search Tools

Turn off Hover Translation (关闭翻译)

Chemotherapy Change Improves Pancreatic Cancer Outcomes ...

<https://www.cancer.gov/news-events/cancer-currents...>

Altering Chemotherapy Improves Outcomes in Early-Stage Pancreatic Cancer. In stage IIA pancreatic cancer, the cancer has not spread and the tumor can be removed surgically. Results from two clinical trials are expected to improve the outlook for people diagnosed with early-stage pancreatic cancer that can be treated with surgery.

Neoadjuvant Chemo Ups Survival in Pancreatic Cancer

<https://www.medscape.com/viewarticle/908082>

Jan 22, 2019 - Neoadjuvant chemotherapy significantly improved overall survival in a cohort of Japanese patients and could be a new treatment option, but ...

Search Tools

Turn off Hover Translation (关闭取词)


PEOPLE ALSO ASK

- Can chemo be used to shrink pancreatic cancer?
- How can chemotherapy help early stage pancreatic cancer?
- Can neoadjuvant patients live longer after surgery?
- Can chemotherapy be given before surgery?

19 Feb 2021 08:00PM

1871 words • 4 matches • 2 sources

FAQ

 iThenticate®

61336-Review-check.docx

Quotes Included
Bibliography Included

7%
SIMILARITY

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 61336

Manuscript Type: CASE REPORT

Change in neoadjuvant chemotherapy could alter the prognosis of patients with pancreatic adenocarcinoma: A case report

Meyer A *et al.* Chemotherapy prognosis in Patients with PDAC

Alberto Meyer, Bárbara J Carvalho, Kayo A Medeiros, Leonardo Z Pipek, Fernanda Nascimento, Milena Suzuki, João Munhoz, Leandro R Iuamoto, Luiz A Carneiro-D'Albuquerque, Wellington Andraus

Abstract

Match Overview

1

Internet 74 words
crawled on 13-Mar-2020
www.futuremedicine.com

4%

2

Internet 57 words
crawled on 27-Aug-2020
www.wjgnet.com

3%

PAGE: 1 OF 8

Text-Only Report

国内版 国际版

Change in neoadjuvant chemotherapy could alter the prognosis of p



ALL

IMAGES

VIDEOS

60,400 Results

Any time ▾

Chemotherapy Change Improves Pancreatic Cancer ...

<https://www.cancer.gov/news-events/cancer-currents...> ▾

Altering Chemotherapy Improves Outcomes in Early-Stage Pancreatic Cancer. In stage IIA pancreatic cancer, the cancer has not spread and the tumor can be removed surgically. Results from two clinical trials are expected to improve the outlook for people diagnosed with early-stage pancreatic cancer that can be treated with surgery.

Neoadjuvant Chemo Ups Survival in Pancreatic Cancer

<https://www.medscape.com/viewarticle/908082> ▾

Jan 22, 2019 - Neoadjuvant chemotherapy significantly improved overall survival in a cohort of Japanese patients and could be a new treatment option, but the regimen is ...

PEOPLE ALSO ASK

Can chemotherapy shrink pancreatic cancer? ▾

What is the prognosis of exocrine pancreatic cancer? ▾

How long is chemo treatment for pancreatic cancer? ▾

Can pancreatic cancer be cured? ▾

Feedback

Complete Response after Treatment with Neoadjuvant ...

<https://www.hindawi.com/journals/crionm/2017/7834702> ▾

As recently shown, patients who undergo a prolonged time interval after neoadjuvant chemoradiation with ongoing chemotherapy are more likely to have an improved pathological response at time of surgical resection, which is associated with improved median overall survival; an apparent benefit observed in this case report (30-month survival) .

Cited by: 4

Author: Tiffany A. Pompa, William F. Morano, Ch...

Publish Year: 2017

A phase II trial proposal of total neoadjuvant treatment ...