

Endoscopic findings of dyspeptic patients unresponsive to ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5175035>

Dyspepsia is a group of symptoms including pain, bloating, early satiety, postprandial upper abdominal fullness, nausea, loss of appetite, pyrosis, regurgitation, and belching.

Dyspepsia is an important clinical problem in that it is very frequently seen in the community which is also one the reasons for seeking medical help.

Cited by: 1 Author: Mestan Sahin, Canan Akbulut, Can D...

Publish Year: 2015

Endoscopic Findings in Persistent Dyspepsia in Secondary ...

<https://www.ncbi.nlm.nih.gov/pubmed/21561680>

Search Toc

Turn off Hover

ALL IMAGES VIDEOS

Add the Olive with Bing extension

45,600 Results Any time

(PDF) Clinically Significant Endoscopic Findings in a ...

https://www.researchgate.net/publication/225296109... Jun 12, 2012 - clinically significant endoscopic findings in primary care patients with uninvestigated dyspepsia: the Canadian Adult Dyspepsia Empiric Treatment-Prompt Endoscopy (CADET-PE) study.

Clinically Significant Endoscopic Findings in a Multi ...

https://link.springer.com/article/10.1007/s10620-012-2256-7 - Jun 12, 2012 - A cross-sectional study of consecutive adult patients attending an open access endoscopy list for the primary indication of dyspepsia was conducted. Independent epidemiological and clinical factors for CSEF were determined prospectively. Cited by: 15 Author: Sanjiv Mahadeva, Khean-Lee Goh Publish Year: 2012

Cureus | Endoscopic Findings in Patients With ...

https://www.cureus.com/articles/43769-endoscopic-findings-in-patients-with... - Oct 26, 2020 - Clinically significant endoscopic findings were found in 91.8% of study subjects. The proportion of subjects with significant findings in the oesophagus, stomach, and duodenum were 73.5%, 33.3%, and 14%, respectively. On sub-analyzing this data, oesophagitis was found to be the most common oesophageal abnormality (33.1%). Author: Ahmad Badi, Vamanjore A. Naushad, N... Publish Year: 2020

(PDF) What Is the Prevalence of Clinically Significant ...

https://www.academia.edu/10466607/What_Is_the... - Prevalence of Clinically Significant Endoscopic Findings in Individuals With Dyspepsia According to Study Location The prevalence of erosive esophagitis was reported in 3 studies conducted in Asia, 8,18,20 and 4 studies in the West. 9,16,17,19 The pooled prevalence of erosive esophagitis was lower in Asian studies (2.7%, 95% CI, 1.2%-4.8%), compared with Western studies (25.0%, 95% CI, 4.0%-57.0%) ...

What Is the Prevalence of Clinically Significant ...

https://www.sciencedirect.com/science/article/pii/S154235651000563X - Oct 01, 2010 - Prevalence of Clinically Significant Endoscopic Findings in Individuals Regardless of Symptom Status. Six studies reported the prevalence of endoscopic findings in 4075 individuals, regardless of symptom status (Supplementary Table 2) 9, 18, 19, 20, 21, 22 The most common finding was erosive esophagitis, with a pooled prevalence of 9.9%. The pooled prevalence of peptic ulcer ... Cited by: 141 Author: Alexander C. Ford, Avantika Marwaha, Alle... Publish Year: 2010

Evaluation of the endoscopic findings in patients with ...

https://www.middleeastmedicalportal.com/evaluation... - Apr 14, 2020 - Endoscopy findings consisted of 54% antral gastritis, 20% peptic ulcer, 42% esophagitis (mostly includes grade A), 5% hiatal hernia, 5% gastric metaplasia, 15% prepyloric ulcer, 18% duodenal ulcer, 38% positive for H. pylori, and 1% Barrett's esophagus disease which had no significant relationship with dyspepsia.

1 2 3 4 5 >

Search Tools

Turn off Hover Translation (关闭翻译)

Name of Journal: *World Journal of Clinical Cases*
Manuscript NO: 62352
Manuscript Type: ORIGINAL ARTICLE

Observational Study
Clinically significant endoscopic findings in patients of dyspepsia with no warning symptoms: A cross-sectional study

Mao LQ *et al.* Role of endoscopy in dyspepsia

Li-Qi Mao, Shuang-Shuang Wang, Yan-Lin Zhou, Lin Chen, Lei-Min Yu, Meng Li, Bin Lv

Abstract

BACKGROUND

Dyspepsia is one of the commonest clinical disorder. However, controversy remains over the role of endoscopy in patients with dyspepsia. No studies have evaluated the

Match Overview

1	Crossref 105 words Jeffrey P. Fiorenza, Alex M. Timanow, Walter W. Chan. The Initial Management and Endoscopic Outcomes of Dy...	3%
2	Internet 59 words www.ncbi.nlm.nih.gov	2%
3	Crossref 57 words Chen, S. L., K. A. Grove, J. S. Lee, H. Miwa, H. Suzuki, P. Guo, Y. T. Hao, and M. H. Chen. "Systematic review with	2%
4	Internet 51 words crawled on 22-Sep-2020 www.wjnet.com	2%
5	Crossref 50 words "Yamada's Textbook of Gastroenterology", Wiley, 2015	2%
6	Crossref 49 words "Abstracts", Journal of Gastroenterology and Hepatology, 2016	2%
7	Crossref 48 words Khaled Aboljawad, Antonios Wehbeh, Emad Qayed. "Lo w Prevalence of Clinically Significant Endoscopic Find...	2%
8	Internet 35 words crawled on 15-Jan-2019 www.onlinelibrary.wiley.com	1%

国内版 国际版

Clinically significant endoscopic findings in patients of dyspepsia wi



ALL IMAGES VIDEOS

45,800 Results Any time ▾

(PDF) Clinically Significant Endoscopic Findings in a ...

<https://www.researchgate.net/publication/225296109...>

clinically significant endoscopic findings in primary care patients with uninvestigated dyspepsia: the Canadian Adult Dyspepsia Empiric Treatment-Prompt Endoscopy (CADET-PE) study.

Clinically Significant Endoscopic Findings in a Multi ...

<https://link.springer.com/article/10.1007/s10620-012-2256-7> ▾

Jun 12, 2012 · A cross-sectional study of consecutive adult patients attending an open access endoscopy list for the primary indication of dyspepsia was conducted. Independent epidemiological and clinical factors for CSEF were determined prospectively.

Cited by: 15 Author: Sanjiv Mahadeva, Khean-Lee Goh

Publish Year: 2012

(PDF) What Is the Prevalence of Clinically Significant ...

https://www.academia.edu/10466607/What_Is_the... ▾

Prevalence of Clinically Significant Endoscopic Findings in Individuals With Dyspepsia According to Study LocationThe prevalence of erosive esophagitis was reported in 3 studies conducted in Asia, 8, 18, 20 and 4 studies in the West. 9, 16, 17, 19 The pooled prevalence of erosive esophagitis was lower in Asian studies (2.7%; 95% CI, 1.2%-4.8%), compared with Western studies (25.0%; 95% CI, 4.0%-57.0%) ...

Cureus | Endoscopic Findings in Patients With ...

<https://www.cureus.com/articles/43769-endoscopic-findings-in-patients-with...> ▾

Clinically significant endoscopic findings were found in 91.8% of study subjects. The proportion of subjects with significant findings in the oesophagus, stomach, and duodenum were 73.5%, 33.3%, and 14%, respectively. On sub-analyzing this data, oesophagitis was found to be the most common oesophageal abnormality (33.1%).

Author: Ahmad Badi, Vamanjore A. Naushad, ... Publish Year: 2020

What Is the Prevalence of Clinically Significant ...

<https://www.sciencedirect.com/science/article/pii/S154235651000563X>

Oct 01, 2010 · Prevalence of Clinically Significant Endoscopic Findings in Individuals Regardless of Symptom Status. Six studies reported the prevalence of endoscopic findings in 4075 individuals,