

Match Overview

1	Internet 68 words crawled on 23-Jul-2020 pesquisa.bvsalud.org	2%
2	Internet 39 words crawled on 21-May-2020 journals.lww.com	1%
3	Internet 25 words crawled on 12-Oct-2020 cardiab.biomedcentral.com	1%
4	Internet 24 words crawled on 01-Feb-2020 jtd.amegroups.com	1%
5	Internet 17 words crawled on 03-Mar-2021 academic.oup.com	1%
6	Internet 14 words crawled on 25-Sep-2020 www.dovepress.com	<1%
7	Crossref Posted Content 9 words Qingyi Xu, Shuhua Jia, Maiko Fukasawa, Lin Lin et al. "A cross-sectional study on associations of physical symptoms, he	<1%
8	Crossref 8 words Siew Tzuh Tang, Chia-Hsun Hsieh, Ming-Chu Chiang, Jen-S hi Chen, Wen-Cheng Chang, Wen-Chi Chou, Ming-Mo Hou.	<1%

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 62923

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Self-perceived Burden and ¹influencing factors in patients with cervical cancer administered radiotherapy

Luo T *et al.* Burden in cervical cancer patients

Ting Luo, Rong-Zhi Xie, Yan-Xia Huang, Xiao-Hua Gong, Hui-Ying Qin, Yi-Xiao Wu

Abstract

BACKGROUND

Cervical cancer is the fourth commonest malignancy in women around the world. It represents the second most commonly diagnosed cancer in South East Asian women, and an important cancer death cause in women of developing nations. Data collected in

[PDF] Self-perceived Burden among Advanced Cancer Patients: A ...
www.davidpublisher.com/Public/uploads/Contribute/5f3e06f0734aa.pdf
Abstract: Objective: The aims of this study are to explore the prevalence and the influencing factors of SPB (self-perceived burden) among advanced cancer patients. Methods: One hundred fifty-six patients were recruited from the Oncology Comprehensive wards of a comprehensive hospital in Zhuhai city from January to December 2019.

Patients' self-reported factors influencing cervical ...
https://www.sciencedirect.com/science/article/pii/S235257892030062X
Aug 01, 2020 · 1. Introduction. Cervical cancer is the fourth most common neoplasm affecting women worldwide, with approximately 90% of cases occurring in developing countries (Ferlay et al., 2015). Cervical cancer is a leading contributor to the increased burden of diseases in less developed regions and it is acknowledged as the second most common cancer among women in developing ...
Author: Maryam Kasraeian, Kamran Hessami, H... Publish Year: 2020

Social, demographic and healthcare factors associated with ...
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4735146
Jan 21, 2016 · In spite of cervical cancer being a very common cancer, responsible for 2275 deaths every year (27.2 deaths per 100 000 women), women with cervical cancer in Uganda are an understudied population. 9 Against this background, we set out to examine patient and primary healthcare factors, and stage at diagnosis in women with cervical cancer in ...
Cited by: 25 Author: Amos Deogratus Mwaka, Christopher Orac...
Publish Year: 2016

Factors Associated With Higher Caregiver Burden Among ...
https://journals.lww.com/cancernursingonline/...
Over the past 2 decades, an increasing number of studies have been conducted to measure caregiver burden or to explore multidimensional factors associated with caregiver burden among family caregivers of cancer patients. 12–17 Unfortunately, studies focused on family caregivers of elderly cancer patients are still very rare.

Self-efficacy, social support, and quality of life among ...
https://www.emerald.com/insight/content/doi/10...
In Thailand, most patients with cancer primarily receive in-home care from their family members. However, information regarding the quality of life (QoL) of the primary family-member caregivers is scarce. The purpose of this paper is to assess primary family-member caregivers' QoL and its association with self-efficacy and social support using a Thai version of the Caregiver Quality of Life ...

Informal caregiver burden and influencing factors in ...
https://journals.sagepub.com/doi/full/10.1177/0300060520974927
Dec 01, 2020 · The sex of the informal caregiver, their relationship with the patient, daily duration of care, having chronic health problems and the patient's duration of disease are major factors influencing caregiver burden. Increased physical, mental and social attention and ...

Cervical Cancer Treatment Delays and Associated Factors in ...
https://ascopubs.org/doi/full/10.1200/JGO.18.00199
Jan 29, 2019 · Once they have their FSA, they are usually prioritized for treatment initiation, because several studies have shown that for every day of extension in overall treatment time, there is a loss of approximately 1% of local tumor control, particularly in patients with cervical cancer at stages III to IV. 45,46 However, because such patients often present with a poorer overall clinical condition, they are more likely to have unscheduled interruptions during radiotherapy...

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

TOOLS ▾

The Relationship between Self-Perceived Burden and ...

<https://www.hindawi.com/journals/bmri/2019/6840743> ▾



Abstract

Introduction

Methods

Results

Discussion

Co



At present, the influence factors of posttraumatic growth (PTG) in colorectal cancer (CRC) patients and the relationship between PTG, self-perceived burden (SPB), and resilience are not completely clear. Thus, the present study examined whether resilience and SPB could predict PTG in CRC patients. The role of resilience as a potential mediator was also assessed. Using a cross-sectional design, a convenience sample of 157 CRC patients was selected as subjects, from July to December 2016 in a third-grade hos...

[See more on hindawi.com](#)

Cited by: 4

Author: Chengshuai Zhang, Ruitong Gao, Jiandong ...

Publish Year: 2019

[PDF] Self-perceived Burden among Advanced Cancer Patients: A ...

www.davidpublisher.com/Public/uploads/Contribute/5f3e06f0734aa.pdf

Abstract: Objective: The aims of this study are to explore the prevalence and the **influencing factors** of SPB (self-perceived **burden**) among advanced **cancer patients**. Methods: One hundred fifty-six **patients** were recruited from the Oncology Comprehensive wards of a comprehensive hospital in Zhuhai city from January to December 2019.

PEOPLE ALSO ASK

What is the global burden of cervical cancer? ▾

What is the caregiver burden for elderly cancer patients? ▾

What are the risk factors for cervical cancer? ▾

What are the symptoms of advanced cervical cancer? ▾

Feedback

Human papillomavirus (HPV) and cervical ...

<https://www.who.int/news-room/fact-sheets/detail/...> ▾

Global **burden of cervical cancer** . Worldwide, **cervical cancer** is the fourth most frequent **cancer** in

