

Progressive tumefactive fibroinflammatory lesion of the ...

europepmc.org/articles/PMC3325739 ▾

Jan 01, 2012 · In 1 report on 3 cases, no details were provided on the prescribed radiation dose. 10 In two other cases, 50 Gy in 25 fractions was administered to disease located in the head and neck area. 5;9 In IPT of the base of skull, low dose radiation (20 Gy) proved to be ineffective with poor disease control reported in 6 patients. 23 In our case, it ...

Progressive Tumefactive Fibroinflammatory Lesion of the ...

<https://journals.sagepub.com/doi/10.4081/rt.2012.e12>

Tumefactive fibroinflammatory lesion (TFIL) is a rare benign tumor in the head and neck region. We present a case of a 40-year-old female with a benign but progressive lesion of the infratemporal f...

71,100 Results Any time ▾[Intrasellar Rhabdomyosarcoma: Case Report | Request PDF](#)<https://www.researchgate.net/publication/12061665...>

Request PDF | Intrasellar Rhabdomyosarcoma: Case Report | There has been only one reported case of an intrasellar rhabdomyosarcoma, the origin of which was in the para-nasal sinus. The authors ...

[Rhabdomyosarcoma:: Review for the Ophthalmologist ...](#)<https://www.researchgate.net/publication/10924947...>

Pleomorphic rhabdomyosarcomas is a rare variant, occurring more commonly in adults. We report a rare case of ocular pleomorphic rhabdomyosarcoma in an 18-month-old child with atypical presentation ...

[CT and MRI Features of Recurrent Tumors and Second Primary ...](#)<https://www.ajronline.org/doi/10.2214/ajr.181.3.1810879>

All rhabdomyosarcomas arose in the region previously irradiated, including five tumors that developed in the temporal muscle within the irradiated field and one that involved the contralateral temporal muscle,...

Cited by: 20 Author: Ukihide Tateishi, Tadashi Hasegawa, Kunihi...

Publish Year: 2003

[Neurological manifestations of rhabdomyosarcomas in the ...](#)<https://www.deepdyve.com/lp/springer-journals/...>

Case Report Case 1 Rhabdomyosarcomas are tumors of children. Those involving the middle ear, mastoid and nasopharynx may spread to the cranial cavity and cause neurological symptoms. Accordin...

[Extracranial primary and secondary meningiomas - ScienceDirect](#)<https://www.sciencedirect.com/science/article/pii/S0901502711014056>

Feb 01, 2012 · Extracranial meningiomas arising from the sinonasal tract or temporal muscle with no evidence of an association with an intracranial tumour (also called heterotopic, ectopic, or extracalvarial...

Cited by: 16 Author: Giorgio Iaconetta, Antonio Santella, Marco ...

Publish Year: 2012

[The dural tail sign—beyond meningioma - ScienceDirect](#)<https://www.sciencedirect.com/science/article/pii/S0009926004000741>

Feb 01 2005 · Meningioma is a benign tumour originating from arachnoid cells and frequently adheres to

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 64089

Manuscript Type: CASE REPORT

Adult rhabdomyosarcoma originating in the temporal muscle, invading the skull and meninges: A case report and literature review

Wang GH *et al.* Adult temporal muscle rhabdomyosarcoma

Geng-Huan Wang, He-Ping Shen, Zheng-Min Chu, Jian Shen

Abstract

BACKGROUND

Rhabdomyosarcoma (RMS) is a rare malignant tumor of mesenchymal origin that mainly affects children. Spindle cell/sclerosing rhabdomyosarcoma (SSRMS) is even rarer. As a new subtype, it was added to the latest disease classification of the World Health Organization in 2013. To our knowledge, this case is the first reported adult SSRMS originating from the temporal muscle.

Match Overview

1	Internet 20 words crawled on 05-Mar-2020 www.worldjscience.org	1%
2	Crossref 18 words Jinwei Zhang, Kaimi He, Jialing Wang, Wenming Xu, Lin Ma, Dejiang Huang. "A Prospective Evaluation of the Corneal"	1%
3	Internet 11 words crawled on 03-Nov-2020 www.wjnet.com	1%
4	Internet 11 words crawled on 26-Jan-2021 www.sipublishing.com	1%
5	Internet 11 words crawled on 25-Aug-2020 spublishing.blob.core.windows.net	1%
6	Crossref Posted Content 10 words Ke Zhao, Ke Rao, Xin Chen, Si Chen, Hailong Xu. "Primary Hepatic Cholangiocarcinoma in a Male Patient: A Case Report a	1%
7	Crossref 10 words Sarah N. Damont, Alexander J. Lazar, Julia A. Bridge, Robert S. Benjamin, Jonathan C. Trent. "PAX3/7-FOXO1 fusion st	1%

国内版 国际版

Adult rhabdomyosarcoma originating in the temporal muscle, invad



ALL IMAGES VIDEOS

70,100 Results Any time ▾

Intrasellar Rhabdomyosarcoma: Case Report | Request PDF

<https://www.researchgate.net/publication/12061665...>

Request PDF | Intrasellar Rhabdomyosarcoma: Case Report | There has been only one reported case of an intrasellar rhabdomyosarcoma, the origin of which was in the para-nasal sinus. The authors ...

Rhabdomyosarcoma:: Review for the Ophthalmologist ...

<https://www.researchgate.net/publication/10924947...>

Rhabdomyosarcoma is a malignant soft tissue neoplasm of skeletal muscle origin. Rhabdomyosarcoma is the ... adults. We report a rare case of ocular pleomorphic rhabdomyosarcoma in an 18-month-old ...

Soft tissue sarcoma - Symptoms and causes - Mayo Clinic

<https://www.mayoclinic.org/diseases-conditions/...>

Overview

Symptoms

Causes

Risk Factors

Soft tissue sarcoma is a rare type of cancer that begins in the tissues that connect, support and surround other body structures. This includes muscle, fat, blood vessels, nerves, tendons and the lining of your joints. More than 50 subtypes of soft tissue sarcoma exist. Some types are more likely to affect children, while others affect mostly adults. These tumors can be difficult to diagnose because they may be mistaken for many other types of growths. Soft tissue sarcoma can occur anywhere in...

[See more on mayoclinic.org](#)

Published: Dec 02, 2020

Rhabdomyosarcoma: A clinicopathological study and ...

<https://www.deepdyve.com/lp/wiley/rhabdomyosarcoma...>

Rhabdomyosarcoma: A clinicopathological study and classification of 39 cases

Rhabdomyosarcoma: A clinicopathological study and classification of 39 cases Horn, Robert C.; Enterline, Horatio T. 1958-01-01 00:00:00 has generally been considered an uncommon tumor. Its recognition has been complicated by the fact that many pathologists accept the diagnosis only when such evidences of a high degree ...

Extracranial primary and secondary meningiomas - ...