

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 62282

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Clinical pathological features and risk factors for primary breast cancer patients**

Yeyan Lei *et al.* risk factors for primary breast cancer patients

Ye-Yan Lei, Shuang Bai, Qing-Qing Chen, Xu-Jin Luo, Dong-Mei Li

### Match Overview

1 **Internet** 15 words  
crawled on 18-Apr-2021  
[www.hindawi.com](http://www.hindawi.com)

1%



ALL

IMAGES

VIDEOS

TOOLS ▾

## Risk factors for distant metastasis of patients with ...

<https://pubmed.ncbi.nlm.nih.gov/31113872>

Age > 50, higher clinical stage T and N, and tumor size > 5 cm were independent risk factors for distant metastasis of primary TNBC. Moreover, higher clinical stage T and stage N were independent risk factors for bone metastasis of the patients. TNBC patients with either bone or visceral metastasis have poor survival, with brain metastasis worst of all, though the OS difference was not statistically ...

Cited by: 14

Author: Yi Yao, Yuxin Chu, Bin Xu, Qinyong Hu, ...

Publish Year: 2019

## Clinical and pathological factors influencing ... - BMC Cancer

<https://bmccancer.biomedcentral.com/articles/10.1186/s12885-017-3969-y>

Jan 08, 2018 · In this uniquely large cohort of TNBC patients, we found that a high tumor stage at diagnosis – as defined by tumor size, pathological lymph nodes, and presence of metastasis – is the most important prognostic factor for cancer progression and mortality.

Cited by: 33

Author: Silvana Anna Maria Urru, Silvano Gallus, ...

Publish Year: 2018

## Clinical pathological characteristics of breast cancer ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4644204>

Epidemiological studies suggest that type 2 diabetes mellitus (T2DM) increases breast cancer risk and thereafter goes with an increased mortality. A study by Larsson et al. suggested that females with diabetes had a 20 % increased risk of postmenopausal breast cancer.

Cited by: 4

Author: Li Juanjuan, Wei Wen, Liu Zhongfen, Che...

Publish Year: 2015

## Risk factors for invasive breast cancer when core needle ...

<https://pubmed.ncbi.nlm.nih.gov/21079099>

Hypothesis: A core needle biopsy (CNB) diagnosis of ductal carcinoma in situ (DCIS) may be associated with a final diagnosis of invasive cancer. Preoperative radiologic, clinical, and pathological features may identify patients at high risk of diagnostic upstaging, who may be appropriate candidates for sentinel node biopsy at initial surgery.

Cited by: 97

Author: Emil D. Kurniawan, Allison Rose, Arlene ...

Publish Year: 2010

Clinical pathological features and risk factors for primary breast car



ALL IMAGES VIDEOS TOOLS

### Clinical and pathological factors influencing ... - BMC Cancer

<https://bmccancer.biomedcentral.com/articles/10.1186/s12885-017-3969-y>

Cited by: 35 Author: Silvana Anna Maria Urru, Silvano Gallus, Cris...

Publish Year: 2018

Jan 08, 2018 · In this uniquely large cohort of TNBC patients, we found that a high tumor stage at diagnosis – as defined by tumor size, **pathological** lymph nodes, and presence of metastasis – is the most important prognostic factor for cancer progression and mortality.

### Clinical and Pathologic Factors Matter in Decision Making ...

<https://www.jwatch.org/na53402/2021/04/09/clinical...>

Apr 09, 2021 · Clinicians caring for **patients** with early-stage **breast cancer** long relied on **clinical** and **pathologic factors** to estimate an individual's **risk** for recurrent disease and then acted on that estimate to make adjuvant therapy recommendations.

### Search Tools

Turn on Hover Translation (开启取词)

#### PEOPLE ALSO ASK

Which is a risk factor for breast cancer?

Is there a genetic predisposition for breast cancer?

What are the molecular aspects of breast cancer?

What are the signs and symptoms of breast cancer?



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

9,630,000 Results

Any time ▾

Open links in new tab



## Symptoms & Risk Factors - Breast Pathology | Johns Hopkins ...

<https://pathology.jhu.edu/breast/symptoms-risk-factors> ▾

There are a variety of genetic and environmental factors that can increase a person's risk for breast cancer. These factors include older age, female gender, personal or family history of breast cancer, known genetic mutations, reproductive and **hormonal** factors, and other environmental factors such as exposure to radiation. Hormonal Risk Factors

## Clinical and Pathological Characteristics of Patients with ...

<https://pubmed.ncbi.nlm.nih.gov/33870111>

Materials and methods: This study enrolled **patients** with invasive **breast cancer** who have been evaluated for BRCA1/2 mutations due to the presence of high-risk **factors** admitted to two tertiary referral centers in Turkey. **Clinical** and histopathological **features** were analyzed in BRCA1 mutation carriers, BRCA2 mutation carriers, and non-carriers.

### PEOPLE ALSO ASK

What are the factors that cause breast cancer? ▾

What is the pathologic stage of breast cancer? ▾

Is there a genetic predisposition for breast cancer? ▾

What are the molecular aspects of breast cancer? ▾

Feedback

## Clinical and Pathological Characteristics of Patients with ...

<https://europepmc.org/article/PMC/PMC8025729> ▾

Among **patients** diagnosed with invasive **breast cancer** between 2015 and 2020 in two tertiary referral centers in Istanbul, **patients** with **breast cancer** who have been evaluated for BRCA1/2 mutations due to the presence of high-risk **factors** including younger age at diagnosis (<40 years old), male sex, bilateral localization of the tumor, and ...

## Clinical and pathological factors influencing ... - BMC Cancer

<https://bmcancer.biomedcentral.com/articles/10.1186/s12995-017-2060-y>