







Rectal bleeding caused by a syphilitic inflammatory mass.

https://www.ncbi.nlm.nih.gov/pubmed/30696637

Jan 28, 2019 · A 47-year-old man presented with fatigue, decrease appetite, abdominal pain and rectal **bleeding**. His colonoscopy revealed a single, firm, raised, centrally ulcerated mass at the anorectal junction. During this same admission, he was diagnosed with HIV and **syphilis**, found to have multiple hepatic lesions and positive cerebrospinal fluidvenereal ...

Cited by: 2 Author: Joao Serigado, Eugene Lewis, Grace Kim

Publish Year: 2019

Syphilis Causing Lichenoid Esophagitis: ACG Case Reports ...

https://journals.lww.com/acgcr/Fulltext/2020/06000/...

Extragenital syphilis has been reported as either single case reports or small case series. 1 Syphilis of the stomach has been reported in only 1% of the cases. 2 Most cases of syphilis involving the GI tract have been reported in the rectum 3 and oral cavity. 4,5 ljiri et al in 2016 described syphilis of the whole GI tract, but they did not mention any involvement of the esophagus. 6 After extensive research, we found only few case reports that described esophageal syphilis ...



Feedback

Gastrointestinal bleeding caused by jejunal angiosarcoma ... https://pubmed.ncbi.nlm.nih.gov/33083419

Case summary: Here, we report the case of a 57-year-old man who presented with hematochezia for 1 mo. The patient had a medical history of chronic viral hepatitis B for 15 years. The causes of gastrointestinal bleeding were initially diagnosed as esophagogastric variceal bleeding or portal hypertensive gastropathy before endoscopy.