

[PDF] [Malignant fibrous histiocytoma arising in previous ...](#)<https://acsjournals.onlinelibrary.wiley.com/doi...>

lignant **fibrous histiocytoma** was diagnosed, and in December 1982, a revision of the **amputation stump** was done. A bulging multinodular **tumor** mass, measuring 5 X 4 X 3 cm, was found in the subcutaneous tissue. It extended up into the dermis and down to the deep fascia, which it did not penetrate. The cut surfaces were white to gray.

[Malignant fibrous histiocytoma of the mandible in the ...](#)<https://www.sciencedirect.com/science/article/pii/S0268000907000108>

May 01, 2008 · **Malignant fibrous histiocytoma** (MFH) is a pleomorphic high-grade tumor with histiocytic, fibroblastic and myofibroblastic features that was first introduced in the 1960s as a clinical entity . Several variants of MFH have been reported .

Cited by: 8

Author: Ali Hossein Mesgarzadeh, Ramin Mo...

Publish Year: 2008

[Malignant fibrous histiocytoma arising in previous ...](#)<https://www.deepdyve.com/lp/wiley/malignant...> ▾

Jan 01, 1984 · **Malignant fibrous histiocytoma** arising in previous **surgical** sites. Report of two **cases** Inoshita, Tsuyoshi; Youngberg, George A. 1984-01-01 00:00:00 Two patients who developed **malignant fibrous histiocytoma** in previous **surgical** sites are reported. One patient developed **malignant fibrous histiocytoma** in an **amputation** site, and the other patient ...

[Transformation of a giant cell tumor of bone into a ...](#)<https://link.springer.com/article/10.1007/s12570-013-0179-1> ▾

Apr 17, 2013 · We **report** a case of **GCT** with transformation into a secondary **malignant fibrous histiocytoma** that developed after local **GCT** recurrence and had no association to adjuvant radiation therapy. **Case report** A 59-year-old man underwent, in 1994, surgical curettage of a benign **GCT** located in the left distal femur followed by phenol cauterization and filling of **bony** defect with **bone** ...

Author: J. Bruns, O. Yazigee, G. Delling, ...

Publish Year: 2013

Malignant fibrous histiocytoma of the mandible in the ...<https://www.sciencedirect.com/science/article/pii/S0268000907000108>

May 01, 2008 · Introduction. Malignant fibrous histiocytoma (MFH) is a pleomorphic high-grade tumor with histiocytic, fibroblastic and myofibroblastic features that was first introduced in the 1960s as a clinical entity. Several variants of MFH have been reported. Although it has been hypothesized that these variants manifest common morphological patterns of a heterogeneous group of poorly differentiated ...

Cited by: 8

Author: Ali Hossein Mesgarzadeh, Ramin Mostofi Z...

Publish Year: 2008

Pleomorphic Sarcoma in a Patient with Osteopetrosis<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7536008>

Keywords: bone scintigraphy, humerus, malignant fibrous histiocytoma, osteopetrosis, undifferentiated pleomorphic sarcoma of bone CASE REPORT A 53-year-old woman with a past medical history of delayed development, hypothyroidism, and hyperlipidemia presented to the Emergency Department with shoulder pain and immobility.

Author: Kevin C McGill, Daria Motamedi, Nirma ...

Publish Year: 2020

Forequarter amputation for malignant tumours of the upper ...<https://europepmc.org/article/MED/19554191>

Jan 01, 2007 · Needle-core biopsy of the mass was suggestive of malignant fibrous histiocytoma (Figure 3). After a multidisciplinary discussion involving medical and radiation oncology, it was decided that the patient required surgical resection before radiation. He underwent forequarter amputation via an anterior approach (Figure 4). Surgical pathology ...

Cited by: 9

Author: Vanessa Dimas, Jennifer Kargel, John Bau...

Search Tools

Turn off Hover Translation (关闭取词)

国内版

国际版

Malignant fibrous histiocytoma of the bone in a traumatic amputat



ALL

IMAGES

VIDEOS

12,100 Results

Any time ▾

Malignant fibrous histiocytoma of the mandible in the ...

<https://www.sciencedirect.com/science/article/pii/S0268000907000108>

May 01, 2008 · Introduction. **Malignant fibrous histiocytoma** (MFH) is a pleomorphic high-grade tumor with histiocytic, fibroblastic and myofibroblastic features that was first introduced in the 1960s as a clinical entity. Several variants of MFH have been reported. Although it has been hypothesized that these variants manifest common morphological patterns of a heterogeneous group of poorly differentiated ...

Cited by: 8

Author: Ali Hossein Mesgarzadeh, Ramin Mostofi...

Publish Year: 2008

Forequarter amputation for malignant tumours of the upper ...

<https://europepmc.org/article/PMC/PMC2698809> ▾

Jan 01, 2007 · Needle-core biopsy of the mass was suggestive of **malignant fibrous histiocytoma** (Figure 3). After a multidisciplinary discussion involving medical and radiation oncology, it was decided that the patient required surgical resection before radiation. He underwent forequarter **amputation** via an anterior approach (Figure 4). Surgical pathology ...

Diaphyseal medullary stenosis-bone malignancy syndrome ...

<https://www.ncbi.nlm.nih.gov/medgen/350613> ▾

From OMIM Diaphyseal medullary stenosis with **malignant fibrous histiocytoma** is an autosomal dominant bone dysplasia characterized by pathologic fractures due to abnormal cortical growth and diaphyseal medullary stenosis. The fractures heal poorly, and there is progressive bowing of the lower extremities. In 2 families, affected individuals also showed a limb-girdle myopathy, with muscle ...

Pleomorphic Sarcoma in a Patient with Osteopetrosis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7536008>

Keywords: bone scintigraphy, humerus, **malignant fibrous histiocytoma**, osteopetrosis, undifferentiated pleomorphic sarcoma of bone CASE REPORT A 53-year-old woman with a past medical history of delayed development, hypothyroidism, and hyperlipidemia presented to the Emergency Department with shoulder pain and immobility.

Author: Kevin C McGill, Daria Motamedi, Nim...

Publish Year: 2020

Bone Sarcomas - Cancer Network

<https://www.cancernetwork.com/view/bone-sarcomas> ▾