(PDF) Adrenocortical Carcinoma: The Management of ...

https://www.researchgate.net/publication/262841950...



Case presentation: We report a 47-year-old female patient diagnosed with stage IV ACC with liver metastases, who developed symptoms of acute adrenocortical dysfunctions after hepatic arterial ...

Estimated Reading Time: 5 mins

Liver Resection and Ablation for Metastatic Adrenocortical ...

https://www.researchgate.net/publication/49818109...

Case presentation: We report a 47-year-old female patient diagnosed with stage IV ACC with liver metastases, who developed symptoms of acute adrenocortical dysfunctions after hepatic arterial ...

Pulmonary embolism after arterial chemoembolization for ...

https://www.researchgate.net/publication/271596272...

Background: Acute lung injury (ALI) is a rare but life-threatening pulmonary complication of transcatheter arterial chemoembolization (TACE) for hepatocellular carcinoma (HCC).

The incidence and outcome of major complication following ... https://journals.lww.com/md-journal/fulltext/2016/...

included. Major **complication** was defined as admission to a hospital for therapy, an unplanned increase in the level of care, prolonged hospitalization, permanent adverse sequelae, or death **after** conventional TAE/TACE therapy by Society of Interventional Radiology. During the study period, a total of 2863 TAE/TACE procedures were performed among 1120 patients, and a ...


Ruptured Hepatic Carcinoma After Transcatheter Arterial ...


https://www.researchgate.net/publication/256691744...

Ruptured Hepatic Carcinoma After Transcatheter Arterial Chemoembolization ... HCC after TACE is a rare complication. Objective The aim of this study was to ...

Estimated Reading Time: 5 mins

Prevention and management of infectious complications of





Rare complication of acute adrenocortical dysfunction in adrenoco







ALL

IMAGES

VIDEOS

2.290 Results

Any time ▼

Case Report: Complete Necrosis of a Large Adrenocortical ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8121102

Apr 30, 2021 · Introduction. Adrenocortical cancer (ACC) is a rare malignancy with mostly bad prognosis. Most cases are diagnosed late, in the presence of distant metastases (1–8). The medical treatment options for ACC are rather limited, as mitotane is the only available drug specific for the adrenal cortex and its use associated with severe side effects and a narrow therapeutic range ().

Author: Gergely Huszty, Attila Doros, Katalin ... Pu

Publish Year: 2021

Successful transcatheter arterial embolization for ...

https://www.sciencedirect.com/science/article/pii/S193004332100073X

Apr 01, 2021 · She underwent emergency transcatheter arterial embolization (TAE) to stabilize her general condition before undergoing elective surgery. To our knowledge, this is the first pediatric case of a ruptured ACT treated with TAE, which has only been reported so far in 2 adult patients with adrenocortical carcinoma and adenoma [3,4]. Case description

Author: Miki Yoshida, Hiroaki Takahashi, Yuni ... Publish Year: 2021

Successful transcatheter arterial embolization for ...

https://europepmc.org/article/PMC/PMC7900638 -

Feb 16, 2021 - Introduction. Pediatric adrenocortical tumor (ACT) is extremely rare and has an incidence of 0.1-0.4 cases per million individuals [1]. It should be treated as a malignant neoplasm, although histopathologically, it is difficult to classify it as a carcinoma or adenoma. Rupture is a rare complication of ACTs.

Author: Miki Yoshida, Hiroaki Takahashi, Yuni ... Publish Year: 2021

Case Report: Complete Necrosis of a Large Adrenocortical ...

https://www.frontiersin.org/articles/10.3389/fendo.2021.677187 -

Introduction

Case Presentation

Discussion and Conclusion

Dat >

Adrenocortical cancer (ACC) is a rare malignancy with mostly bad prognosis. Most cases are diagnosed late, in the presence of distant metastases (1–8). The medical treatment options for ACC are rather limited, as mitotane is the only available drug specific for the adrenal cortex and its use associated with severe side effects and a narrow therapeutic range (1). Systemic combination chemotherapy with