


## See results for Bile Duct Cancer (Medical Condition) A cancer that begins in the bile duct, a duct which carries bile and digestive fluid through liver. This causes jaundice, fever, abdominal pain, sudden w...

Search Tools Turn off Hover Translation (美团取调)



## Isolated IgG4-related sclerosing cholangitis misdiagnosed ...

https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/28202062

Feb 15, 2017. The most common cause of perihilar obstruction is cholangiocarcinoma, especially in Thailand. Benign perihilar stricture represents less than 20% of cases. IgG4-related disease and IgG4-related sclerosing cholangitis, however, have been receiving increased ...

Cited by: 12 Author: Narongsak Rungsakulkij, Pattana Sornm...

Publish Year: 2017

## Case report: A female case of isolated IgG4-related ...

https://pubmed.ncbi.nlm.nih.gov/28422840

Here we report a female case of isolated IgG4 related sclerosing cholangitis mimicking cholangiocarcinoma. Patient concerns: A 58-year-old woman complaint of one-month history of jaundice and right upper quadrant discomfort, and the biliary reconstruction showed full-length wall thickening and segmental stenosis.

Cited by: 5 Author: Jianchun Xiao, Guanqiao Li, Gang Yang, ...

Publish Year: 2017



Feedback

## Isolated IgG4-related sclerosing cholangitis misdiagnosed ... https://bmcsurg.biomedcentral.com/articles/10.1186/s12893-017-0214-1

Feb 15, 2017 · The most common cause of perihilar obstruction is cholangiocarcinoma, especially in Thailand. Benign perihilar stricture represents less than 20% of cases. IgG4-related disease and IgG4-related sclerosing cholangitis, however, have been receiving increased recognition. Isolated IgG4-