



Severe bleeding after sinus floor elevation using the ...

https://pubmed.ncbi.nlm.nih.gov/23000712

Severe bleeding after sinus floor elevation using the transcrestal technique: a case report Eur J Oral Implantol. Autumn 2012;5(3):287-91. Authors ... The patient reported to the emergency care unit a few hours after the surgery with ongoing bleeding and progressive swelling of especially the right side of the midface. The buccal swelling did ...

Cited by: 12 Author: Simon Storgard Jensen, Jacob Eriksen, ...

Publish Year: 2012

Uretero-aortic fistula: a case report

https://pubmed.ncbi.nlm.nih.gov/21318990

We report a case of uretero-aortic fistula following prolonged ureteral stenting in the left ureter which crossed over the aorta toward the cutaneous ureterostomy stoma. A 59 year-old woman presented massive bleeding from the left cutaneous ureterostomy of the single stoma for bilateral ureters. The ...

Cited by: 1 Author: Takeshi Nomoto, Yoshiyuki Tanaka, Kous...

Publish Year: 2009



Feedback

Massive upper gastrointestinal bleeding ... - BMJ Case Reports https://casereports.bmj.com/content/12/5/e228654 •

May 01, 2019 · A biliary **fistula** which may occur spontaneously or **after surgery**, is an abnormal communication from the biliary system to an organ, cavity or free surface. Spontaneous biliary-enteric **fistula** is a rare complication of gallbladder pathology, with over 90% of them secondary to cholelithiasis. Approximately 6% are due to perforating peptic ulcers. Symptoms of biliary-enteric **fistula** varies ...

Cited by: 2 Author: Dinesh Kumar Vadioaloo, Guo Hou Loo, ...