

Circa 1.380 risultati (0,72 secondi)

Metabolic encephalopathy in beta-ketothiolase deficiency: the ...

Mancanti: outside irreversible damage

Beta-Ketothiolase Deficiency Presenting with Metabolic Stroke ...

Mancanti: ~~irreversible~~ | Deve includere: irreversible

<https://www.researchgate.net> \ 31 . Traduci questa pagina

Cerebral White Matter

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3753195>

Lesions of the cerebral white matter (WM) result in focal neurobehavioral syndromes, neuropsychiatric phenomena, and dementia. The cerebral WM contains fiber pathways that convey axons linking cerebral cortical areas with each other and with subcortical structures, facilitating the distributed neural circuits that subserve sensorimotor function, intellect, and emotion.

Cited by: 403 Author: Jeremy D. Schmahmann, Eric E. Smith, Flor...
Publish Year: 2008

Search Tools

[Turn off Hover Translation](#)

Neuropsychiatric Aspects of Infectious Diseases ...

<https://www.sciencedirect.com/science/article/pii/S0749070408000432>

Oct 01, 2008 · Subcortical brain structures, such as the **basal ganglia** and periventricular white matter, are most affected. If unchecked, this immune cascade leads to neuronal cell apoptosis. Effective suppression of CNS viral replication and the resulting immune activation has the potential, however, to reverse at least some of the neuropathologic changes.

Cited by: 12 Author: Stephen J. Ferrando, Zachary Freyberg
Publish Year: 2008

Cerebral White Matter

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3753195>

Lesions of the cerebral white matter (WM) result in focal neurobehavioral syndromes, neuropsychiatric phenomena, and dementia. The cerebral WM contains fiber pathways that convey axons linking cerebral cortical areas with each other and with subcortical structures, facilitating the distributed neural circuits that subserve sensorimotor function, intellect, and emotion.

Cited by: 403 Author: Jeremy D. Schmahmann, Eric E. Smith, Flor...
Publish Year: 2008

Search Tools

[Turn off Hover Translation](#)

Neuropsychiatric Aspects of Infectious Diseases ...

<https://www.sciencedirect.com/science/article/pii/S0749070408000432>

Oct 01, 2008 · Subcortical brain structures, such as the **basal ganglia** and periventricular white matter, are most affected. If unchecked, this immune cascade leads to neuronal cell apoptosis. Effective suppression of CNS viral replication and the resulting immune activation has the potential, however, to reverse at least some of the neuropathologic changes.

Cited by: 12 Author: Stephen J. Ferrando, Zachary Freyberg
Publish Year: 2008

国内版

国际版

Emergence of lesions outside of the basal ganglia and irreversible



ALL

IMAGES

VIDEOS

11,800 Results

Any time ▾

Cerebral White Matter

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3753195>

Lesions of the cerebral white matter (WM) result in focal neurobehavioral syndromes, neuropsychiatric phenomena, and dementia. The cerebral WM contains fiber pathways that convey axons linking cerebral cortical areas with each other and with subcortical structures, facilitating the distributed neural circuits that subserve sensorimotor function, intellect, and emotion.

Cited by: 403

Author: Jeremy D. Schmahmann, Eric E. Smith, ...

Publish Year: 2008

Encephalitic Arboviruses: Emergence, Clinical Presentation ...

<https://link.springer.com/article/10.1007/s13311-016-0443-5> ▾

May 24, 2016 · Arboviruses are arthropod-borne viruses that exhibit worldwide distribution, contributing to systemic and neurologic infections in a variety of geographical locations. Arboviruses are transmitted to vertebral hosts during blood feedings by mosquitoes, ticks, biting flies, mites, and nits. While the majority of arboviral infections do not lead to neuroinvasive forms of disease, they are among ...

Cited by: 63

Author: Hamid Salimi, Matthew D. Cain, Robyn S...

Publish Year: 2016

Estimated Reading Time: 8 mins

Neuropsychiatric Aspects of Infectious Diseases ...

<https://www.sciencedirect.com/science/article/pii/S0749070408000432>

Oct 01, 2008 · Subcortical brain structures, such as the basal ganglia and periventricular white matter, are most affected. If unchecked, this immune cascade leads to neuronal cell apoptosis. Effective suppression of CNS viral replication and the resulting immune activation has the potential, however, to reverse at least some of the neuropathologic changes.

Cited by: 12

Author: Stephen J. Ferrando, Zachary Freyberg

Publish Year: 2008

Enzymes involved in branched-chain amino acid metabolism ...

<https://link.springer.com/article/10.1007/s00726-017-2412-7> ▾

Mar 21, 2017 · Brain MRI may show brain atrophy, occipital periventricular white matter lesions, and alterations of the basal ganglia, but could be normal (Seaver et al. 2011). Deficiency of 17-β-hydroxysteroid dehydrogenase type 10 leads to accumulation of its substrate, 2-methyl-3-hydroxybutyryl-coA.

Cited by: 69

Author: María M Adeva-Andany, Laura López-Mas...