

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 59246

**Manuscript Type:** CASE REPORT

**Primary aldosteronism due to bilateral micronodular hyperplasia and concomitant subclinical Cushing's syndrome: A case report**

Teragawa H *et al.* Primary aldosteronism and subclinical Cushing's syndrome

Hiroki Teragawa, Chikage Oshita, Yuichi Orita, Kunihiro Hashimoto, Hirofumi Nakayama, Yuto Yamazaki, Hironobu Sasano

#### Abstract

##### BACKGROUND

Adrenal incidentaloma (AI) has been frequently encountered in the clinical setting. It has been shown that primary aldosteronism (PA) or subclinical Cushing's syndrome (SCS) are the representative causative diseases of AI. However, the coexistence of PA and SCS has been reportedly observed. Recently, we encountered a case of AI, in which PA and SCS coexisted, confirmed by histopathological examinations after a laparoscopic

#### Match Overview

There are no matching sources for this report.

Primary aldosteronism due to bilateral micronodular hyperplasia an



ALL IMAGES VIDEOS

103,000 Results Any time

## Molecular Basis of Primary Aldosteronism and Adrenal ...

<https://academic.oup.com/jes/article/4/9/bvaa075/5864675>

Jun 29, 2020 · 2. Benign Adrenal Tumors Associated with **Primary Hyperaldosteronism**. The 2 most common causes of **primary hyperaldosteronism** (PA) are **aldosterone**-producing adenomas (APAs), also called Conn adenomas, or **bilateral** adrenal **hyperplasia** (BAH). APAs are usually unilateral, small in size (1-2 cm), and diagnosed in patients aged 40 to 50 years .

**Author:** Patricia Vaduva, Fideline Bonnet, Jérô... **Publish Year:** 2020

## REFERENCES - UpToDate

<https://www.uptodate.com/contents/cushings...>

A **case** of Adrenocorticotrophic hormone -independent **bilateral** adrenocortical macronodular **hyperplasia** **concomitant** with **primary aldosteronism**. BMC Surg 2017; 17:97. Doppman JL, Chrousos GP, Papanicolaou DA, et al. Adrenocorticotropin-independent macronodular adrenal **hyperplasia**: an uncommon cause of **primary** adrenal hypercortisolism.

Search Tools

Turn off Hover Translati

## Images of Primary Aldosteronism Due To Bilateral Micronodu...

<bing.com/images>





Primary aldosteronism due to bilateral micronodular hyperplasia an



Sign in



ALL

IMAGES

VIDEOS

121,000 Results

Any time ▾

## Molecular Basis of Primary Aldosteronism and Adrenal ...

<https://academic.oup.com/jes/article/4/9/bvaa075/5864675> ▾

Jun 29, 2020 · 2. Benign Adrenal Tumors Associated with **Primary Hyperaldosteronism**. The 2 most common causes of **primary hyperaldosteronism** (PA) are **aldosterone**-producing adenomas (APAs), also called Conn adenomas, or **bilateral** adrenal **hyperplasia** (BAH). APAs are usually unilateral, small in size (1-2 cm), and diagnosed in patients aged 40 to 50 years .

**Author:** Patricia Vaduva, Fideline Bonnet, Jérôm... **Publish Year:** 2020

## Diagnosis and treatment of primary aldosteronism ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/joim.12831>

Sep 25, 2018 · **Aldosterone**-producing adenoma (APA) and **bilateral** idiopathic **hyperaldosteronism** (IHA) are the two most common subtypes of PA (Table 1) 11. A less common form of PA, unilateral **hyperplasia** or **primary** adrenal **hyperplasia** (PAH), is caused by **micronodular** or macronodular **hyperplasia** of the zona glomerulosa of predominantly one adrenal gland.

**Cited by:** 61

**Author:** William Francis Young

**Publish Year:** 2019

## Search Tools

Turn off Hover Translation (关闭取词)

## Images of Primary Aldosteronism Due to Bilateral Micronodular Hy...

[bing.com/images](https://bing.com/images)

ALL IMAGES VIDEOS MAPS NEWS SHOPPING

120,000 Results

Any time ▾



## [Molecular Basis of Primary Aldosteronism and Adrenal ...](#)

<https://academic.oup.com/jes/article/4/9/bvaa075/5864675> ▾

Jun 29, 2020 · 2. Benign Adrenal Tumors Associated with **Primary Hyperaldosteronism**. The 2 most common causes of **primary hyperaldosteronism** (PA) are **aldosterone**-producing adenomas (APAs), also called Conn adenomas, or **bilateral** adrenal **hyperplasia** (BAH). APAs are usually unilateral, small in size (1-2 cm), and diagnosed in patients aged 40 to 50 years .

**Author:** Patricia Vaduva, Fideline Bonnet, Jérôm... **Publish Year:** 2020

## [Diagnosis and treatment of primary aldosteronism ...](#)

<https://onlinelibrary.wiley.com/doi/full/10.1111/joim.12831>

Sep 25, 2018 · **Aldosterone**-producing adenoma (APA) and **bilateral** idiopathic **hyperaldosteronism** (IHA) are the two most common subtypes of PA (Table 1) 11. A less common form of PA, unilateral **hyperplasia** or **primary** adrenal **hyperplasia** (PAH), is caused by **micronodular** or macronodular **hyperplasia** of the zona glomerulosa of predominantly one adrenal gland.

**Cited by:** 61 **Author:** William Francis Young

**Publish Year:** 2019

### PEOPLE ALSO ASK

How is primary aldosteronism diagnosed? ▾

What is unilateral adrenal hyperplasia? ▾

What is primary aldosteronism? ▾

Feedback

## [Cushing's syndrome due to primary bilateral macronodular ...](#)

<https://www.uptodate.com/contents/cushings-syndrome...> ▾