



All

Images

Videos

翻译成中文

关闭取词

105,000 Results

Any time ▾

Adjuvant chemotherapy for resected colorectal cancer ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4716056>

Jan 14, 2016 · Core tip: Surgical resection is the only option of cure for patients with metastatic colorectal cancer (CRC). The risk of recurrence within 18 mo after metastasectomy is about 75% and the liver is the main organ involved. However, there is still no standard treatment and the effective role of adjuvant therapy remains controversial.

Cited by: 18

Author: Brandi G, De Lorenzo S, Nannini M, Curti...

Publish Year: 2016

XELOX vs. FOLFOX in metastatic colorectal cancer: An ...

https://www.researchgate.net/publication/294105237_XELOX_vs_FOLFOX_in_metastatic...

Request PDF on ResearchGate | XELOX vs. FOLFOX in metastatic colorectal cancer: An updated meta-analysis | This meta-analysis aims to evaluate chemotherapy with XELOX (capecitabine plus ...

[PDF] Original Article The efficacy of XELOX and FOLFOX ...

www.ijcem.com/files/ijcem0034336.pdf

The efficacy of XELOX and FOLFOX adjuvant chemotherapy in stage III colorectal cancer patients with low preoperative serum albumin levels Po-Ming Chen^{1,2}, Chun-Yi Lin^{2,3}, ... In a large-scale meta-analysis, Landre et al. (2015) reported that the addition of oxaliplatin .

Efficacy of capecitabine versus 5-fluorouracil in ...

<https://academic.oup.com/annonc/article/22/12/2604/213377> ▾

Mar 17, 2011 · The consistency of these findings with the overall meta-analysis result suggests that the non-inferiority of capecitabine- versus 5-FU-containing chemotherapy can be extended to all patient subsets considered within the analysis. Most of the patients included in the meta-analysis had either colon or colorectal cancer (95%).

Cited by: 84

Author: J. Cassidy, L. Saltz, C. Twelves, E. Van C...

Publish Year: 2011

Author: Cassidy, J.

Three-month XELOX Regimen May Be Non-inferior to 6 ...

<https://www.cancertherapyadvisor.com/home/news/conference-coverage/american-society-of...> ▾

Jun 06, 2017 · An analysis suggested that a shorter, 3-month course of adjuvant chemotherapy for some patients with stage III colon cancer may yield comparable outcomes to the standard 6 ...

Author: Alexander M. Castellino, PhD

Therapeutic effect of endostar combined with chemotherapy ...

Name of Journal: *World Journal of Meta-Analysis*

Manuscript NO: 49553

Manuscript Type: META-ANALYSIS

The long-term efficacy of capecitabine plus oxaliplatin chemotherapy on stage III colon cancer: A meta-analysis

Fu HT *et al.* The XELOX chemotherapy for colon cancer

Hong-Tao Fu, Jing-Jing Tian, Ying-Ying Xu, Jia-Xin Fu, Shao-Ling Nie, Yan-Yan Tang, Ping Chen, Liang Zong

Abstract

Match Overview

1	Internet 221 words crawled on 28-Jul-2019 www.dovepress.com	5%
2	Internet 149 words crawled on 16-Dec-2018 www.jcancer.org	4%
3	Crossref 118 words C M Lund, D Nielsen, C Dehlendorf, A B Christiansen, F Rønholdt, J S Johansen, K K Vistisen. "Efficacy and toxic...	3%
4	Internet 99 words crawled on 16-Sep-2019 bmccancer.biomedcentral.com	2%
5	Internet 64 words crawled on 10-Jan-2017 springerplus.springeropen.com	2%
6	Internet 28 words crawled on 27-Jul-2019 www.ncbi.nlm.nih.gov	1%
7	Crossref 18 words Pectasides, Dimitrios, Vasilios Karavasilis, George Papaxoinis, Georgia Gourgioti, Thomas Makatsoris, Georgia	<1%
8	Crossref 17 words Jonathan M. Loree, Aaron Sha, Maryam Soleimani, Hagen F. Kennecke et al. "Survival Impact of CAPOX Versus ...	<1%
9	Crossref 15 words Matthew J Munro, Susrutha K Wickremesekera, Lifeng Peng, Swee T Tan, Tinte Itinteang. "Cancer stem cells in ...	<1%
10	Internet 15 words crawled on 25-Oct-2013 wehi.edu.au	<1%
11	Internet 13 words crawled on 01-Mar-2019 breast-cancer-research.biomedcentral.com	<1%



国内版

国际版

The long-term efficacy of capecitabine plus oxaliplatin chemo



All

Images

Videos

关闭取词

26,700 Results

Any time ▾

Capecitabine as Adjuvant Treatment for Stage III Colon Cancer

<https://www.nejm.org/doi/full/10.1056/NEJMoa043116>

Jun 30, 2005 · We randomly assigned a total of 1987 patients with resected stage III colon cancer to receive either oral capecitabine (1004 patients) or bolus fluorouracil plus leucovorin (Mayo Clinic regimen; 983 patients) over a period of 24 weeks.

Cited by: 1309

Author: Chris Twelves, Alfred Wong, Marek P. No...

Publish Year: 2005

Oxaliplatin-Based Adjuvant Chemotherapy for Stage III ...

<https://jamanetwork.com/journals/jamaoncology/fullarticle/2749687> ▾

Sep 12, 2019 · The ACHIEVE trial was an open-label, multicenter randomized phase 3, noninferiority trial conducted in Japan. Patients who had undergone potentially curative resection for stage III colon cancer were randomly assigned 1:1 to receive either 6 months or 3 months of adjuvant oxaliplatin-based chemotherapy (mFOLFOX6 or CAPOX).

Cited by: 1

Author: Takayuki Yoshino, Takeharu Yamanaka, ...

Publish Year: 2019

Author: Takayuki Yoshino, MD, Phd

Chemotherapy use in stage III colon cancer: a National ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4543852>

Methods: This is a retrospective study of stage III colon cancer patients (n = 207,718) diagnosed between 2000 and 2011 in the National Cancer Data Base (NCDB). The NCDB contains ~70% of new cancer diagnosis from >1500 American College of Surgeons accredited cancer programs in the United States and Puerto Rico.

Cited by: 7

Author: Smrity Upadhyay, Sumit Dahal, Vijaya R...

Publish Year: 2015

Effect of Adjuvant Capecitabine or Fluorouracil, with or ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4435787>

These data add to the evidence that oxaliplatin plus capecitabine or leucovorin and fluorouracil is the standard of care for the adjuvant treatment of stage III colon cancer, offering physicians flexibility to treat patients according to the patients' overall physical performance and preference.

Cited by: 82

Author: Hans-Joachim Schmoll, Chris Twelves, W...

Publish Year: 2014



All

Images

Videos

关闭取词

26,500 Results

Any time ▾

Capecitabine as Adjuvant Treatment for Stage III Colon Cancer

<https://www.nejm.org/doi/full/10.1056/NEJMoa043116>

Jun 30, 2005 · We randomly assigned a total of 1987 patients with **resected stage III colon cancer** to receive either **oral capecitabine** (1004 patients) or **bolus fluorouracil plus leucovorin** (Mayo Clinic regimen; 983 patients) over a period of 24 weeks.

Cited by: 1309

Author: Chris Twelves, Alfred Wong, Marek P. No...

Publish Year: 2005

Oxaliplatin-Based Adjuvant Chemotherapy for Stage III ...

<https://jamanetwork.com/journals/jamaoncology/fullarticle/2749687> ▾

Sep 12, 2019 · The ACHIEVE trial was an **open-label**, multicenter randomized phase 3, noninferiority trial conducted in Japan. Patients who had undergone potentially curative resection for **stage III colon cancer** were randomly assigned 1:1 to receive either 6 months or 3 months of **adjuvant oxaliplatin-based chemotherapy** (mFOLFOX6 or CAPOX).

Cited by: 1

Author: Takayuki Yoshino, Takeharu Yamanaka, ...

Publish Year: 2019

Author: Takayuki Yoshino, MD, PhD

Duration of Adjuvant Chemotherapy for Stage III Colon Cancer

<https://www.nejm.org/doi/full/10.1056/NEJMoa1713709>

Mar 29, 2018 · QUICK TAKE **Duration of Chemotherapy for Colon Cancer** 01:40. Since 2004, **oxaliplatin** with a fluoropyrimidine has been standard **adjuvant chemotherapy** in patients with **stage III colon cancer**.

Cited by: 120

Author: Axel Grothey, Alberto F. Sobrero, Anthon...

Publish Year: 2018

Capecitabine in the management of colorectal cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3097797>

Mar 24, 2011 · Díaz-Rubio E, Tabernero J, Gómez-España A, et al. Spanish Cooperative Group for the Treatment of Digestive Tumors Trial Phase III study of **capecitabine plus oxaliplatin** compared with **continuous-infusion fluorouracil plus oxaliplatin** as **first-line therapy** in **metastatic colorectal cancer**: final report of the Spanish Cooperative Group for the Treatment of Digestive Tumors Trial.

Cited by: 66

Author: Bradford R Hirsch, S Yousuf Zafar

Publish Year: 2011