

Name of Journal: *World Journal of Meta-Analysis*

Manuscript NO: 59547

Manuscript Type: META-ANALYSIS

Effects of antithrombotic agents on post-operative bleeding after endoscopic resection of gastrointestinal neoplasms and polyps: A systematic review and meta-analysis

Xiang BJ *et al.* Antithrombotic agents and ER

Bing-Jie Xiang, Yu-Hong Huang, Min Jiang, Cong Dai

Match Overview

1	Internet 286 words crawled on 06-Oct-2019 f6publishing.blob.core.windows.net	3%
2	Internet 145 words crawled on 18-Dec-2019 www.ueg.eu	2%
3	Crossref 105 words Jiaqi Dong, Kunyan Wei, Jiaqi Deng, Xi Zhou, Xiaomei Huang, MingMing Deng, Muhan Lü. "Effects of antithrombotic th ...	1%
4	Internet 40 words crawled on 20-Oct-2018 www.thieme-connect.de	<1%
5	Crossref 32 words Ki Bae Bang, Hyun Deok Shin. "Endoscopic treatment of s ... gery or procedure-related gastrointestinal bleeding", Gastro	<1%
6	Internet 32 words crawled on 18-Feb-2019 eprints.whiterose.ac.uk	<1%
7	Crossref 12 words "UEG Week 2016 Poster Presentations", United European G astroenterology Journal, 2016	<1%
8	Internet 11 words crawled on 12-Oct-2020 www.karger.com	<1%
9	Internet 11 words crawled on 02-May-2020 journals.sagepub.com	<1%
10	Internet 10 words crawled on 17-Aug-2020 mafiadoc.com	<1%
11	Internet 10 words crawled on 10-Sep-2018 www.thieme-connect.com	<1%



ALL

IMAGES

VIDEOS

12,200 Results

Any time ▾

[Effects of antithrombotic therapy on bleeding after ...](#)

<https://www.sciencedirect.com/science/article/pii/S0016510717321107>

Nov 01, 2017 · **Bleeding** is the most common adverse event **after endoscopic** submucosal dissection (ESD). Although several studies have reported on the use of **antithrombotic agents** and post-ESD **bleeding**, many issues remain controversial. We conducted a **meta-analysis** and **systematic review** to evaluate the **effects of antithrombotic** therapy on post-ESD **bleeding**.

Cited by: 14

Author: Jiaqi Dong, Kunyan Wei, Jiaqi Deng, Xi Z...

Publish Year: 2017

[Postoperative bleeding in patients on antithrombotic ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5558119>

Aug 14, 2017 · **METHODS**. A total of 2488 gastric **neoplasms** in 2148 consecutive patients treated between May 2001 and June 2016 were studied. The **antithrombotic agents** were categorized into antiplatelet **agents**, anticoagulants, and other **antithrombotic agents**, and we included combination therapies [e.g., dual antiplatelet therapy (DAPT)].The risk factors associated with post-ESD **bleeding**, namely, **antithrombotic** ...

Cited by: 13

Author: Chiko Sato, Kingo Hirasawa, Ryonho Koh...

Publish Year: 2017

[Bleeding after endoscopic submucosal dissection: Risk ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/27468187>

Jul 14, 2016 · **Core tip**: **Antithrombotic agents** and large **resection** are known to be significant risk factors for post-endoscopic submucosal dissection (post-ESD) **bleeding**, and as the indications for **antithrombotic agents** increase, and the indications for **endoscopic resection** are expanded, endoscopists have a chance to face an increasing number of patients with a high risk of post-ESD **bleeding**.

[Comparison of the effects of antithrombotic therapy on ...](#)

<https://www.sciencedirect.com/science/article/pii/S0016510718329869>

Feb 01, 2019 · **Endoscopic resection** (ER) is recognized as an effective treatment for superficial gastric neoplasm without metastasis because of the high en bloc **resection** rate and low local recurrence rate. 1 With regard to its safety and efficacy, the indications for ER have broadened in the field of therapeutic endoscopy. However, post-ER **bleeding** is one of the major adverse events **after** ER, and previous ...

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

60,200 Results

Any time ▼

Effects of antithrombotic therapy on bleeding after ...

<https://www.sciencedirect.com/science/article/pii/S0016510717321107>

Nov 01, 2017 · **Bleeding** is the most common adverse event **after endoscopic** submucosal dissection (ESD). Although several studies have reported on the use of **antithrombotic agents** and post-ESD **bleeding**, many issues remain controversial. We conducted a **meta-analysis** and **systematic review** to evaluate the **effects of antithrombotic** therapy on post-ESD **bleeding**.

Cited by: 14

Author: Jiaqi Dong, Kunyan Wei, Jiaqi Deng, Xi Z...

Publish Year: 2017

Postoperative bleeding in patients on antithrombotic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5558119>

Aug 14, 2017 · METHODS. A total of 2488 gastric **neoplasms** in 2148 consecutive patients treated between May 2001 and June 2016 were studied. The **antithrombotic agents** were categorized into antiplatelet **agents**, anticoagulants, and other **antithrombotic agents**, and we included combination therapies [e.g., dual antiplatelet therapy (DAPT)]. The risk factors associated with post-ESD **bleeding**, namely, **antithrombotic** ...

Cited by: 13

Author: Chiko Sato, Kingo Hirasawa, Ryonho Ko...

Publish Year: 2017

Bleeding after endoscopic submucosal dissection: Risk ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/27468187>

Jul 14, 2016 · Core tip: **Antithrombotic agents** and large **resection** are known to be significant risk factors for post-**endoscopic** submucosal dissection (post-ESD) **bleeding**, and as the indications for **antithrombotic agents** increase, and the indications for **endoscopic resection** are expanded, endoscopists have a chance to face an increasing number of patients with a high risk of post-ESD **bleeding**.

Comparison of the effects of antithrombotic therapy on ...

<https://www.sciencedirect.com/science/article/pii/S0016510718329869>

Feb 01, 2019 · **Endoscopic resection** (ER) is recognized as an effective treatment for superficial gastric neoplasm without metastasis because of the high en bloc **resection** rate and low local recurrence rate. 1

With regard to its safety and efficacy, the indications for ER have broadened in the field of therapeutic