

Wiley-Blackwell

- [illegible]

Received 15 January 2004; revised 10 March 2004; accepted 10 March 2004

© 2007 The Authors
Journal compilation © 2007 Blackwell Publishing Ltd

[illegible]

© 2004 Blackwell Publishing Ltd *Journal of Internal Medicine* 255: 111–118

Correlation between metastatic lymph node ratio and prognosis in patients with extrahepatic cholangiocarcinoma

Jian-Wei Zhang, Yun-Miao Chu, Zheng-Min Lai, Xiao-Ling Tang, Ying-Tai Chen, Chang-Feng Wang, Xu Chen

Fig. 1 *Phragmites* distribution in the coastal marshes of the Sacramento-San Joaquin River Delta, California, 1990–1999

Aims: To investigate the prognostic value of metastatic lymph node ratio (MLNR) in extrahepatic cholangiocarcinoma (ECC) patients undergoing radical resection.

METHODS Seventy-eight patients with BCC were enrolled. Correlations between various clinicopathological factors and prognosis were investigated by



Correlation between metastatic lymph node ratio and prognosis in patients with



学术搜索

找到约 2,530 条结果 (用时0.09秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[\[HTML\] Staging, resectability, and outcome in 225 patients with hilar cholangiocarcinoma](#)[nih.gov 中的 \[HTML\]](#)

WR Jarnagin, Y Fong, RP DeMatteo, M Gonen... - Annals of ..., 2001 - ncbi.nlm.nih.gov

... involvement by tumor; although useful to some extent, it does not correlate with resectability or ...

In addition, the correlation between more locally advanced tumors (ie, higher T stage) and ... Bacterial

colonization of bile at the time of resection correlated strongly with preoperatively ...

被引用次数: 887 相关文章 所有 7 个版本 引用 保存

[Lymph node ratio after curative surgery for intrahepatic cholangiocarcinoma](#)

D Tamandl, K Kaczirek, B Gruenberger... - British Journal of ..., 2009 - Wiley Online Library

... to evaluate the prognostic ability of LNR, the relationship between LNR and ... In contrast, the

UICC/AJCC classifications discriminate only between lymph node-positive and ... Number of lymph

node metastases is a significant prognostic factor in intrahepatic cholangiocarcinoma. ...

被引用次数: 44 相关文章 所有 5 个版本 引用 保存

[Adequate lymph node assessment for extrahepatic bile duct adenocarcinoma](#)

K Ito, H Ito, PJ Allen, M Gonen, D Klimstra... - Annals of ..., 2010 - journals.lww.com

... Skip Navigation [Links](#) Home > April 2010 - Volume 251 - Issue 4 > Adequate Lymph Node

Assessment ... the classification of hilar or distal was based on the relationship between the tumor ...

R0 resection and were classified as N0, a linear positive correlation between TLNC and ...

被引用次数: 55 相关文章 所有 5 个版本 引用 保存

[Evaluation of long-term survival after hepatic resection for metastatic colorectal cancer: a multifactorial model of 929 patients](#)

M Rees, PP Tekkis, FKS Welsh, T O'Rourke... - Annals of ..., 2008 - journals.lww.com

... Skip Navigation [Links](#) Home > January 2008 - Volume 247 - Issue 1 > Evaluation of Long-term ...

Background: No published studies investigated collectively the inter-relation of factors related

to ... resectable metastases, with variable 5-year survival rates ranging between 25% and ...

被引用次数: 470 相关文章 所有 4 个版本 引用 保存

[Intrahepatic cholangiocarcinoma: prognostic factors after surgical resection](#)

A Guglielmi, A Ruzzenente, T Campagnaro... - World journal of ..., 2009 - Springer

Further statistical analyses were carried out to identify the relation between macroscopic gross

[网页](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 229,000 条结果 (用时 0.24 秒)

Google 学术 : Correlation between metastatic lymph node ratio and prognosis in patients with extrahepatic cholangiocarcinoma

... outcome in 225 patients with hilar cholangiocarcinoma - Jarnagin - 被引用次数 : 888

... curative surgery for intrahepatic cholangiocarcinoma - Tamandl - 被引用次数 : 45

Adequate lymph node assessment for extrahepatic bile ... - Ito - 被引用次数 : 55

Prognostic significance of lymph node ratio after resection of ...www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)作者 : A Guglielmi - 2011 - 被引用次数 : 25 - [相关文章](#)

Lymph node (LN) metastases are a major negative prognostic factor for ... Five-year survival for patients above and below the LNR cut-off value of 0.25 was 0% ... originating from either intra- or extrahepatic bile.¹ Accordingly it is classified by ... for perihilar cholangiocarcinoma, the relationship between lymph nodes status ...

Get PDF (600K) - Wiley Online Libraryonlinelibrary.wiley.com/doi/10.1002/bjs.9752/pdf - [翻译此页](#)

patients undergoing surgery for distal cholangiocarcinoma. However, the value of these ... 2 Correlation between lymph node ratio and total lymph node count ...

Prognostic impact of lymph node metastasis - Wiley Online ...onlinelibrary.wiley.com > ... > Early View > Abstract - [翻译此页](#)

2015年1月22日 - All patients with distal cholangiocarcinoma who underwent surgical resection Correlation between lymph node ratio and total lymph node count, stratified The number of metastatic lymph nodes in extrahepatic bile duct ...

Assessment of nodal status for perihilar cholangiocarcinoma ...www.thehbsn.org > Home > Vol 2, No 5 (October 2013) ▾ [翻译此页](#)

Unfortunately the presence of lymph node metastases is frequent and it is ... in patients with positive LNs, with an expected 5-year survival between 0% and ...