

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 15108

Columns: ORIGINAL ARTICLES

Basic Study

Novel isolated cecal pouch model for endoscopic observation in rats

Kurodo Koshino, Nobuo Kanai, Masayuki Yamato, Teruo Okano, Masakazu

Yamamoto

Match Overview

1 **CrossCheck** 26 words
Yano, Kazuo, Kenichiro Tsuyuki, Natsumi Watanabe, Hiroshi Kasanuki, and Masayuki Yamato. "The regulation of all ...

1%

[网页](#) [图片](#) [视频](#) [新闻](#) [更多](#) [搜索工具](#)

找到约 1,590,000 条结果 (用时 0.61 秒)

Google 学术: Novel isolated cecal pouch model for endoscopic observation in rats

... of robenacoxib: a novel selective inhibitor of ... - King - 被引用次数: 36

Treatment of diversion colitis with short-chain-fatty acid ... - Harig - 被引用次数: 591

Prophylaxis of pouchitis onset with probiotic therapy: a ... - Gionchetti - 被引用次数: 882

Stricture of ileo-anal J pouch anastomosis increased the ... www.researchgate.net/.../231224610_Stricture_of_ileo-anal_J_... - 翻译此页

The abnormal endoscopic findings included isolated pouch ulcer (n = 29, ... the intensity of pouchitis in an experimental rat model ... ic fecal residue in the pouch in patients and experimental ... has been observed that endoscopic dilatation of the stric- come after restorative proctocolectomy in pigs: comparing a novel

Probiotics and prebiotics in chronic inflammatory bowel ... www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: JB Ewaschuk - 2006 - 被引用次数: 132 - 相关文章

2006年10月7日 - Patients with a recent ileo-anal pouch anastomosis who responded to probiotic However, no difference was observed in endoscopic disease activity. ... The ingested probiotic bacteria are not detectable in the cecal content, yet the isolated from goat milk reduce intestinal inflammation in a rat model of ...

[PDF] Intestinal stasis associated bowel inflammation - World ... www.wjgnet.com/1007-9327/5/518.pdf - 翻译此页

作者: S Komatsu - 被引用次数: 5 - 相关文章

1999年9月22日 - by intestinal stasis and/or an altered fecal stream, resulting from "cul de sac" structures (blind loop or pouch) in the intestinal tract, bowel obstruction

网页 图片 购物 视频 新闻 更多 ▾ 搜索工具

找到约 1,560,000 条结果 (用时 0.50 秒)

Google 学术 : Novel isolated cecal pouch model for endoscopic observation in rats

... of robenacoxib: a novel selective inhibitor of ... - King - 被引用次数 : 36

Treatment of diversion colitis with short-chain-fatty acid ... - Harig - 被引用次数 : 591

Prophylaxis of pouchitis onset with probiotic therapy: a ... - Gionchetti - 被引用次数 : 883

Stricture of ileo-anal J pouch anastomosis increased the ...

www.researchgate.net/.../231224610_Stricture_of_ileo-anal_J_... - 翻译此页

The abnormal endoscopic findings included isolated pouch ulcer (n = 29, ... the intensity of pouchitis in an experimental rat model ... ic fecal residue in the pouch in patients and experimental ... has been observed that endoscopic dilatation of the stricture ... come after restorative proctocolectomy in pigs: comparing a novel

Probiotics and prebiotics in chronic inflammatory bowel ...

[www.ncbi.nlm.nih.gov/.../Literature/PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/.../Literature/PubMed%20Central%20(PMC)) - 翻译此页

作者 : JB Ewaschuk - 2006 - 被引用次数 : 132 - 相关文章

2006年10月7日 - Patients with a recent ileo-anal pouch anastomosis who responded to probiotic However, no difference was observed in endoscopic disease activity. ... The ingested probiotic bacteria are not detectable in the cecal content, yet the isolated from goat milk reduce intestinal inflammation in a rat model of ...

[PDF] Intestinal stasis associated bowel inflammation - World ...

www.wjgnet.com/1007-9327/5/518.pdf - 翻译此页

作者 : S Komatsu - 被引用次数 : 5 - 相关文章

1999年9月22日 - by intestinal stasis and/or an altered fecal stream, resulting from "cul de sac" structures (blind loop or pouch) in the intestinal tract, bowel obstruction or ... novel insights into the molecular and cellular basis ... finding are specific and endoscopic observation is ... has been presented in experimental models of.