


15428-Review

BY ZHONG-XIN LI

Quotes Excluded
Bibliography Excluded0%
SIMILAR

Match Overview



Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 15428

Columns: ORIGINAL ARTICLE

Basic Study

Hemodynamics and vasoactive substance levels during renal congestion

Zhong-Xin Li, Man-Cai Wang, You-Cheng Zhang, Jie Mao, Mo Chen, Rui Ni, Feng-Xian Wei, Gen-Nian Wang, Ling-Yi Zhang

Abstract

AIM: To explore hemodynamics and vasoactive substance levels during renal vein

There are no matching sources for this document.

[网页](#) [图片](#) [新闻](#) [视频](#) [更多](#) [搜索工具](#)

找到约 270,000 条结果 (用时 0.51 秒)

Google 学术: Hemodynamics and vasoactive substance levels during renal congestion

... capillary wedge pressure in the intensive care unit - Fortia - 被引用次数: 176

Role of humoral factors in the intestinal hyperemia ... - Benoit - 被引用次数: 227

Vasoactive factors and hemodynamic mechanisms in ... - Gatta - 被引用次数: 54

Deleterious Effects of Increased Intra-Abdominal Pressure ...

www.hindawi.com/journals/an/2014/731657/ • [翻译此页](#)

作者: Z Armaly - 2014

2014年10月9日 - Most ADHF hospitalizations stem from congestion in patients refractory to ... The importance of locally released vasoactive substances in the regulation of ... Moreover, kidney function at both the renal hemodynamic level and ...

[PDF] Deleterious Effects of Increased Intra-Abdominal Pressur...

downloads.hindawi.com/journals/an/2014/731657.pdf • [翻译此页](#)

作者: Z Armaly - 2014

2014年10月9日 - kidney excretory function and renal hemodynamics are not fully understood. The current review ... axis in CHF is arginine vasopressin (AVP), whose circulating levels are ... Most ADHF hospitalizations stem from congestion in patients ... activation and release of vasoactive substances resulting in low renal ...

Relationship Between B-Type Natriuretic Peptides and ...

content.onlinejacc.org/article.aspx?articleid=1136577 • [翻译此页](#)作者: PR Fortia - 2005 - 被引用次数: 173 - [相关文章](#)

2005年5月17日 - Hemodynamics were recorded simultaneously with blood sampling for BNP and NT-proBNP. Peptide levels were approximately four-fold greater in

网页 图片 新闻 视频 购物 更多 搜索工具

找到约 264,000 条结果 (用时 0.48 秒)

Google 学术 : Hemodynamics and vasoactive substance levels during renal congestion

... capillary wedge pressure in the intensive care unit - Forfia - 被引用次数 : 176

Role of humoral factors in the intestinal hyperemia ... - Benoit - 被引用次数 : 227

Vasoactive factors and hemodynamic mechanisms in ... - Gatta - 被引用次数 : 54

Deleterious Effects of Increased Intra-Abdominal Pressure ...

www.hindawi.com/journals/an/2014/731657/ ▾ 翻译此页

作者 : Z Armaly - 2014 - 相关文章

2014年10月9日 - Most ADHF hospitalizations stem from congestion in patients refractory to The importance of locally released vasoactive substances in the regulation of Moreover, kidney function at both the renal hemodynamic level and ...

[PDF] Deleterious Effects of Increased Intra-Abdominal Pressur...

downloads.hindawi.com/journals/an/2014/731657.pdf ▾ 翻译此页

作者 : Z Armaly - 2014

2014年10月9日 - kidney excretory function and renal hemodynamics are not fully understood. The current review axis in CHF is arginine vasopressin (AVP), whose circulating levels are ... Most ADHF hospitalizations stem from congestion in patients ... activation and release of vasoactive substances resulting in low renal ...

Relationship Between B-Type Natriuretic Peptides and ...

content.onlinejacc.org/article.aspx?articleid=1136577 ▾ 翻译此页

作者 : PR Forfia - 2005 - 被引用次数 : 173 - 相关文章

2005年5月17日 - Hemodynamics were recorded simultaneously with blood sampling for BNP and NT-proBNP. ... Peptide levels were approximately four-fold greater in patients with the presence of renal failure, independent of pulmonary congestion Alterations in circulating vasoactive substances in the critically ill—a ...