

Match Overview

[x] [y]

1	CrossCheck (17 words) The first part of the manuscript is the first, the second part is the second, and the third part is the third.	1%
2	Publishers (12 words) The first part of the manuscript is the first, the second part is the second, and the third part is the third.	1%
3	Publishers (12 words) The first part of the manuscript is the first, the second part is the second, and the third part is the third.	1%
4	CrossCheck (12 words) The first part of the manuscript is the first, the second part is the second, and the third part is the third.	1%
5	CrossCheck (12 words) The first part of the manuscript is the first, the second part is the second, and the third part is the third.	1%

Name of journal: World Journal of Gastroenterology

ESPJ Manuscript NO: 15261

Column: CASE REPORT

Esophageal subepithelial lesion diagnosed as malignant gastrointestinal neuroectodermal tumor

Sang Bum Kim, Si Hyung Lee, Mi Jin Gu

Abstract

A 25-year-old male visited our hospital with a complaint of aggravating dysphagia and odynophagia for a few days. Esophagogastroduodenoscopy showed huge bulging mucosa with intact surface causing luminal narrowing at 35cm from incisor teeth. Endoscopic ultrasonography showed about 35 mm sized irregular margined in-homogeneous hypoechoic lesion with obscure layer of origin. Endoscopic



Esophageal subepithelial lesion diagnosed as malignant gastrointestinal neuro



学术搜索

找到约 274 条结果 (用时0.06秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Gastrointestinal neuroendocrine/neuroectodermal tumors

CM Fenoglio-Preiser - American Journal of Clinical ..., 2001 - ajcp.ascpjournals.org

... **Esophagus** Too few to estimate <0.4%-0.5% <1% of **esophageal** neoplasms in ... **esophageal** carcinoid **tumors** display various architectural patterns, including organoid, trabecular, and sheet ... 81.52 This suggests that the **tumors** arise from **subepithelial** neuroendocrine complexes ...

被引用次数: 31 相关文章 所有 5 个版本 引用 保存

171.67.112.51 中的 [PDF]

Synovial sarcoma of the upper digestive tract: a report of two cases with demonstration of the X; 18 translocation by fluorescence in situ hybridization

SD Billings, LF Meisner, OW Cummings, E Tejada - Modern Pathology, 2000 - nature.com

... A, low power view of the gastric **tumor** demonstrating extension of the **tumor** into the superficial **subepithelial** stroma. ... **Esophageal** carcinosarcoma: a genetic analysis. ... Carcinosarcoma of the **esophagus**: an immunohistochemical and electron microscopic study. ...

被引用次数: 44 相关文章 所有 5 个版本 引用 保存

nature.com 中的 [HTML]

A retroperitoneal bronchogenic cyst with malignant change

SM Sullivan, S Okada, M Kudo... - Pathology ..., 1999 - Wiley Online Library

... **Esophageal** cyst, another foregut-derived **lesion**, can be readily ruled out, because it has two distinct layers of smooth ... Further, the possibility of Müllerian cyst is also remote because it lacks **subepithelial** basement membrane and its mucin is negative for PAS staining. ...

被引用次数: 81 相关文章 所有 5 个版本 引用 保存

[HTML] **Granular cell tumors of the gastrointestinal tract: Questions and answers**

F Radaelli, G Minoli - Gastroenterology & hepatology, 2009 - ncbi.nlm.nih.gov

... Granular cell **tumor** of the **esophagus**: three case reports and review of the literature. Turk J Gastroenterol. ... A prospective study comparing endoscopy and EUS in the evaluation of GI **subepithelial** masses. ... Endosonographic features of **esophageal** granular cell **tumors**. ...

被引用次数: 4 相关文章 所有 2 个版本 引用 保存

nih.gov 中的 [HTML]

Immunohistochemical localization of ganglioside GD3 in human malignant melanoma, epithelial tumors, and normal tissues

WG Donald, MB Dizon, A Kaurth... - Cancer research, 1985 - AACR

aacrjournals.org 中的 [PDF]

网页 图片 新闻 视频 购物 更多 搜索工具

找到约 34,700 条结果 (用时 0.20 秒)

Google 学术 : Esophageal subepithelial lesion diagnosed as malignant gastrointestinal neuroectodermal tumor

Gastrointestinal neuroendocrine/neuroectodermal ... - Fenoglio-Preiser - 被引用次数 : 31

Synovial sarcoma of the upper digestive tract: a report ... - Billings - 被引用次数 : 44

A retroperitoneal bronchogenic cyst with malignant ... - Sullivan - 被引用次数 : 81

[PDF] MAGIC 2015 Poster Presentation List - Multi-national Allia...

www.magicSYMPOSIUM.org/abstract/2015.../2015_poster.pdf 翻译此页

A Case of Esophageal Subepithelial Lesion Diagnosed as Malignant Gastrointestinal Neuroectodermal Tumor. Byung Ik Jang. Yeungnam University. Korea.

Granular Cell Tumors of the Gastrointestinal Tract

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) 翻译此页

作者 : F Radaelli - 2009 - 被引用次数 : 4 - 相关文章

Granular cell tumors (GCTs) are rare, usually benign tumors, that can be located ...

esophageal tumors.³ Involvement in other gastrointestinal localizations such as ...

reliable for detecting the nature and origin of a subepithelial mass, and GCTs are ... In

a review of 183 cases, 8 lesions (4%) were malignant and all of these 8 ...

제64회 대한소화기내시경학회 추계학술대회 일시 : 2014년 ...

www.gie.or.kr/event/meeting/old_data/2014.../2014_64.html 翻译此页

Role of endoscopy in the diagnosis of upper GI SET, 이항락 한양의대 14, Efficacy

of Endoscopic Incisional Biopsy on Gastric Sub-Epithelial Tumor to Diagnosis and

Treatment, Jun Bong Kim 12, A Case of Esophageal Subepithelial Lesion

Diagnosed as Malignant Gastrointestinal Neuroectodermal Tumor, Jay Song.

Endoscopic mucosal resection for granular cell tumor ...

www.jghc.org > Home > Vol. 10 翻译此页