

## Match Overview

[icon] [icon]

1	Copied/Plagiarized: 17 words Context: 2.8 similarity, 100% match, 100% similarity, 100% match, 100% similarity, 100% match	1%
2	Copied/Plagiarized: 17 words Context: 2.8 similarity, 100% match, 100% similarity, 100% match, 100% similarity, 100% match	1%

Name of Journal: World Journal of Gastroenterology

ENP's Manuscript NO: 15846

Column: LETTERS TO THE EDITOR

**Transjugular intrahepatic portosystemic shunt as bridge-to-surgery in refractory gastric antral vascular ectasia**

Aymen Ben, Viridiana Otero, Aurélie Plesner, Patrice Vallon, Xavier Drey

**Abstract**

Gastric antral vascular ectasia (GAVE) may cause gastrointestinal bleeding (GIB). The treatment of GAVE relies on endoscopic approaches such as electrocoagulation (argon plasma coagulation, laser therapy, heater probe therapy, radiofrequency ablation), cryotherapy, and band ligation. In refractory cases, antrectomy may be considered. In the event of an associated cirrhosis and portal hypertension, it has been suggested that antrectomy could be an option, provided the mortality risk isn't considered too great.



Transjugular intrahepatic portosystemic shunt as bridge-to-surgery in refi



网页

图片

视频

新闻

更多 ▾

搜索工具

找到约 20,600 条结果 (用时 0.30 秒)

### The Transjugular Intrahepatic Portosystemic Shunt in the ...

[www.hindawi.com/journals/ijh/2012/167868/](http://www.hindawi.com/journals/ijh/2012/167868/) ▾ 翻译此页

作者: G Pomier-Layrargues - 2012 - 被引用次数: 14 - 相关文章

2012年5月18日 - The **transjugular intrahepatic portosystemic shunt** (TIPS) was used for the first ..... **Gastric Antral Vascular Ectasia** (GAVE) .... It has no role in these patients except for highly selected cases as a **bridge** to liver transplantation. .... In **refractory** cases, **surgical** side-to-side portacaval shunt has been used in the ...

### Peripheral Vascular Interventions - 第 457 页 - Google 图书结果

<https://books.google.com.hk/books?isbn=0781786878> - 翻译此页

Krishna Kandarpa - 2008 - Medical

Portal hypertensive gastropathy and **gastric** 107. **antral vascular ectasia** (GAVE) ....

**Transjugular intrahepatic portosystemic shunts**: comparison with ... The North American Study for the Treatment of **Refractory** Ascites. .... **Transjugular intrahepatic portosystemic shunt** for Budd-Chiari syndrome after failure of **surgical** shunting.

### Transjugular intrahepatic portosystemic shunts and portal ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: S Siramolpiwat - 2014 - 相关文章

2014年12月7日 - In patients with **refractory** ascites, there is growing evidence that TIPS not only ... **Transjugular intrahepatic portosystemic shunt** (TIPS) is a percutaneously ... changes similar to those observed in **surgical** portosystemic shunts. .... **Gastric antral vascular ectasia** (GAVE), also known as watermelon stomach, ...

### World J Gastroenterol - World Journal of Gastroenterology

[www.wjgnet.com/1007-9327/7/170.asp](http://www.wjgnet.com/1007-9327/7/170.asp) ▾ 翻译此页

作者: M Schepke - 被引用次数: 7 - 相关文章

This **transjugular portosystemic intrahepatic** stent **shunt** (TIPS) functions as a side to

[网页](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 20,600 条结果 (用时 0.71 秒)

### The Transjugular Intrahepatic Portosystemic Shunt in the ...

[www.hindawi.com/journals/ijh/2012/167868/](http://www.hindawi.com/journals/ijh/2012/167868/) ▾ [翻译此页](#)

作者 : G Pomier-Layrargues - 2012 - 被引用次数 : 14 - [相关文章](#)

2012年5月18日 - The **transjugular intrahepatic portosystemic shunt** (TIPS) was used for the first ..... **Gastric Antral Vascular Ectasia** (GAVE) ... It has no role in these patients except for highly selected cases as a **bridge** to liver transplantation. ... In **refractory** cases, **surgical** side-to-side portacaval shunt has been used in the ...

### Peripheral Vascular Interventions - 第 457 页 - Google 图书结果

<https://books.google.co.uk/books?isbn=0781786878> - [翻译此页](#)

Krishna Kandarpa - 2008 - Medical

Portal hypertensive gastropathy and **gastric** 107. **antral vascular ectasia** (GAVE) ... **Transjugular intrahepatic portosystemic shunts**: comparison with ... The North American Study for the Treatment of **Refractory** Ascites. ... **Transjugular intrahepatic portosystemic shunt** for Budd-Chiari syndrome after failure of **surgical** shunting.

### Transjugular intrahepatic portosystemic shunts and portal ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者 : S Siramolpiwat - 2014 - [相关文章](#)

2014年12月7日 - In patients with **refractory** ascites, there is growing evidence that TIPS not only ... **Transjugular intrahepatic portosystemic shunt** (TIPS) is a percutaneously ... changes similar to those observed in **surgical** portosystemic shunts. .... **Gastric antral vascular ectasia** (GAVE), also known as watermelon stomach, ...

### World J Gastroenterol - World Journal of Gastroenterology

[www.wjgnet.com/1007-9327/7/170.asp](http://www.wjgnet.com/1007-9327/7/170.asp) ▾ [翻译此页](#)

作者 : M Schepke - 被引用次数 : 7 - [相关文章](#)

This **transjugular portosystemic intrahepatic stent shunt** (TIPS) functions as a side to side **shunt**, ... of liver cirrhosis, namely recurrent intestinal bleeding and **refractory** ascites. ... TIPS insertion does not require abdominal **surgery** and has a lower .... hypertensive gastropathy and patients with a **gastric (antral) vascular ectasia** ...