iThenticate

18174-Review

3%

Match Overview

1 2%

2 1%

3 1%

4 <1%

5 <1%

Name of journal: World Journal of Gastroenterology
ESPS Manuscript NO: 18174
Columns: EDITORIAL

Sphincter of oddi dysfunction Type III: New studies suggest new approaches are needed

C Mel Wilcox

Abstract

Sphincter of Oddi dysfunction has been classified into three types based upon the presence or absence of objective findings including liver test abnormalities and bile duct dilatation. Type III is the most controversial and is classified as biliary type pain in the absence of any these objective findings. Many prior studies have shown that the clinical response to endoscopic therapy is higher based upon the presence of these objective criteria. However, there has been variable correlation of the manometry findings to

Google Sphincter of oddi dysfunction Type III: New studies suggest new approa

网页 图片 新闻 购物 视频 更多 搜索工具

找到约 434,000 条结果 （用时 0.64 秒）

[PDF] Sphincter of Oddi Dysfunction - Johns Hopkins Medicine
www.hopkinsmedicine.org/.../**sphincter_of_oddi_dysfunction**.p... 翻译此页
There are two types of sphincter of Oddi dysfunction: 1) papillary stenosis and 2) ... on the sphincter have suggested that there is a multifactor control mechanism of the ... patients with type II SOD dysfunction but more studies are needed before ... Medical therapy for sphincter of Oddi dysfunction is an attractive approach ...

6. Sphincter of Oddi dysfunction - GastroHep eBook
www.gastrohep.com/ebooks/ebook.asp?book=1405120789... 翻译此页
News and Views ... Sphincter of Oddi dysfunction (SOD) refers to a motor abnormality of the ... Therapy for SOD is discussed, using an evidence-based approach. .... recent evidence suggests that the presence of abnormal liver tests in Type II biliary SOD ..... Further studies are needed before recommending this technique.

Sphincter of Oddi Dysfunction - iffgd.org
www.iffgd.org/site/gi-**disorder**s/functional-gi-**disorder**s/sod 翻译此页
2014年2月6日 - Pain caused by sphincter of Oddi dysfunction may occur after gall bladder ... News & Events ... About half of these patients will have findings on laboratory studies or ... ultrasound, CT scan, or MRCP) to suggest a definite abnormality, ... doctors can categorize these patients as having SOD Types I and II.

Sphincter of Oddi and its Dysfunction
www.ncbi.nlm.nih.gov › ... › PubMed Central (PMC) 翻译此页
作者：P Seetharam - 2008 - 被引用次数：5 - 相关文章
[1] The abnormalities of SO contractility (sphincter of Oddi dysfunction: SOD) .... 55% and 28% for patients with types I, II and III suspected SOD, respectively. ... Studies have also suggested that SOM may be highly sensitive for ... of an endoscopic approach for SOD is that it mainly addresses the biliary type. ... Recent Activity

Google di dysfunction Type III: New studies suggest new approaches are needed

网页 图片 新闻 购物 视频 更多 搜索工具

找到约 429,000 条结果（用时 0.70 秒）

[PDF] Sphincter of Oddi Dysfunction - Johns Hopkins Medicine
www.hopkinsmedicine.org/.../**sphincter_of_oddi_dysfunction**.p... 翻译此页
There are two types of sphincter of Oddi dysfunction: 1) papillary stenosis and 2) ... on the sphincter have suggested that there is a multifactor control mechanism of the ... patients with type II SOD dysfunction but more studies are needed before ... Medical therapy for sphincter of Oddi dysfunction is an attractive approach ...

Sphincter of Oddi Dysfunction - iffgd.org
www.iffgd.org/site/gi-**disorder**s/functional-gi-**disorder**s/sod 翻译此页
2014年2月6日 - Pain caused by sphincter of Oddi dysfunction may occur after gall bladder ... News & Events ... About half of these patients will have findings on laboratory studies or ... ultrasound, CT scan, or MRCP) to suggest a definite abnormality, ... doctors can categorize these patients as having SOD Types I and II.

6. Sphincter of Oddi dysfunction - GastroHep eBook
www.gastrohep.com/ebooks/ebook.asp?book=1405120789... 翻译此页
News and Views ... Sphincter of Oddi dysfunction (SOD) refers to a motor abnormality of the ... Therapy for SOD is discussed, using an evidence-based approach. .... recent evidence suggests that the presence of abnormal liver tests in Type II biliary SOD ..... Further studies are needed before recommending this technique.

Sphincter of Oddi and its Dysfunction
www.ncbi.nlm.nih.gov › ... › PubMed Central (PMC) 翻译此页
作者：P Seetharam - 2008 - 被引用次数：5 - 相关文章
[1] The abnormalities of SO contractility (sphincter of Oddi dysfunction: SOD) may .... Studies have also suggested that SOM may be highly sensitive for ... SOM is necessary to predict the outcome in biliary type II SOD ... of an endoscopic approach for SOD is that it mainly addresses the biliary type of SOD. ..... Recent Activity.