

## Match Overview

[Icon] [Icon]

- |   |   |     |
|---|---|-----|
| 1 | CrossCheck 11/2014<br>Title: Metastatic Pancreatic Cancer in Young Patients   | 1%  |
| 2 | CrossCheck 12/2014<br>Title: Role of PD-1 Inhibitors in the Treatment of Metastatic Pancreatic Cancer: A Review of the Literature and Future Perspectives | <1% |
| 3 | CrossCheck 11/2014<br>Title: Role of PD-1 Inhibitors in the Treatment of Metastatic Pancreatic Cancer: A Review of the Literature and Future Perspectives | <1% |

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 15779

Columns: CASE REPORT

**Metastatic pancreatic carcinoma with neoadjuvant FOLFIRINOX therapy and R0 resection.**

Sophie Schuster, Peter Kropf, Joannin Ramez, Gerald Antoch, Wilfried Trudel, Koenig, Dieter Häussinger, Dirk Graf

### Abstract

Patients with metastatic carcinoma of the pancreas have a very poor prognosis, and long-term survival cannot be expected. This case report describes two patients with an initial diagnosis of metastatic pancreatic cancer, both with hepatic metastases and one with an additional peritoneal carcinomatosis. Initially, both patients were treated intravenously with the FOLFIRINOX chemotherapy regimen.



Metastasized pancreatic carcinoma with neoadjuvant FOLFIRINOX therapy



网页

新闻

图片

视频

更多

搜索工具

找到约 11,500 条结果 (用时 0.76 秒)

### Google 学术: Metastasized pancreatic carcinoma with neoadjuvant FOLFIRINOX therapy and R0 resection

... versus gemcitabine for metastatic pancreatic cancer - Conroy - 被引用次数: 1175

... locally advanced pancreatic adenocarcinoma - Hosein - 被引用次数: 65

FOLFIRINOX in locally advanced pancreatic cancer: ... - Faris - 被引用次数: 41

### Neoadjuvant FOLFIRINOX application in borderline ...

[www.ncbi.nlm.nih.gov/pubmed/25501072](http://www.ncbi.nlm.nih.gov/pubmed/25501072) - 翻译此页

作者: A Paniccia - 2014 - 相关文章

Neoadjuvant FOLFIRINOX application in borderline resectable pancreatic adenocarcinoma: a retrospective cohort study. Paniccia A(1) ... The primary endpoints were completion of therapy to surgery and negative resection margin (R0) rate.

### Neoadjuvant treatment of borderline resectable and non ...

[annonc.oxfordjournals.org/content/early/.../annonc.mdt239.full](http://annonc.oxfordjournals.org/content/early/.../annonc.mdt239.full) - 翻译此页

作者: V Heinemann - 2013 - 被引用次数: 22 - 相关文章

2013年7月12日 - In non-metastatic, non-resectable pancreatic cancer, treatment ... R0 resection after neoadjuvant FOLFIRINOX was achieved in 5 of 18 (28%) ...

### Treatment of locoregional disease: adjuvant versus ...

[annonc.oxfordjournals.org/content/23/suppl\\_10/x141.full](http://annonc.oxfordjournals.org/content/23/suppl_10/x141.full) - 翻译此页

作者: V Heinemann - 2012 - 被引用次数: 2 - 相关文章

Disadvantages of neoadjuvant therapy are tumor progression during ... after the R0 resection of PC prolongs disease-free and OS. adjuvant treatment should be started ... FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer.



eatatic carcinoma with neoadjuvant FOLFIRINOX therapy and R0 resection



网页 新闻 购物 图片 视频 更多 ▾ 搜索工具

找到约 11,400 条结果 (用时 0.62 秒)

### Google 学术 : Metastasized pancreatic carcinoma with neoadjuvant FOLFIRINOX therapy and R0 resection

... versus gemcitabine for metastatic pancreatic cancer - Conroy - 被引用次数 : 1180

... locally advanced pancreatic adenocarcinoma - Hosein - 被引用次数 : 65

FOLFIRINOX in locally advanced pancreatic cancer: ... - Faris - 被引用次数 : 41

### Neoadjuvant FOLFIRINOX application in borderline ...

[www.ncbi.nlm.nih.gov/pubmed/25501072](http://www.ncbi.nlm.nih.gov/pubmed/25501072) - 翻译此页

作者 : A Paniccia - 2014 - 相关文章

Neoadjuvant FOLFIRINOX application in borderline resectable pancreatic adenocarcinoma: a retrospective cohort study. Paniccia A(1) ... The primary endpoints were completion of therapy to surgery and negative resection margin (R0) rate.

### Neoadjuvant treatment of borderline resectable and non ...

[annonc.oxfordjournals.org/content/early/.../annonc.mdt239.full](http://annonc.oxfordjournals.org/content/early/.../annonc.mdt239.full) ▾ 翻译此页

作者 : V Heinemann - 2013 - 被引用次数 : 22 - 相关文章

2013年7月12日 - In non-metastatic, non-resectable pancreatic cancer, treatment .....

R0 resection after neoadjuvant FOLFIRINOX was achieved in 5 of 18 (28%) ...

### Treatment of locoregional disease: adjuvant versus ...

[annonc.oxfordjournals.org/content/23/suppl\\_10/x141.full](http://annonc.oxfordjournals.org/content/23/suppl_10/x141.full) ▾ 翻译此页

作者 : V Heinemann - 2012 - 被引用次数 : 2 - 相关文章

Tumors are either confined to the pancreas (T1 and T2) or extend beyond the ... of pancreatic adenocarcinoma is mandatory if neoadjuvant treatment is planned [7]. ... Disadvantages of neoadjuvant therapy are tumor progression during ... the R0 resection of PC prolongs disease-free and OS. adjuvant treatment should be ...